

Pacific Journal of Mathematics

**CONVOLUTION TRANSFORMS WHOSE INVERSION
FUNCTION HAS COMPLEX ROOTS IN A WIDE ANGLE**

ZEEV DITZIAN

CONVOLUTION TRANSFORMS WHOSE INVERSION FUNCTION HAS COMPLEX ROOTS IN A WIDE ANGLE

ZEEV DITZIAN

In this paper a class of convolution transforms:

$$(1.1) \quad f(x) = \int_{-\infty}^{\infty} G(x-t)\varphi(t)dt$$

whose kernels $G(t)$ satisfy

$$(1.2) \quad G(t) = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} [E(s)]^{-1} \cdot e^{st} ds$$

where

$$(1.3) \quad E(s) = \prod_{k=1}^{\infty} (1 - s/a_k) \quad \text{or} \quad E(s) = \prod_{k=1}^{\infty} (1 - s/a_k) \exp(s \operatorname{Re} a_k^{-1})$$

will be treated. Investigation of properties will be carried for the subclass defined by the restriction on a_k as follows:

(a) For some ψ , $0 < \psi < \pi/2$

$$\min_{n=0,1,2} |n\pi - \arg a_k| \leq \psi \quad \text{where} \quad a_k \neq 0.$$

(b) For some $0 < q < 1$

$$\text{and integer } l \quad |a_{k+l}| \geq q^{-l} |a_k| \quad \text{for all } k \geq k_0.$$

It should be mentioned that the restriction (a) on the argument of a_k is much weaker than those used in other subclasses of convolution transforms defined by (1.1), (1.2) and (1.3) that were investigated before.

I. I. Hirschman and D. V. Widder [4] treated a class of transforms for which $\arg a_k$ tend to either 0 or π . J. Dauns and D. V. Widder [1] and the author [2] studied the case $E(s) = \prod_{k=1}^{\infty} (1 - s^2/a_k^2)$ for which $|\arg a_k| \leq \psi < \pi/4$, that is: The sequence of roots contains pairs of a_k and $-a_k$. A milder way of coupling was introduced by the author [3]. The question that arises is: Can we relax the restriction on the argument of the a_k 's and still have the transforms and their inversion formulae? Of course it was shown [1, p. 442] that in some simple cases the analogous inversion formula to that of Hirschman and Widder does not hold. Examples can be given to show that in some cases (1.2) does not converge. Here a restriction on the growth of the roots is given (b) which assures us of the convergence of (1.2) and helps us to prove that

$$(1.4) \quad \lim_{m \rightarrow \infty} P_m(D)f(x) = \prod_{k=1}^{\infty} (1 - a_k^{-1}D)f(x) = \varphi(x) \quad \text{a.e.}$$

where

$$D \equiv \frac{d}{dx}.$$

We shall assume for convenience that $|a_k| \leq |a_{k+1}|$ and also that $E(s) = \prod_{k=1}^{\infty} (1 - s/a_k)$ since treating $E(s) = \prod_{k=1}^{\infty} (1 - s/a_k) \exp(s \operatorname{Re} a_k^{-1})$ would mean only shifting the argument t of $G(t)$ by the number $\sum \operatorname{Re} a_k^{-1}$.

It should be noted that the harder part of the proof is an estimate of $E(s)$ (§ 2) and an estimate on $|G_m(t)| = |P_m(D)G(t)|$ (in § 4) the results achieved for the later had not been published for the nonvoid intersection of our class and the class of variation diminishing transforms.

2. An estimate for $E(s)$.

THEOREM 2.1. *Suppose that $E(s) = \prod_{k=1}^{\infty} (1 - s/a_k)$ and the sequence $\{a_k\}$ satisfies conditions (a) and (b) and let $0 < \eta < \pi/2 - \psi$, then there exist $A(n) > 0$ and $B(n) > 0$ such that*

$$(2.1) \quad |E(re^{i\theta})| \geq (A(n) + B(n)r^{2n})^{1/2}$$

for any n and r uniformly for $\psi + \eta \leq \theta \leq \pi - \psi - \eta$ and $\pi + \psi + \eta \leq \theta \leq 2\pi - \psi - \eta$.

Proof. Without loss of generality we may assume $|a_k| \leq |a_{k+1}|$. We define $\varphi_k = \arg a_k$ and have

$$\begin{aligned} |1 - re^{i\theta}/|a_k|e^{i\varphi_k}|^2 &= 1 - 2\frac{r}{|a_k|} \cos(\theta - \varphi_k) + \frac{r^2}{|a_k|^2} \\ &\geq 1 - 2\frac{r}{|a_k|} \cos \eta + \frac{r^2}{|a_k|^2} = \left[1 - \left(1 + \frac{1}{2} \tan^2 \eta\right)^{-1}\right] \\ (2.2) \quad &+ \left[\left(1 + \frac{1}{2} \tan^2 \eta\right)^{-1} - 2\frac{r}{|a_k|} \cos \eta + \frac{r^2}{|a_k|^2} \cos^2 \eta \left(1 + \frac{1}{2} \tan^2 \eta\right)\right] \\ &+ \frac{1}{2} \frac{r^2}{|a_k|^2} \sin^2 \eta \geq \frac{1}{2} \cdot \frac{\tan^2 \eta}{1 + \frac{1}{2} \tan^2 \eta} + \frac{1}{2} \frac{r^2}{|a_k|^2} \sin^2 \eta. \end{aligned}$$

Therefore

$$\prod_{k=1}^n \left| 1 - r e^{i\theta}/a_k \right|^2 \geq \left(\frac{\frac{1}{2} \tan^2 \eta}{1 + \frac{1}{2} \tan^2 \eta} \right)^n + \left(\frac{\sin^2 \eta}{2} \right)^n \frac{r^{2n}}{|a_1|^2 \dots |a_n|^2} \equiv A_1(n) + B_1(n)r^{2n} .$$

To complete the proof it is enough to show that there exists a constant $c > 0$ independent of r and θ (in its specified angle) such that:

$$\left| \prod_{k=1}^{\infty} (1 - r e^{i\theta}/a_k) \right|^2 \geq c > 0 .$$

We shall write

$$\prod_{k=n+1}^{\infty} \dots = \left(\prod_{k=n+1}^{n_1(r)} \dots \right) \cdot \left(\prod_{k=n_1(r)+1}^{n_2(r)} \dots \right) \cdot \left(\prod_{k=n_2(r)+1}^{\infty} \dots \right) \equiv I_1(r) \cdot I_2(r) \cdot I_3(r) .$$

Choose $n_1(r)$ as the largest integer satisfying

$$k \leq n_1(r) , \quad |a_k| < r/2 \cos \eta .$$

If $n_1(r) < n + 1$ then $I_1(r) = 1$; otherwise

$$I_1(r) = \left| \prod_{k=n+1}^{n_1(r)} (1 - r e^{i\theta}/a_k) \right|^2 \geq \prod_{k=n+1}^{n_1(r)} \left(1 - \frac{2r}{|a_k|} \cos \eta + \frac{r^2}{|a_k|^2} \right) \geq 1 .$$

We choose $n_2(r)$ as $n_2(r) = \min \{l : l \geq n + 1, k > l \text{ implise } |a_k| > 4r \cos \eta\}$ and therefore have

$$I_3(r) \geq \prod_{k=n_2(r)+1}^{\infty} \left(1 - 2 \frac{r}{|a_k|} \cos \eta + \frac{r^2}{|a_k|^2} \right) \geq \prod_{k=n_2(r)+1}^{\infty} \left(1 - 2 \frac{r}{|a_k|} \cos \eta \right) .$$

Using condition (b) and the definition of $n_2(r)$ we obtain

$$I_3(r) \geq \prod_{n=0}^{\infty} \left(1 - \frac{1}{2} q^n \right)^l = c_1(q) > 0 \quad \text{for } 0 < q < 1 .$$

We shall estimate $I_2(r)$ by (2.2) as follows

$$\begin{aligned} I_2(r) &\geq \prod_{k=n_1(r)+1}^{n_2(r)} \left(\frac{1}{2} \frac{\tan^2 \eta}{1 + \frac{1}{2} \tan^2 \eta} + \frac{r^2}{|a_k|^2} \cdot \frac{\sin^2 \eta}{2} \right) \\ &\geq \left(\frac{1}{2} \frac{\tan^2 \eta}{1 + \frac{1}{2} \tan^2 \eta} \right)^{n_2(r) - n_1(r) - 1} \end{aligned}$$

(if $n_1(r) < n + 1$ then instead of $n_2(r) - n_1(r) - 1$ we should write $n_2(r) - n - 1$). We can estimate $n_2(r) - n_1(r) - 1$ from above as follows; we shall find the smallest m satisfying $q^{-m} > 4r \cos \eta / r(2 \cos \eta)^{-1} = 8 \cos^2 \eta$ which we call m_0 , by (b) $m_0 \cdot l > n_2(r) - n_1(r)$.

Combining these results

$$\begin{aligned} \left| \prod_{k=n+1}^{\infty} (1 - r e^{i\theta} / a_k) \right|^{-2} &= I_1(r) \cdot I_2(r) I_3(r) \\ &\geq \prod_{n=1}^{\infty} \left(1 - \frac{1}{2} q^n \right)^l \cdot \left(\frac{1}{2} \frac{\tan^2 \eta}{1 + \frac{1}{2} \tan^2 \eta} \right)^{m_0 \cdot l} = c > 0. \end{aligned}$$

COROLLARY 2.1.a. *Under assumptions (a) and (b) the kernel function $G(t)$ satisfies*

$$(3.2) \quad E(s)^{-1} = \int_{-\infty}^{\infty} e^{-st} G(t) dt$$

$G(t) \in C^\infty(-\infty, \infty)$ and

$$(3.3) \quad G^{(n)}(t) = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} \frac{s^n e^{st} ds}{E(s)}.$$

REMARK 2.1.b. In Theorem 2.1 it is shown that if (a) and (b) are satisfied then for $0 < \eta < \pi/2 - \psi$

$$(3.4) \quad |E(r e^{i\theta})| \geq C(q, \eta)$$

for

$$\left\{ \theta : \left| \theta - \frac{\pi}{2} \right| < \frac{\pi}{2} - \psi - \eta \right\} \cup \left\{ \theta : \left| \theta - \frac{3\pi}{2} \right| < \frac{\pi}{2} - \psi - \eta \right\}$$

and $C(q, \eta)$ does not depend on r or the smallest $|a_k|$.

3. Asymptotic estimate of $G(t)$. Define α_1 and α_2 by:

$$(3.1) \quad \begin{aligned} \alpha_1 &= \max \{ \operatorname{Re} a_k, -\infty \mid \operatorname{Re} a_k < 0 \} \quad \text{and} \\ \alpha_2 &= \min \{ \operatorname{Re} a_k, \infty \mid \operatorname{Re} a_k > 0 \}. \end{aligned}$$

THEOREM 3.1. *Suppose $E(s) = \prod_{k=1}^{\infty} (1 - s/a_k)$, the sequence $\{a_k\}$ satisfies conditions (a) and (b) and let $G(t)$ be defined by (1.2), then:*

(i) $\alpha_1 = -\infty$ implies

$$G^{(n)}(t) = o(e^{ct}) \qquad t \rightarrow \infty$$

for all $c, c < 0$.

(ii) $\alpha_1 > -\infty$ implies

$$G^{(n)}(t) = \sum_{l=1}^L P_l(t) \exp(a_{k(l)}t) + o(e^{ct}) \quad t \rightarrow \infty$$

where $\operatorname{Re} a_{k(l)} = \alpha_1$, $1 \leq l \leq L$ ($a_{k(j)} \neq a_{k(i)}$ for $i \neq j$ and if $a_k \neq a_{k(l)}$, $1 \leq l \leq L$ then $\operatorname{Re} a_k \neq \alpha_1$), $P_l(t)$ are polynomials of degree μ_l where $\mu_l + 1$ is the multiplicity of $a_{k(l)}$ in $\{a_k\}$ and $c < \alpha_1$.

(iii) $\alpha_2 = \infty$ implies

$$G^{(n)}(t) = o(e^{ct}) \text{ for all } c, c > 0. \quad t \rightarrow -\infty$$

(iv) $\alpha_1 < \infty$ implies

$$G^{(n)}(t) = \sum_{l=L+1}^{L+M} P_l(t) \exp(a_{k(l)}t) + o(e^{ct}) \quad t \rightarrow -\infty$$

where $\operatorname{Re} a_{k(l)} = \alpha_2$ for $L + 1 \leq l \leq L + M$, $P_l(t)$ are as in (b) and $c > \alpha_2$.

Proof. The proof follows the well established method of Hirschman and Widder [5, p. 108]. In order to use this method it is enough to show that

$$|E(\sigma + i\tau)|^{-1} = O(|\tau|^{-n}) \quad |\tau| \rightarrow \infty$$

uniformly for $-A \leq \sigma < A$ for every finite A . By Theorem 2.1 we have for $|\tau|/|\sigma| > \tan(\psi + \eta)$ and therefore for $|\tau| > A \tan(\psi + \eta)$ (where $\eta > 0$, $\psi + \eta < \pi/2$ and ψ is defined in condition (a))

$$|E(\sigma + i\tau)|^{-1} \leq (A(n) + B(n)|\sigma + i\tau|^{2n})^{-1/2} \leq B(n)^{-1/2} |\tau|^{-n}.$$

4. $G_m(t)$ and properties. Define $G_m(t)$ by

$$(4.1) \quad E_m(s)^{-1} = \int_{-\infty}^{\infty} e^{-st} G_m(t) dt$$

where

$$(4.2) \quad E_m(s) = \prod_{k=m+1}^{\infty} (1 - s/a_k).$$

By Theorem 2.1 and Corollary 2.1.a we have $G_m(t) \in C^\infty(-\infty, -\infty)$ and for $m = 0, 1, 2, \dots$

$$(4.3) \quad G_m^{(n)}(t) = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} \frac{s^n e^{st}}{E_m(s)} ds.$$

The asymptotic estimates of Theorem 3.1 for $G_m(t)$ that satisfies condition (a) and (b) will be useful; however the following new estimate will be essential for the proof of the inversion formula.

THEOREM 4.1. *Let conditions (a) and (b) hold and $|a_k| \leq |a_{k+1}|$, then*

$$(4.4) \quad \begin{aligned} |G_m(t)| &\leq M_1 |a_{m+1}| \exp\left(-\frac{1}{2} \cos \psi |a_{m+1}t|\right) \\ &\quad + M_2 |a_{m+2}| \exp\left(-\frac{1}{2} \cos \psi |a_{m+2}t|\right). \end{aligned}$$

Proof. We shall divide the proof of (4.4) into two cases

(I) $|a_{m+2}| \leq 4 |a_{m+1}| / \cos \psi$

(II) $|a_{m+2}| \geq 4 |a_{m+1}| / \cos \psi$.

To show (4.4) in Case I we write $d_m = 1/2 \cos \psi |a_{m+1}|$ and since $d_m < |a_{m+1}| \cos \psi$ we have

$$(4.5) \quad G_m(t) = \frac{1}{2\pi i} \int_{d_m - i\infty}^{d_m + i\infty} \frac{e^{st}}{E_m(s)} ds = \frac{1}{2\pi i} \int_{-d_m - i\infty}^{-d_m + i\infty} \frac{e^{st}}{E_m(s)} ds .$$

Using the first integral we have

$$(4.6) \quad |G_m(t)| \leq \frac{1}{2\pi} e^{d_m t} \int_{-\infty}^{\infty} |E_m(d_m + i\tau)|^{-1} d\tau .$$

To estimate $G_m(t)$ we have to estimate $E_m(d_m + i\tau)$

$$\begin{aligned} |E_m(d_m + i\tau)| &= \prod_{i=1}^2 \frac{1}{|a_{m+i}|} |a_{m+i} - d_m - i\tau| \cdot \prod_{k=m+3}^{\infty} \left| 1 - \frac{d_m + i\tau}{a_k} \right| \\ &\equiv I_1(\tau) \cdot I_2(\tau) \end{aligned}$$

$$(4.7) \quad \begin{aligned} &|a_{m+i}|^{-2} |a_{m+i} - d_m - i\tau|^2 \\ &= (\operatorname{Re} a_{m+i} - d_m)^2 |a_{m+i}|^{-2} + (\tau - \operatorname{Im} a_{m+i})^2 |a_{m+i}|^{-2} \\ &\geq \frac{1}{4} \cos^2 \psi + (\tau^2 - 2\tau \operatorname{Im} a_{m+i} + (\operatorname{Im} a_{m+i})^2) |a_{m+i}|^{-2} \\ &\geq \frac{1}{8} \cos^2 \psi + \frac{1}{8} \frac{\cot^2 \psi}{1 + \frac{1}{8} \cot^2 \psi} \tau^2 |a_{m+i}|^{-2} \\ &\quad + \left(\left[1 + \frac{1}{8} \cot^2 \psi \right]^{-1} \tau^2 - 2\tau \operatorname{Im} a_{m+i} \right. \\ &\quad \left. + \left(1 + \frac{1}{8} \cot^2 \psi \right) (\operatorname{Im} a_{m+i})^2 \right) |a_{m+i}|^{-2} \geq A(\psi) + B(\psi) \tau^2 |a_{m+i}|^{-2} \end{aligned}$$

where $A(\psi) = 1/8 \cos^2 \psi$ and

$$B(\psi) = \frac{1}{8} \frac{\cot^2 \psi}{1 + \frac{1}{8} \cot^2 \psi} .$$

Using (4.7) and since $|a_{m+2}| \geq |a_{m+1}|$

$$I_1(\tau) \geq \{A(\psi)^2 + 2A(\psi)B(\psi) |a_{m+2}|^{-2} \tau^2 + B(\psi)^2 |a_{m+2}|^{-4} \tau^4\}^{1/2} \geq k_1(1 + \tau^2 |a_{m+1}|^{-2}) .$$

To estimate $I_2(\tau)$ we recall from Remark 2.1.b that when $d_m + i\tau = re^{i\theta}$ and $\psi + \eta \leq \theta \leq 2/\pi$ or $3\pi/2 \leq \theta \leq 2\pi - \psi - \eta$, that is when $|\tau| \geq d_m \tan(\psi + \eta)$, $I_2(\tau) \geq c > 0$.

When $|\tau| \leq d_m \tan(\psi + \eta)$ we choose $n_1 \geq m + 3$ such that $|a_{n_1}| \geq 2d_m(1 + \tan(\psi + \eta))$ and therefore

$$I_2(\tau) \geq \left\{ \prod_{k=m+3}^{n_1} A(\psi) \right\}^{1/2} \cdot \prod_{k=n_1+1}^{\infty} \left(1 - \frac{d_m(1 + \tan(\psi + \eta))}{|a_k|} \right) \geq A(\psi)^{(n_1-m-3)/2} \cdot \prod_{n=0}^{\infty} \left(1 - \frac{1}{2} q^n \right)^2$$

and since by condition (b) $n_1 - m - 3$ is bounded regardless of m $I_2(\tau) \geq c_1$.

Therefore (4.6) and (4.7) yield

$$|G_m(t)| \leq C_2 e^{d_m t} \int_{-\infty}^{\infty} [1 + |a_{m+1}|^{-2} \tau^2]^{-1} d\tau \leq M_1 |a_{m+1}| \exp\left(\frac{1}{2} \cos \psi |a_{m+1}| t\right) .$$

This estimation though correct for all t is valuable only for $t \leq 0$; for $t \geq 0$ we obtain the result taking the second integral of (4.5) into consideration.

In the Case II, $|a_{m+2}| > 4|a_{m+1}|/\cos \psi$, therefore

$$|\operatorname{Re} a_{m+1}| \leq |a_{m+1}| < \frac{1}{4} |\operatorname{Re} a_{m+2}| .$$

To prove our result for $t \leq 0$ we use the method of Theorem 3.1 and obtain

$$(4.8) \quad G_m(t) = a_{m+1} \frac{e^{a_{m+1}t}}{E_{m+1}(a_{m+1})} + \int_{k_m - i\infty}^{k_m + i\infty} \frac{e^{st} ds}{E_m(s)}$$

when $\operatorname{Re} a_{m+1} < 0$ and

$$(4.9) \quad G_m(t) = \int_{k_m - i\infty}^{k_m + i\infty} \frac{e^{st} ds}{E_m(s)}$$

when $\operatorname{Re} a_{m+1} > 0$; where $k_m = 1/2 |a_{m+2}| \cos \psi$.

Using (4.8) and (4.9) we obtain

$$(4.10) \quad |G_m(t) \leq |a_{m+1}| \cdot \frac{e^{|\operatorname{Re} a_{m+1}|t}}{E_{m+1}(a_{m+1})} + e^{k_m t} \int_{-\infty}^{\infty} [E_m(k_m + i\tau)]^{-1} d\tau$$

$$|E_{m+1}(a_{m+1})| \geq \prod_{k=2}^{\infty} \left(1 - \left| \frac{a_{m+1}}{a_{m+k}} \right| \right) \geq \prod_{n=0}^{\infty} \left(1 - \frac{1}{2} q^n \right)^i .$$

Considerations already used in this theorem show that

$$|E_{m+2}(k_m + i\tau)| \geq C > 0 .$$

Since $|a_{m+1}| \leq |a_{m+2}|$ and $|\operatorname{Re} a_{m+1} - k_m| \geq 1/2 k_m > |a_{m+1}|$ we have

$$\left| 1 - \frac{k_m + i\tau}{a_{m+1}} \right| \cdot \left| 1 - \frac{k_m + i\tau}{a_{m+2}} \right| \geq (A(\psi) + B(\psi)\tau^2 |a_{m+2}|^{-2}) .$$

Recalling that

$$E_m(k_m + i\tau) = E_{m+2}(k_m + i\tau) \cdot \left(1 - \frac{k_m + i\tau}{a_{m+1}}\right) \left(1 - \frac{k_m + i\tau}{a_{m+2}}\right)$$

the proof of Case II for $t \leq 0$ follows immediately. The proof when $t \geq 0$ is similar taking $-k_m$ instead of k_m .

THEOREM 4.2. *If conditions (a) and (b) are satisfied and $|a_k| \leq |a_{k+1}|$, then*

$$(4.4) \quad |G'_m(t)| \leq \sum_{i=1}^3 N_i |a_{m+i}|^2 \exp \left(-\frac{1}{2} \cos \psi |a_{m+i}t| \right) .$$

Proof. Since $(1 - a_k^{-1}D)G_m(t) = G_{m+1}(t)$ we have

$$G'_m(t) = -a_{m+1}G_{m+1}(t) + |a_{m+1}||G_m(t)| .$$

Using Theorem 4.1 for both m and $m + 1$ we obtain

$$\begin{aligned} |G'_m(t)| &\leq |a_{m+1}||G_{m+1}(t)| + |a_{m+1}||G_m(t)| \\ &\leq M_1 |a_{m+1}|^2 \exp \left(-\frac{1}{2} \cos \psi |a_{m+1}t| \right) \\ &\quad + M_2 |a_{m+1}||a_{m+2}| \exp \left(-\frac{1}{2} \cos \psi |a_{m+2}t| \right) \\ &\quad + M_1 |a_{m+1}||a_{m+2}| \exp \left(-\frac{1}{2} \cos \psi |a_{m+2}t| \right) \\ &\quad + M_2 |a_{m+1}||a_{m+3}| \exp \left(-\frac{1}{2} \cos \psi |a_{m+3}t| \right) \end{aligned}$$

which yields (4.4) easily.

REMARK 4.1.a. If in Theorems 4.1 and 4.2 the restriction, $|a_{m+1}| < L|a_m|$ for some $L > 1$, is added the proofs become obviously shorter and involve only the first term (in each theorem).

REMARK 4.1.b. It can be proved that if conditions (a) and (b) are satisfied and the multiplicity of a_{m+1} and a_{m+2} in $\{a_k\}$ is one then:

$$(a) \quad |G_m(t)| \leq M|a_{m+1}| \exp(-|\operatorname{Re} a_{m+1}||t|)$$

and

$$(b) \quad |G'_m(t)| \leq K|a_{m+1}|^2 \exp(-|\operatorname{Re} a_{m+1}||t|) + N|a_{m+1}||a_{m+2}| \exp(-|\operatorname{Re} a_{m+2}||t|).$$

These results are better than those of Theorem 4.1 and 4.2 but the proof I have uses those theorems. Since Theorems 4.1 and 4.2 are sufficient for the inversion result, I will not prove here these generalizations.

5. Inversion theorems. The results we shall obtain will correspond to the following two different situations: (1) Both α_1 and α_2 are finite. (2) Either α_1 or α_2 is non finite. (α_1 and α_2 were defined in § 3).

THEOREM 5.1. *Suppose:*

(1) *Conditions (a) and (b) are satisfied.*

(2) *The constants α_1 and α_2 are finite, $\left| \int_0^t \varphi(v)dv \right| \leq Ke^{(\alpha_2 - \varepsilon)t}$ for $t \geq 0$ and $\left| \int_t^0 \varphi(v)dv \right| \leq Ke^{(\alpha_1 + \varepsilon)t}$ for $t \leq 0$ for some $\varepsilon > 0$, and $\varphi(t) \in L_1(A, B)$ for all A, B satisfying $-\infty < A < B < \infty$.*

(3) *At a point $x \int_0^h [\varphi(x + y) - \varphi(x)]dy = o(h)$ $h \rightarrow 0$. Then*

$$(5.1) \quad \lim_{m \rightarrow \infty} P_m(D)f(x) = \varphi(x).$$

Proof. By Theorem 3.1 and assumption 2 we derive the uniform convergence in an interval $a \leq x \leq b$ of $\int_{-\infty}^{\infty} G^{(m)}(x - t)\varphi(t)dt$ and therefore

$$(5.2) \quad \frac{d^n}{dx^n} f(x) = \int_{-\infty}^{\infty} G^{(n)}(x - t)\varphi(t)dt$$

and

$$P_m(D)f(x) = \int_{-\infty}^{\infty} G_m(x - t)\varphi(t)dt.$$

To complete the proof we remember that $\int_{-\infty}^{\infty} G_m(t)dt = 1$ and therefore

$$\begin{aligned} |P_m(D)f(x) - \varphi(x)| &= \left| \int_{-\infty}^{\infty} G_m(x-t)[\varphi(t) - \varphi(x)]dt \right| \\ &= \left| \left\{ \int_{-\infty}^{x-\delta} + \int_{x-\delta}^{x+\delta} + \int_{x+\delta}^{\infty} \right\} G'_m(x-t)\alpha(t)dt \right| = I_1 + I_2 + I_3 \end{aligned}$$

where $\alpha(t)$ is given by $\alpha(t) = \int_x^t [\varphi(v) - \varphi(x)]dv$. Using (3) one can choose δ so that for $(x-t) \leq \delta$ $|\alpha(t)| \leq \varepsilon|x-t|$ and therefore

$$\begin{aligned} |I_2| &\leq \varepsilon \int_{x-\delta}^{x+\delta} |G'_m(x-t)||x-t| dt = \varepsilon \int_{-\delta}^{\delta} |G'_m(v)||v| dv \\ &\leq \varepsilon 2 \sum_{i=1}^3 N_i |a_{m+i}|^2 \cdot \int_0^{\delta} v \exp\left(-\frac{1}{2} \cos \psi |a_{m+i}| v\right) dv \\ &\leq \varepsilon 2 \sum_{i=1}^3 N_i \left\{ \delta |a_{m+i}| \exp\left(-\frac{1}{2} \cos \psi |a_{m+i}| \delta\right) \cdot 2 \cdot (\cos \psi)^{-1} \right. \\ &\quad \left. + 4 (\cos \psi)^{-2} \right\}. \end{aligned}$$

For any fixed δ $|a_{m+i}| \exp(-1/2 \cos \psi |a_{m+i}| \delta) = o(1)$ $m \rightarrow \infty$. Using (2) $|\alpha(t)| \leq K e^{(\alpha_2 - \varepsilon)t}$ for $t \geq 0$ and $|\alpha(t)| \leq K e^{(\alpha_1 + \varepsilon)t}$ for $t \leq 0$ and therefore

$$\begin{aligned} |I_3| &\leq K e^{(\alpha_1 + \varepsilon)(x+\delta)} \int_{x+\delta}^{\max(x+\delta, 0)} \sum_{i=1}^3 N_i |a_{m+i}|^2 \exp\left(\frac{1}{2} \cos \psi |a_{m+i}| t\right) \\ &\quad + \sum_{i=1}^3 K N_i |a_{m+i}|^2 \int_{\max(x+\delta, 0)}^{\infty} \exp\left(-\frac{1}{2} \cos \psi |a_{m+i}| t\right) e^{(\alpha_2 - \varepsilon)t} dt. \end{aligned}$$

Since $\lim_{x \rightarrow \infty} x e^{-ax} = 0$ for $a > 0$ we obtain $I_3 = o(1)m \rightarrow \infty$ and similarly $I_2 = o(1)m \rightarrow \infty$.

REMARK 5.1.a. Condition (2) can be replaced by a milder condition (2*) when there are only simple roots on $\text{Re } z = \alpha_1$ and $\text{Re } z = \alpha_2$. (2*) $|\alpha(t)| \leq K\chi(t)e^{\alpha_2 t}$ for $t \leq 0$ and $|\alpha(t)| \leq K\chi(t)e^{\alpha_1 t}$ for $t \geq 0$ where $\chi(t) > 0$ and $\int_{-\infty}^{\infty} \chi(t)dt < \infty$. In the proof the only change is in showing the uniform convergence (on a finite interval) of (5.2).

If for some $G(t)$ $\alpha_1 = -\infty$ then for $G^*(t)$, $G^*(t) \equiv G(-t)$ $\alpha_2 = \infty$ and vice versa. We shall treat therefore such kernels for which $\alpha_1 = -\infty$. For the inversion result we shall need the following lemma.

LEMMA. 5.2. *If conditions (a) and (b) are satisfied and $\alpha_1 = -\infty$, then $G(t) = 0$ for $t \geq 0$.*

Proof. Let $[1 - s/a_i]^{-1} = \int_{-\infty}^{\infty} e^{-st} g_i(t) dt$, then since $\text{Re } a_i > 0$

$$g_i(t) = \begin{cases} a_i e^{a_i t} & t < 0 \\ 0 & t > 0 \end{cases}$$

Define $G_m^*(t) = g_1 * g_2 * \dots * g_m(t)$, it is clear that $G_m^*(t) = 0$ for $t > 0$ (by induction) and that

$$G_m^*(t) = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} e^{st} \left[\prod_{k=1}^m \left(1 - \frac{s}{a_k} \right) \right]^{-1} ds .$$

We have, for all m

$$\begin{aligned} G(t) &= \int_{-\infty}^{\infty} G_m^*(v) G_m(t - v) dv = \int_{-\infty}^0 G_m^*(v) G_m(t - v) dv \\ |G_m^*(t)| &\leq |g_1(\cdot)| * |g_2(\cdot)| * \dots * |g_m(t)| \\ &= \prod_{i=1}^m \frac{|a_i|}{\text{Re } a_i} h_1 * \dots * h_m(t) \end{aligned}$$

where

$$h_i(t) = \begin{cases} \text{Re } a_i \exp(\text{Re } a_i t) & t < 0 \\ 0 & t > 0 \end{cases} .$$

It is well known that $|h_1 * \dots * h_m(t)| \leq \min_{1 < k \leq m} \text{Re } a_k$ (see [5, p. 138]) and that for $m \geq m_0$ $\min_{1 < k \leq m} \text{Re } a_k = \alpha_2$. Therefore for $m \geq m_0$

$$|G_m^*(t)| \leq \frac{\alpha_2}{\cos^m \psi} \quad (|a_i| \leq (\cos \psi)^{-1} |\text{Re } a_i|) .$$

Since we have

$$\begin{aligned} |G(t)| &\leq \alpha_2 \cdot (\cos \psi)^{-m} \int_{-\infty}^0 |G_m(t - v)| dv \\ &= \alpha_2 (\cos \psi)^{-m} \cdot \int_t^{\infty} |G_m(v)| dv \end{aligned}$$

using Theorem 4.1 we obtain for $t > 0$

$$\begin{aligned} |G(t)| &\leq \alpha_2 (\cos \psi)^{-m} \cdot 2(\cos \psi)^{-1} \left[M_1 \exp\left(-\frac{1}{2} \cos \psi |a_{m+1}| |t|\right) \right. \\ &\quad \left. + M_2 \exp\left(-\frac{1}{2} \cos \psi |a_{m+2}| |t|\right) \right] . \end{aligned}$$

Condition (b) implies for every $t \neq 0$

$$(\cos \psi)^{-m} \exp\left(-\frac{1}{2} \cos \psi |a_{m+i}| |t|\right) = o(1) \quad m \rightarrow \infty$$

for $i = 1, 2, \dots$. Therefore $G(t) = o(1)$ $m \rightarrow \infty$ for $t > 0$ and being independent of m $G(t) = 0$ for $t > 0$. Since $G(t) \in C^\infty(-\infty, \infty)$ $G(t) = 0$ for $t = 0$ also.

THEOREM 5.3. *Suppose:*

(1) *Conditions (a) and (b) are satisfied.*

(2) $\alpha_1 = -\infty$, $\varphi(t)$ is defined for $t \geq M$ and $\varphi(t) \in L_1(M, R)$ for all $R < \infty$ and $\int_M^t \varphi(t) dt \leq K e^{(\alpha_2 - \varepsilon)t}$.

(3) *Conditions (3) of Theorem 5.1 is satisfied.*

Then for $x > M$

$$(5.3) \quad \lim_{m \rightarrow \infty} P_m(D)f(x) = \varphi(x) .$$

Proof. The proof is almost identical to that of Theorem 5.1, but for the convergence of

$$\int_{-\infty}^{\infty} G^{(n)}(x-t)\varphi(t)dt \quad \text{and} \quad \int_{-\infty}^{\infty} G_m(x-t)\varphi(t)dt$$

we have to use also Lemma 5.2 (remembering that $G_m(t)$ satisfies also conditions (a) and (b) and $\alpha_1 = -\infty$ and therefore $G_m(t) = 0$ for $t \geq 0$).

REFERENCES

1. J. Dauns and D. V. Widder, *Convolution transforms whose inversion functions have complex roots*, Pacific J. Math. (2) **15** (1965), 427-442.
2. Z. Ditzian, *On asymptotic estimates for kernels of convolution transforms*, Pacific J. Math. (2) **21** (1967), 249-254.
3. ———, *On a class of convolution transforms*, Pacific J. Math. **25** (1968), 83-107.
4. I. I. Hirschman and D. V. Widder, *Convolution transforms with complex kernels*, Pacific J. Math. **1** (1951), 211-225.
5. ———, *The convolution transform*, Princeton, 1955.

Received November 27, 1967. This work was partially supported by NRC grant A-4816.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R. PHELPS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California 90024.

Each author of each article receives 50 reprints free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$ 8.00; single issues, \$ 3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$ 4.00 per volume; single issues \$ 1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners of publishers and have no responsibility for its content or policies.

| | |
|--|-----|
| Charles A. Akemann, <i>Invariant subspaces of $C(G)$</i> | 421 |
| Dan Amir and Zvi Ziegler, <i>Generalized convexity cones and their duals</i> | 425 |
| Raymond Balbes, <i>On (J, M, m)-extensions of order sums of distributive lattices</i> | 441 |
| Jan-Erik Björk, <i>Extensions of the maximal ideal space of a function algebra</i> | 453 |
| Frank Castagna, <i>Sums of automorphisms of a primary abelian group</i> | 463 |
| Theodore Seio Chihara, <i>On determinate Hamburger moment problems</i> | 475 |
| Zeev Ditzian, <i>Convolution transforms whose inversion function has complex roots in a wide angle</i> | 485 |
| Myron Goldstein, <i>On a paper of Rao</i> | 497 |
| Velmer B. Headley and Charles Andrew Swanson, <i>Oscillation criteria for elliptic equations</i> | 501 |
| John Willard Heidel, <i>Qualitative behavior of solutions of a third order nonlinear differential equation</i> | 507 |
| Alan Carleton Hindmarsh, <i>Pick's conditions and analyticity</i> | 527 |
| Bruce Ansgar Jensen and Donald Wright Miller, <i>Commutative semigroups which are almost finite</i> | 533 |
| Lynn Clifford Kurtz and Don Harrell Tucker, <i>An extended form of the mean-ergodic theorem</i> | 539 |
| S. P. Lloyd, <i>Feller boundary induced by a transition operator</i> | 547 |
| Henry B. Mann, Josephine Mitchell and Lowell Schoenfeld, <i>A new proof of the maximum principle for doubly-harmonic functions</i> | 567 |
| Robert Einsohn Mosher, <i>The product formula for the third obstruction</i> | 573 |
| Sam Bernard Nadler, Jr., <i>Sequences of contractions and fixed points</i> | 579 |
| Eric Albert Nordgren, <i>Invariant subspaces of a direct sum of weighted shifts</i> | 587 |
| Fred Richman, <i>Thin abelian p-groups</i> | 599 |
| Jordan Tobias Rosenbaum, <i>Simultaneous interpolation in H_2. II</i> | 607 |
| Charles Thomas Scarborough, <i>Minimal Urysohn spaces</i> | 611 |
| Malcolm Jay Sherman, <i>Disjoint invariant subspaces</i> | 619 |
| Joel John Westman, <i>Harmonic analysis on groupoids</i> | 621 |
| William Jennings Wickless, <i>Quasi-isomorphism and TFM rings</i> | 633 |
| Minoru Hasegawa, <i>Correction to "On the convergence of resolvents of operators"</i> | 641 |