

# Pacific Journal of Mathematics

**COMMUTATIVE SEMIGROUPS WHICH ARE ALMOST FINITE**

BRUCE ANSGAR JENSEN AND DONALD WRIGHT MILLER

## COMMUTATIVE SEMIGROUPS WHICH ARE ALMOST FINITE

BRUCE A. JENSEN AND DONALD W. MILLER

**Semigroups satisfying certain finiteness conditions are studied. It is shown that an infinite semigroup  $S$  every proper subsemigroup of which is finite is a group; thus in particular if  $S$  is commutative then it is isomorphic to the group  $Z(p^\infty)$  for some prime  $p$ . An infinite commutative semigroup every proper homomorph of which is finite is shown to be imbeddable in an infinite cyclic group with zero element adjoined and its structure is described.**

In his monograph on infinite abelian groups Kaplansky [2] includes the following two exercises concerning an infinite abelian group  $G$ :

(I) If every proper subgroup of  $G$  is finite then, for some prime  $p$ ,  $G$  is isomorphic to  $Z(p^\infty)$ .

(II) If every proper homomorph of  $G$  is finite then  $G$  is an infinite cyclic group.

The converse of each of these implications is, of course, also true.

It is natural to ask what conclusions can be drawn for commutative semigroups under analogous hypotheses. In §1 it is shown that if  $S$  is an infinite semigroup each of whose proper subsemigroups is finite then  $S$  is a group. Thus in particular if  $S$  is commutative the conclusion of (II) is obtained.

A semigroup  $S$  is said to be homomorphically finite, or, for brevity, *HF*, if  $S$  is infinite while each proper homomorph of  $S$  is finite. Section 2 is devoted to showing that an infinite commutative semigroup  $S$  is *HF* if and only if  $S$  is imbeddable in an infinite cyclic group. A description of all such semigroups is given.

The notation and terminology used in this paper follow that of Clifford and Preston [1].

1. Infinite semigroups whose proper subsemigroups are finite. We begin with a lemma.

**LEMMA 1.** *Let  $S$  be an infinite semigroup having no proper infinite subsemigroups. Then:*

- (i)  $S$  is periodic;
- (ii)  $S^2 = S$ ;
- (iii)  $S$  is not nil.

*Proof.* (i) If  $a \in S$  then either  $\langle a \rangle$ , the subsemigroup of  $S$

generated by  $a$ , is finite or  $S = \langle a \rangle$  is cyclic. In the latter case, however,  $S$  contains the proper infinite subsemigroup  $\langle a^2 \rangle$ , contrary to hypothesis.

(ii) If  $x \in S \setminus S^2$ , the complement of  $S^2$  in  $S$ , then  $S \setminus \{x\}$  is a proper infinite subsemigroup of  $S$ . Hence  $S = S^2$ .

(iii) If  $S$  contains no zero element, (iii) holds by default. Hence suppose that  $0$  is a zero element of  $S$  and that  $S$  is nil. By (ii) we can choose  $a \in S$  such that  $aS \neq 0$ . Hence either  $aS$  is finite or  $aS = S$ . If  $aS = S$  then  $S = a^n S$  for every positive integer  $n$  so, since  $S$  is nil,  $S = a^k S = 0S = 0$  for some positive integer  $k$ , a contradiction. Hence assume that  $aS = \{x_0, x_1, \dots, x_n\}$ , where  $n > 0$ ,  $x_0 = 0$  and  $x_i \neq x_j$  for  $i \neq j$ . For  $i = 0, 1, \dots, n$ , define

$$S_i = \{y \in S \mid ay = x_i\}$$

and define a binary relation  $\leq$  on the set  $\mathcal{S}$  of all  $S_i$  by stipulating that  $S_i \leq S_j$  if and only if there exists  $s$  in  $S^1$  such that  $x_j = x_i s$ . Clearly  $\leq$  is reflexive and transitive on  $\mathcal{S}$ . Moreover suppose  $S_i \leq S_j$  and  $S_j \leq S_i$ , say  $x_j = x_i s$  and  $x_i = x_j t$ , where  $s, t \in S^1$ . Then

$$x_j = x_j (ts)^k, \quad k = 1, 2, 3, \dots$$

Since  $S$  is nil this implies that either  $x_j = 0$ , whence also  $x_i = 0$ , or  $ts \notin S$ . In the latter case,  $s = t = 1$  so again  $x_i = x_j$ . Thus  $\leq$  is a partial ordering of  $\mathcal{S}$ .

Evidently  $S_i \leq S_0$  for  $i = 0, 1, \dots, n$ . Moreover there must exist an integer  $N, 1 \leq N \leq n$ , such that

$$(1) \quad S_i \leq S_N \text{ implies } i = N, \text{ all } S_i \in \mathcal{S}.$$

Let  $y \in S_N$ . By (ii)  $y = uv$  for some  $u, v \in S$ . Since  $\mathcal{S}$  describes a partition of  $S$ ,  $u \in S_i$  for exactly one  $i, 0 \leq i \leq n$ . Therefore  $x_N = ay = auv = x_i v$  so  $S_i \leq S_N$  whence, by (1),  $i = N$  and  $x_N = x_N v$ . Consequently  $x_N = x_N v^k$  for  $k = 1, 2, 3, \dots$  so  $x_N = 0 = x_0$ , contrary to  $N > 0$ . This establishes (iii).

**THEOREM 1.** *If  $S$  is an infinite semigroup each of whose proper subsemigroups is finite then  $S$  is a group.*

*Proof.* Let  $A_1 = \{x \in S \mid xS \text{ is finite}\}$  and  $A_2 = \{x \in S \mid xS = S\}$ . For  $i = 1, 2$ , if  $A_i \neq \emptyset$  (the null set) then  $A_i$  is a subsemigroup of  $S$ . Thus, since  $A_1$  and  $A_2$  partition  $S$ , either  $A_1 = S$  and  $A_2 = \emptyset$  or vice versa. An analogous argument on the principal left ideals of  $S$  leads to the conclusion that  $S$  satisfies exactly one of the following:

- (i)  $xS$  and  $Sx$  are finite, all  $x \in S$ ;
- (ii)  $xS$  is finite and  $Sx = S$ , all  $x \in S$ ;

- (iii)  $xS = S$  and  $Sx$  is finite, all  $x \in S$ ;
- (iv)  $xS = Sx = S$ , all  $x \in S$ .

Denote the set of idempotents of  $S$  by  $E$ ; since  $S$  is periodic,  $E \neq \emptyset$ .

Case (i). If  $E$  is finite then  $S^1ES^1$  is a finite ideal of  $S$  and the Rees factor semigroup  $S/S^1ES^1$  is an infinite semigroup having only finite proper subsemigroups. However some power of each element of  $S$  is idempotent so  $S/S^1ES^1$  is nil. Since this contradicts Lemma 1,  $E$  must be infinite.

For each  $e \in E$  define the subsemigroups  $L_e$  and  $R_e$  of  $S$  by

$$L_e = \{x \in S \mid xe = e\}, \quad R_e = \{y \in S \mid ey = e\}.$$

If  $L_e = L_f = R_e = R_f = S$  for some  $e, f \in E$  then  $e = ef = f$ . Hence, since  $E$  is infinite, there must exist an  $e$  in  $E$  such that either  $L_e \neq S$  or  $R_e \neq S$ . Assume the former; then  $L_e$  is finite. Therefore  $E' = E \setminus L_e$  is an infinite subset of  $S$  so  $E'$  generates  $S$ . Consequently there exist elements  $f_1, f_2, \dots, f_k$  of  $E'$  such that  $f_1 f_2 \dots f_k = e$ . Thus  $f_1 e = e$  so  $f_1 \in L_e$ , contradicting  $f_1 \in E'$ . The assumption that  $R_e \neq S$  leads to a similar contradiction, so case (i) is eliminated.

Cases (ii) and (iii) are left-right duals so only one of them need be considered. Suppose then that for each  $x$  in  $S$ ,  $Sx$  is finite and  $xS = S$ . Then  $ex = x$  for all  $e \in E$  so  $E$  is a right zero subsemigroup of  $S$ . Thus if  $e \in E$  and  $E' = E \setminus \{e\}$  then  $E'$  is either empty or a proper subsemigroup of  $S$ . In either case we conclude that  $E$  is finite.

Suppose  $E = \{e\}$ . Then, since  $S$  is periodic, there corresponds to each element  $x$  of  $S$  a positive integer  $n = n(x)$  such that  $x^n = e$ . Therefore  $xe = x^{n+1} = ex = x$  so  $Se = S$ , contradicting the finiteness of  $Se$ . Hence  $E$  has order  $k > 1$ , say  $E = \{e_1, \dots, e_k\}$ . For each  $i$ ,  $1 \leq i \leq k$ , define  $T_i = \{x \in S \mid e_i \in \langle x \rangle\}$ . By the periodicity of  $S$ , the set  $\{T_1, \dots, T_k\}$  is a partition of  $S$ . If  $x \in T_i$ , say  $x^n = e_i$ , then, as above, it follows that  $xe_i = x$ , so that  $T_i \subseteq U_i = \{x \in S \mid xe_i = x\}$ ,  $i = 1, \dots, k$ . Conversely let  $x \in U_i$  and suppose  $x^n = e_j$  for some  $n > 0$  and  $j$ ,  $1 \leq j \leq k$ . As above,

$$e_j = x^n = x^n e_i = e_j e_i = e_i.$$

Thus  $i = j$  so  $U_i \subseteq T_i$ . Therefore  $T_i = U_i$  is a subsemigroup of  $S$  for  $i = 1, \dots, k$ . But since  $S = \bigcup^k T_i$  and  $k > 1$  it follows that at least one of the  $T_i$  is an infinite proper subsemigroup of  $S$ , a contradiction.

This leaves only case (iv). Thus  $S$  is a group.

Combining Theorem 1 with Kaplansky's Exercise (I) we then have the following result.

**THEOREM 2.** *If  $S$  is an infinite commutative semigroup each of*

whose proper subsemigroups is finite then  $S$  is isomorphic to the group  $Z(p^\infty)$  for some prime  $p$ .

2. **Commutative HF semigroups.** An HF (or homomorphically finite) semigroup is defined to be an infinite semigroup each of whose noninjective homomorphisms has finite image. An HF group is an HF semigroup which is also a group, e.g., the infinite cyclic group or any infinite simple group. The following two results are immediate consequences of these definitions.

LEMMA 2. *Every proper nonzero ideal of an HF semigroup  $S$  has finite complement in  $S$ .*

LEMMA 3. *If  $S$  is an infinite semigroup then either all or none of  $S, S^1, S^0$  and  $(S^1)^0$  are HF semigroups.*

L. Rédei [3, Satz 82] has given essentially the following characterization of the subsemigroups of the additive semigroup  $N$  of positive integers.

LEMMA 4. (Rédei). *Let  $N$  be the additive semigroup of all positive integers and let  $d, r \in N$ . Define*

$$(2) \quad I = \{nd \mid n \in N, nd \geq r\}$$

and let  $A$  be any subset of  $dN \setminus I$  such that  $A + A \subseteq A \cup I$ . Then  $S = S(d, r, A)$  is a subsemigroup of  $N$ , and every subsemigroup of  $N$  is so obtainable. Furthermore, for suitable choice of  $r'$  and  $A'$ ,  $S(d, r, A) \cong S(1, r', A')$ .

Rédei's result can easily be extended to the additive group  $Z$  of all integers.

LEMMA 5. *If  $S$  is a nonzero subsemigroup of  $Z$  then either  $S$  is isomorphic to  $Z$  or, for suitable  $r \in N$  and  $A \subseteq rN$ ,  $S$  is isomorphic to  $S(1, r, A)$  with or without an adjoined identity.*

*Proof.* In view of the isomorphism between the subsemigroups  $N$  and  $-N = (-1)N$  of  $Z$  we need only consider those subsemigroups of  $Z$  which contain both a positive and a negative integer. Let  $S$  be such a semigroup and let  $S_1 = S \cap N, S_2 = S \cap (-N)$ . For  $i = 1, 2$  it follows from Lemma 4 that  $S_i$  is isomorphic to  $S(d_i, r_i, A_i)$  for suitable  $d_i, r_i, \in N$  and  $A_i \subseteq r_i N \setminus I_i$ , where  $I_i = \{nd_i \mid n \in N, nd_i \geq r_i\}$ . Moreover  $ud_1 \in S_1$  and  $-vd_2 \in S_2$ , and hence  $ud_1 - vd_2 \in S$ , for all sufficiently large integers  $u$  and  $v$ . Therefore  $(d_1, d_2) \in S$  so  $d_1 = d_2$ . It then follows that  $S$  is the cyclic subgroup of  $Z$  generated by  $d$ .

Commutative *HF* semigroups can now be characterized.

**THEOREM 3.** *Let  $S$  be an infinite commutative semigroup. Then  $S$  is homomorphically finite if and only if  $S$  is imbeddable in an infinite cyclic group with adjoined zero. If this is the case then either  $S$  is itself an infinite cyclic group or  $S$  is isomorphic to a subsemigroup  $\bar{S}$  of the additive semigroup of all nonnegative integers with zero element  $\infty$  adjoined. In the latter event there exist positive integers  $a_1, \dots, a_k$  and  $r$  such that*

$$\bar{S} = \{a_1, a_2, \dots, a_k\} \cup \{n \mid n \geq r\},$$

possibly with adjoined zero element  $\infty$ .

*Proof.* Let  $S$  be a subsemigroup of the additive group of integers with adjoined zero  $\infty$ . By Lemma 3 there is no loss of generality in assuming that  $\infty \notin S$ . Hence by Lemma 5 either  $S$  is an infinite cyclic group, and thus is homomorphically finite, or  $S$  is isomorphic to some semigroup  $S(1, r, A)$ , so that  $S = I \cup A$ , where  $I = \{n \mid n \in \mathbb{N}, n \geq r\}$  is an ideal of  $S$ . Assuming the latter, let  $\sigma$  be a nontrivial congruence on  $S$  which is not one-to-one, so that  $a \sigma b$  for two distinct elements  $a, b$  of  $S$ . Then  $(na) \sigma (nb)$  for all  $n > 0$  so  $\sigma$  is not one-to-one on  $I$ , whence we can assume that  $a, b \in I$ , with  $a < b$ . Then  $(a + (r + k)) \sigma (b + (r + k))$  for each  $k \geq 0$ .

Define  $m = b - a$  and let  $x, y \in I$ , with  $x, y \geq a + r$  and  $x \equiv y \pmod{m}$ , say  $x = a + s, y = a + s + tm$ , where  $s, t \in \mathbb{N}$  and  $s \geq r$ . Since  $a \sigma b$  then  $(a + s) \sigma (b + s)$ , i.e.,  $(a + s) \sigma (a + s + m)$ . Thus by induction  $(a + s) \sigma (a + s + tm)$  so  $x \sigma y$ . It follows that the factor semigroup  $I/\sigma$  is finite so by the finiteness of  $A, S/\sigma$  is also finite. Therefore  $S$  is an *HF*-semigroup.

Conversely let  $S$  be a commutative *HF* semigroup. For each  $c$  in  $S$  define the congruence  $\sigma_c$  on  $S$  by

$$a \sigma_c b \text{ if and only if } ac = bc, \text{ all } a, b \in S.$$

If there exists an element  $c$  in  $S$  such that  $S/\sigma_c$  is finite then the ideal  $Sc$  of  $S$  would also be finite, which, in the light of Lemma 2, would contradict the assumption that  $S$  is infinite unless  $Sc = 0$ .

Suppose  $Sc = 0$  and let  $J = \{x \in S \mid Sx = 0\}$ . Then  $J$  is an ideal of  $S$  so either  $J = 0$  or  $S/J$  is finite. In the latter case, we conclude that  $S^2$  is also finite; thus  $S^2 = 0$  since  $S^2$  and  $S/S^2$  cannot both be finite. However it is evident that the condition  $S^2 = 0$  cannot hold in an *HF* semigroup  $S$ , so  $J = 0$ . Hence  $Sc = 0$  only if  $c = 0$ .

Thus  $S/\sigma_c$  is infinite, and  $\sigma_c$  is one-to-one, for all  $c$  in  $S \setminus \{0\}$  so  $S$  is a commutative cancellative semigroup, possibly with an adjoined zero. In any event,  $S$  contains no proper zero divisors.

Let  $T = S \setminus 0$  or  $T = S$  according as  $S$  does or does not contain a zero element. Let  $G$  denote the group of quotients of  $T$  and regard  $T$  as a subsemigroup of  $G$ , in the usual manner. Suppose  $\sigma$  is a congruence on  $G$  which is not one-to-one, say  $(a/b) \sigma (c/d)$ , where  $a, b, c, d \in S$  and  $ad \neq bc$ . Then  $((a/b)bd) \sigma ((c/d)bd)$ , i.e.,  $(ad) \sigma (bc)$ . Consequently  $\sigma'$ , the restriction of  $\sigma$  to  $S$ , is not one-to-one on  $T$  so  $T/\sigma'$  is finite.

For  $x \in T$  let  $[x]$  and  $[x]'$  denote the  $\sigma$ -class of  $G$  and the  $\sigma'$ -class of  $T$ , respectively, containing  $x$ . Then the homomorphism of  $T/\sigma'$  into  $G/\sigma$  defined by  $[x]' \rightarrow [x]$ , all  $x \in T$ , is injective. Thus  $T/\sigma'$  is cancellative and hence is a finite abelian group. It is readily verified that the mapping of  $S/\sigma'$  defined by  $[x] \rightarrow [x]'$ , all  $x \in S$ , is an isomorphism of  $S/\sigma'$  onto  $G/\sigma$ . Therefore  $G/\sigma$  is also finite so  $G$  is an abelian *HF*-group. Thus by Kaplansky's Exercise II,  $G$  is cyclic.

An application of Lemma 5 now completes the proof.

#### REFERENCES

1. A.H. Clifford and G.B. Preston, *The Algebraic Theory of Semigroups*, Vol. 1, American Mathematical Society, Providence, Rhode Island, 1961.
2. I. Kaplansky, *Infinite Abelian Groups*, The University of Michigan Press, Ann Arbor, Michigan, 1954.
3. L. Rédei, *Theorie der Endlich Erzeugbaren Kommutativen Halbgruppen*, B. G. Teubner Verlagsgesellschaft Leipzig, 1963.

Received October 31, 1967. Under the partial support of a N.S.F. Nebraska Cooperative Fellowship.

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. ROYDEN  
Stanford University  
Stanford, California

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. R. PHELPS  
University of Washington  
Seattle, Washington 98105

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

\* \* \*

AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL WEAPONS CENTER

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California 90024.

Each author of each article receives 50 reprints free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$ 8.00; single issues, \$ 3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$ 4.00 per volume; single issues \$ 1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners of publishers and have no responsibility for its content or policies.



Charles A. Akemann, <i>Invariant subspaces of <math>C(G)</math></i> .....	421
Dan Amir and Zvi Ziegler, <i>Generalized convexity cones and their duals</i> ....	425
Raymond Balbes, <i>On <math>(J, M, m)</math>-extensions of order sums of distributive lattices</i> .....	441
Jan-Erik Björk, <i>Extensions of the maximal ideal space of a function algebra</i> .....	453
Frank Castagna, <i>Sums of automorphisms of a primary abelian group</i> .....	463
Theodore Seio Chihara, <i>On determinate Hamburger moment problems</i> .....	475
Zeev Ditzian, <i>Convolution transforms whose inversion function has complex roots in a wide angle</i> .....	485
Myron Goldstein, <i>On a paper of Rao</i> .....	497
Velmer B. Headley and Charles Andrew Swanson, <i>Oscillation criteria for elliptic equations</i> .....	501
John Willard Heidel, <i>Qualitative behavior of solutions of a third order nonlinear differential equation</i> .....	507
Alan Carleton Hindmarsh, <i>Pick's conditions and analyticity</i> .....	527
Bruce Ansgar Jensen and Donald Wright Miller, <i>Commutative semigroups which are almost finite</i> .....	533
Lynn Clifford Kurtz and Don Harrell Tucker, <i>An extended form of the mean-ergodic theorem</i> .....	539
S. P. Lloyd, <i>Feller boundary induced by a transition operator</i> .....	547
Henry B. Mann, Josephine Mitchell and Lowell Schoenfeld, <i>A new proof of the maximum principle for doubly-harmonic functions</i> .....	567
Robert Einsohn Mosher, <i>The product formula for the third obstruction</i> .....	573
Sam Bernard Nadler, Jr., <i>Sequences of contractions and fixed points</i> .....	579
Eric Albert Nordgren, <i>Invariant subspaces of a direct sum of weighted shifts</i> .....	587
Fred Richman, <i>Thin abelian <math>p</math>-groups</i> .....	599
Jordan Tobias Rosenbaum, <i>Simultaneous interpolation in <math>H_2</math>. II</i> .....	607
Charles Thomas Scarborough, <i>Minimal Urysohn spaces</i> .....	611
Malcolm Jay Sherman, <i>Disjoint invariant subspaces</i> .....	619
Joel John Westman, <i>Harmonic analysis on groupoids</i> .....	621
William Jennings Wickless, <i>Quasi-isomorphism and TFM rings</i> .....	633
Minoru Hasegawa, <i>Correction to "On the convergence of resolvents of operators"</i> .....	641