

# Pacific Journal of Mathematics

**THIN ABELIAN  $p$ -GROUPS**

FRED RICHMAN

## THIN ABELIAN $p$ -GROUPS

FRED RICHMAN

**An abelian  $p$ -group  $G$  is thin if every map from a torsion complete group to  $G$  is small. The class of thin groups is shown to be closed under arbitrary direct sums and extensions and hence any direct sum of countable reduced  $p$ -groups is thin. An example shows that, unlike for groups with no elements of infinite height, a reduced group may contain no unbounded torsion complete subgroups and still fail to be thin. Finally, these groups are used to settle questions in a certain relative homological algebra.**

This investigation was motivated by the study of the relative homological algebra induced in the category of abelian  $p$ -groups (or abelian groups) by declaring the torsion complete  $p$ -groups to be projective. An exact sequence  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$  is proper in this setting if every map from a torsion complete  $p$ -group  $G$  to  $C$  lifts to a map from  $G$  to  $B$ . In particular, since cyclic  $p$ -groups are torsion complete, the sequence  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$  must be  $p$ -pure, i.e.,  $p^n B \cap A = p^n A$  for every positive integer  $n$ . This condition alone enables us to lift any map from  $G$  to  $C$  which factors through a direct sum of cyclic  $p$ -groups. Thus if every map from a torsion complete  $p$ -group to  $C$  factors through a direct sum of cyclic groups the notions of  $p$ -pure and proper coincide.

Pierce [5] called a homomorphism  $\phi$  from a  $p$ -group  $G$  *small* if the kernel of  $\phi$  contained a large subgroup. A *large subgroup* is a subgroup of the form  $\sum p^{n_k} G[p^{n_k}]$  where  $\{n_k\}$  is an increasing sequence of nonnegative integers. If  $L$  is a large subgroup then  $G/L$  is a direct sum of cyclics. (This is clear for  $G$  a direct sum of cyclics. If  $B$  is a basic subgroup of  $G$  then  $L \cap B$  is large in  $B$ , since  $B$  is pure, and  $B + L = G$  by the Baer decomposition. Hence  $G/L = (B + L)/L \cong B/L \cap B$  is a direct sum of cyclics<sup>1</sup>.) Conversely, Megibben [3] has shown that any map from a torsion complete  $p$ -group to a direct sum of cyclics is small. It is therefore of some interest to study  $p$ -groups to which any homomorphism from a torsion complete group is small.

There is an analogy here with the notion of a slender torsion free group as defined by Łoś. A group  $G$  is *slender* if any map from the direct product of countably many infinite cyclic groups annihilates cofinitely many of these cyclic groups. The idea in both cases is to examine groups for which any homomorphism from a fixed, rather

---

<sup>1</sup> I am indebted to Doyle Cutler for this important but little publicized observation.

tightly held together group must have a rather large kernel (a fixed unbounded torsion complete group may be selected, instead of considering the whole class). In either case the point of the study is perhaps more to understand the structure of the fixed group than the class to which it gives rise.

All groups considered in this paper will be abelian.

2. Thin groups. We formally introduce the main characters:

DEFINITION. A  $p$ -group  $G$  is *thin* if every map from a torsion complete group to  $G$  is small.

Our class is clearly closed under subgroups. It is also closed under extensions and direct sums.

THEOREM 1. *If  $H$  and  $G/H$  are thin then so is  $G$ .*

*Proof.* Let  $T$  be torsion complete and  $\phi: T \rightarrow G$ . Then  $T \rightarrow G \rightarrow G/H$  is small so for every  $k$  there exists an  $n_k$  such that  $\phi(L) \subseteq H$  where  $L = \sum p^{n_k} T[p^k]$ . Now  $L$  is torsion complete. Indeed, since  $T/L$  is a direct sum of cyclics we have the exact sequence

$$0 \rightarrow \text{Ext}(Z_{p^\infty}, L) \rightarrow \text{Ext}(Z_{p^\infty}, T)$$

and since  $p^\omega \text{Ext}(Z_{p^\infty}, T) = \text{Pext}(Z_{p^\infty}, T) = 0$  we have

$$\text{Pext}(Z_{p^\infty}, L) = p^\omega \text{Ext}(Z_{p^\infty}, L) = 0$$

and so  $L$  is torsion complete. Thus  $\phi$  restricted to  $L$  is small. Hence  $\phi$  is small.

THEOREM 2. *If  $G = \sum G_\alpha$  and  $G_\alpha$  is thin for all  $\alpha$  then  $G$  is thin.*

*Proof.* Let  $T$  be torsion complete and  $\phi: T \rightarrow G$ . If  $\phi$  is not small then there exist  $x_1, x_2, x_3 \dots \in T[p^n]$ ,  $ht(x_i) \geq i$ , such that  $\phi(x_i) \neq 0$ . Since  $\pi_\alpha \circ \phi$  is small, where  $\pi_\alpha$  is the projection  $G \rightarrow G_\alpha$ , the  $x_j$  may be chosen such that the  $\phi(x_j)$  have disjoint supports. Relabeling (and applying Theorem 1) we may assume that  $\phi(x_i) \in G_i$ . Let  $\pi_i$  be the projection on  $G_i$ . Since  $\pi_i \phi$  is small there is an  $m$  such that  $\pi_i \phi(x_{m+1} + x_{m+2} + \dots) = 0$ . Hence

$$\pi_i \phi(x) = \pi_i \phi(x_1 + \dots + x_m) = \phi(x_i) \neq 0.$$

This is a contradiction since  $\phi(x)$  can have only finitely many nonzero coordinates.

The following is the analogue of the theorem that a direct sum of countable reduced torsion free groups is slender.

**COROLLARY 1.** *Any direct sum of countable reduced  $p$ -groups is thin.*

*Proof.* Clearly  $Z_p$  is thin. Thus any group constructable from  $Z_p$  by a sequence of taking direct sums, subgroups and extensions is also thin. In particular, direct sums of countable reduced groups are thin.

A  $p$ -group is *fully starred* if no subgroup admits a dense subset of smaller cardinality (in the  $p$ -adic topology on the subgroup). Fully starred groups are closed under subgroups, direct sums [4] and extensions [1]. Since  $Z_p$  is fully starred so is any direct sum of countable reduced  $p$ -groups. We may extend Corollary 1 to:

**COROLLARY 2.** *Any fully starred  $p$ -group is thin.*

*Proof.* Let  $G$  be fully starred and  $\phi$  a map from a torsion complete group  $\bar{B}$  to  $G$ . If  $\phi$  is not small there exists a sequence  $x_1, x_2, x_3, \dots \in \bar{B}[p^n]$  such that  $ht(x_i) \rightarrow \infty$  and  $\phi(x_i) \neq 0, i = 1, 2, 3, \dots$ . Let  $H$  be a countable pure subgroup of  $\bar{B}$  containing each  $x_i$ . Then the closure,  $\bar{H}$ , of  $H$  in  $\bar{B}$  is a torsion complete group and  $\phi$  restricted to  $\bar{H}$  is not small. Now  $\phi(\bar{H})$  is not countable by Corollary 1 (a fully starred group must be reduced since 0 is dense in  $Z_{p^\infty}$ ). But  $\phi(H)$  is countable and dense in  $\phi(\bar{H})$ , which is impossible since  $G$  is fully starred.

**3. A counterexample.** If  $G$  contains an unbounded torsion complete subgroup then  $G$  is certainly not thin. Megibben showed that the converse is true for groups without elements of infinite height. In this section we show that this restriction cannot be dropped.

We shall be looking at the socle of a group  $G$  with no elements of infinite height. This socle inherits the  $p$ -adic topology from  $G$ . All topological terminology refers to this topology.

**LEMMA 1.** *Let  $G$  be a  $p$ -group with no elements of infinite height and  $H$  a torsion complete subgroup of  $G$ . Then  $H$  contains a non-discrete complete subgroup of  $G[p]$ .*

*Proof.* Choose  $x_1, x_2, x_3, \dots \in H[p]$  such that  $ht_H(x_{i+1}) > ht_G(x_i)$ . Then  $\sum n_i x_i$  converges (in  $H$  or  $G$ ) and  $ht(\sum n_i x_i) = ht(x_j)$  where  $j = \min \{i \mid p \nmid n_i\}$ . Then  $S = \{\sum n_i x_i \mid n_i \in \mathbb{Z}\}$  is a nondiscrete complete subgroup of  $G[p]$ .

LEMMA 2. *If  $H$  is an extension of a bounded  $p$ -group by a torsion complete  $p$ -group and  $p^o H = 0$  then  $H$  is torsion complete.*

*Proof.* Let  $0 \rightarrow S \rightarrow H \rightarrow \bar{B} \rightarrow 0$  be exact with  $S$  bounded and  $\bar{B}$  torsion complete. Then

$$0 \rightarrow \text{Ext}(Z_{p^\infty}, S) \rightarrow \text{Ext}(Z_{p^\infty}, H) \rightarrow \text{Ext}(Z_{p^\infty}, \bar{B}) \rightarrow 0$$

is exact. Now  $\text{Pext}(Z_{p^\infty}, \bar{B}) = 0$  so  $\text{Ext}(Z_{p^\infty}, \bar{B})$  has no elements of infinite height. Hence  $\text{Pext}(Z_{p^\infty}, H) \subseteq \text{Ext}(Z_{p^\infty}, S)$  is bounded. But  $H$  is the torsion subgroup of  $\text{Ext}(Z_{p^\infty}, H)$  and  $H$  has no elements of infinite height. Hence  $\text{Pext}(Z_{p^\infty}, H) = 0$  and so  $H$  is torsion complete.

THEOREM 3. *Let  $G$  be torsion complete,  $\phi$  a map from  $G$  onto  $K$  such that  $\ker \phi$  is bounded. If  $K$  contains an unbounded torsion complete subgroup  $\bar{B}$  then  $\ker \phi$  either contains or misses a nondiscrete complete subgroup of  $G[p]$ .*

*Proof.* Let  $H = \phi^{-1}(\bar{B})$ . Then  $H$  is torsion complete by Lemma 2. Let  $S = \ker \phi \cap H[p]$ . Since  $\ker \phi$  and  $H[p]$  are closed in  $H$ , so is  $S$ . Let  $H_1$  be a pure subgroup of  $H$  such that  $H_1[p] = S$  (see [2], Lemma 1). Then  $H_1$  is torsion complete ([2], Lemma 2) and so  $H = H_1 \oplus H_2$ . Lemma 1 says that if  $H_1$  is unbounded then  $S$ , and hence  $\ker \phi$ , contains a nondiscrete complete subgroup of  $G[p]$  whereas if  $H_2$  is unbounded then  $S$ , and hence  $\ker \phi$ , misses such a subgroup of  $G[p]$ .

To find our example we need only find a subgroup of the socle of an unbounded torsion complete group which neither misses nor contains a nondiscrete complete subsocle.

THEOREM 4. *Let  $G$  be a torsion complete group with a countable unbounded basic subgroup. Then there exists a subgroup  $S \subseteq G[p]$  such that if  $K \subseteq G[p]$  is nondiscrete complete then  $0 \neq S \cap K \neq K$ .*

*Proof.* Since  $G$  has a countable basic there are  $2^{\aleph_0}$  complete nondiscrete subgroups of  $G[p]$ . Well order these subgroups,  $K_\alpha$ , by the ordinals  $\alpha < 2^{\aleph_0}$ . Construct subgroups  $S_\alpha$  and  $T_\alpha$  of  $G[p]$  inductively by:

$$(1) \quad S_0 = 0 = T_0.$$

$$(2) \quad S_{\alpha+1} = S_\alpha + \langle s_\alpha \rangle, \quad T_{\alpha+1} = T_\alpha + \langle t_{\alpha+1} \rangle$$

where  $s_\alpha \in K_\alpha \setminus (S_\alpha + T_\alpha)$  and  $t_\alpha \in K_\alpha \setminus (S_{\alpha+1} + T_\alpha)$ .

$$(3) \quad S_\beta = \bigcup_{\alpha < \beta} S_\alpha, \quad T_\beta = \bigcup_{\alpha < \beta} T_\alpha \quad \text{at limit ordinals } \beta.$$

The existence of  $s_\alpha$  and  $t_\alpha$  as required in the second step is guaranteed since  $S_\alpha$  and  $T_\alpha$  have cardinality less than  $K_\alpha$ . Set  $S = \bigcup S_\alpha$  and  $T = \bigcup T_\alpha$ . Then  $S \cap T = 0$  but every nondiscrete complete

subsocle  $K_\alpha$  has nontrivial intersection with both  $S$  and  $T$ .

**COROLLARY.** *There exists a reduced  $p$ -group  $A$  such that  $A$  contains no unbounded torsion complete subgroup but  $A$  is not thin.*

*Proof.* Let  $A = G/S$  where  $G$  and  $S$  are as in Theorem 4. Then Theorem 3 implies that  $A$  contains no unbounded torsion complete subgroup whereas the map  $G \rightarrow A$  is certainly not small.

**4. An application.** We return to the relative homological algebra induced on the category of  $p$ -groups (or all groups) by declaring torsion complete  $p$ -groups to be projective. It turns out that the thin  $p$ -groups enable us to characterize the injectives in this algebra without requiring a deep understanding of what ‘‘proper’’ means.

**LEMMA 3.** *If  $G[p]$  is complete then  $p^\omega G$  is  $p$ -divisible.*

*Proof.* Let  $x \in p^\omega G$  and  $p^{i+1}y_i = x$  for  $i = 0, 1, 2, \dots$ . Then  $py_0 = p^{i+1}y_i$  and so  $y_0 - p^i y_i \in G[p]$ . Since  $y_0 - p^i y_i$  is a Cauchy sequence in  $G[p]$  it converges to some element  $z \in G[p]$ . Since  $p^i y_i$  converges to 0 we have  $y_0$  converges to  $z$ , i.e.,  $y_0 = z + g$  where  $g \in p^\omega G$ . Thus  $x = py_0 = pz + pg$  where  $g \in p^\omega G$ .

**COROLLARY.** *A  $p$ -group  $G$  is the direct sum of a torsion complete group and a divisible group if and only if  $G[p]$  is complete.*

*Proof.* The ‘‘only if’’ is clear. The ‘‘if’’ part follows from the lemma, which implies that  $G = R \oplus D$  where  $R$  has no elements of infinite height and  $D$  is divisible, and Lemma 2 of [2] which says that  $R$  is torsion complete.

**THEOREM 5.** *A reduced  $p$ -group  $G$  is injective in the relative homological algebra induced in the category of  $p$ -groups by declaring the torsion complete groups projective  $\iff G$  is torsion complete.*

*Proof.*  $\Leftarrow$  is well known. The torsion complete groups are pure injective in this category.

$\Rightarrow$  We need only show that  $G[p]$  is complete. Let  $B \xrightarrow{\phi} C$  be a pure map of a direct sum of cyclic  $p$ -groups onto a reduced countable group  $C$  such that  $C^1 \neq 0$ . The kernel,  $K$ , of  $\phi$  is a pure direct sum of cyclics. Now  $K[p]$  is not closed in  $B$  since  $C^1 \neq 0$ . Let  $z \in B[p] \setminus K$  be in the closure of  $K[p]$  in  $B$ . Let  $K = K_0 \oplus K_1 \oplus \dots$  where  $K_1$  is a direct sum of cyclic groups of order  $p^{i+1}$ . Then there exist integers

$0 \leq n_1 < n_2 < \dots$  and elements  $0 \neq z_i \in K_{n_i}[p]$  such that

$$ht(z - z_1 - z_2 - \dots - z_i) > p^{n_i}.$$

Choose  $x_i \in K_{n_i}$  such that  $p^{n_i}x_i = z_i$ . Then  $\sum \langle x_i \rangle$  is direct and is a summand of  $K$ ,  $o(x_i) = p^{n_i+1}$  and  $\sum p^{n_i}x_i$  converges to  $z$ .

Let  $\sum g_i$  be a Cauchy series in  $G[p]$ , i.e.,  $ht(g_i) \rightarrow \infty$ . We must show that  $\sum g_i$  converges to an element of  $G[p]$ . We may assume that  $ht(g_i) \geq n_i$ . Thus we can map  $K \rightarrow G$  taking  $p^{n_i}x_i$  to  $g_i$ . This map extends to  $B$  since every map of a torsion complete group into  $C$  is small and hence lifts to  $B$ . The image of  $z$  is then a limit of the Cauchy series  $\sum g_i$ .

Similarly we may characterize the relative injectives in the category of all groups.

**THEOREM 6.** *A reduced group  $G$  is injective in the relative homological algebra induced on the category of groups by declaring the torsion complete  $p$ -groups projective  $\iff G$  is a complete  $Q_p$ -module, where  $Q_p$  is the ring of integers localized at  $p$ .*

*Proof.*  $\Leftarrow$  is well known. These are the reduced  $p$ -pure injectives.

$\Rightarrow$  That  $G$  is a  $Q_p$  module follows from the fact that  $Z \subseteq Q_p$  is proper in this setting (since the quotient is  $p$ -torsion free) and the only reduced images of  $Q_p$  are cyclic  $Q_p$ -modules. By Theorem 5 the torsion subgroup of  $G$  is torsion complete and hence  $p^\omega G$  is divisible by  $\aleph_{\omega+1}$  3. Hence  $G$  has no elements of infinite height. Finally, to show that  $G$  is complete consider  $\sum_{i=1}^\infty Q_p$  and the pure subgroup generated by  $(1, -p, 0, 0, \dots)$ ,  $(0, 1, -p, 0, \dots)$ ,  $(0, 0, 1, -p, 0, \dots)$  etc. If  $\sum g_i$  is a Cauchy series in  $G$ ,  $ht(g_i) \geq i$ , then  $g_i = p^i h_i$  and we can map  $(0, 0, \dots, 1, -p, \dots)$  onto  $h_i$  and extend to  $\sum_{i=1}^\infty Q_p$ . Then  $\sum g_i$  converges to the image of  $(1, 0, 0, \dots)$ .

5. **An aside.** One of the questions which arises in the study of any relative homological algebra is whether a proper (i.e., "pure") subgroup of a projective is again projective. For the relative homological algebra induced in the category of  $p$ -groups by declaring torsion complete groups to be projective, the projectives are direct sums of torsion complete and divisible groups (see [6]). Just what a proper subgroup "is" in this setting is still a little mysterious. The only nontrivial ones which have been exhibited thus far are (ordinary) pure subgroups with thin cokernels. These kinds of subgroups of projectives are again projective. It suffices to consider only direct sums of torsion complete groups. We first prove a pair of very general lemmas.

LEMMA 4. *If  $K$  is a  $p$ -group, the map  $\phi: G \rightarrow K$  is pure and onto  $\Leftrightarrow \phi(p^n G[p]) = p^n K[p]$  for every nonnegative integer  $n$ .*

*Proof.*  $\Rightarrow$  If  $x \in p^n K[p]$  then  $x = p^n y$ . We can find  $g \in G$  of the same order as  $y$  such that  $\phi(g) = y$ . Then  $p^n g \in p^n G[p]$  and  $\phi(p^n g) = x$ .

$\Leftarrow$  Suppose  $y \in K$  and  $p^{n+1}y = 0$ . Then  $p^n y \in p^n K[p]$  and so  $p^n y = \phi(p^n g)$  where  $p^n g \in G[p]$ . Now  $p^n(\phi(g) - y) = 0$  and so, by induction,  $\phi(g) - y = \phi(g_0)$  where  $g_0 \in G$  and  $p^n g_0 = 0$ . Thus  $y = \phi(g - g_0)$  and  $p^{n+1}(g - g_0) = 0$ .

LEMMA 5. *If  $f_1$  and  $f_2$  are maps from  $G_1$  and  $G_2$  to  $H$ , and  $\phi$  is a map from  $G_1$  to  $G_2$  such that  $f_2 \phi = f_1$  then the kernel of  $f_1 \oplus f_2: G_1 \oplus G_2 \rightarrow H$  is  $A \oplus \ker f_2$  where  $A \cong G_1$ .*

*Proof.* Let  $A = \{g_1 - \phi(g_1) \mid g_1 \in G_1\}$ . Then  $A \subseteq \ker(f_1 \oplus f_2)$ . The map taking  $g_1$  to  $g_1 - \phi(g_1)$  is an isomorphism from  $G_1$  to  $A$ . Since  $A \cap G_2 = 0$  we have  $A \cap \ker f_2 = 0$ . It remains to show that  $\ker(f_1 \oplus f_2) \subseteq A + \ker f_2$ . Suppose  $f_1(g_1) + f_2(g_2) = 0$ . Then  $f_2(\phi(g_1) + g_2) = 0$  so  $\phi(g_1) + g_2 \in \ker f_2$ . But  $g_1 + g_2 = g_1 - \phi(g_1) + \phi(g_1) + g_2$ .

THEOREM 7. *Let  $G$  be a direct sum of torsion complete  $p$ -groups and  $K$  a thin  $p$ -group. If  $\phi$  is a pure map from  $G$  onto  $K$  then the kernel of  $\phi$  is a direct sum of torsion complete groups.*

*Proof.* Write  $G = \sum T_i$  where  $T_i$  is torsion complete. Since  $\phi$  induces a small map on  $T_i$  there exists an  $n_i$  such that  $\phi(p^{n_i} T_i[p]) = 0$ . Letting  $B_i$  be a maximal  $p^{n_i}$  bounded summand of  $T_i$  we have  $T_i = A_i \oplus B_i$  where  $B_i$  is bounded and  $\phi(A_i[p]) = 0$ . Since  $\phi(\sum A_i[p]) = 0$  and  $\phi$  is pure onto  $K$  we have, by Lemma 4,  $\phi(p^n \sum B_i[p]) = p^n K[p]$  for every  $n \geq 0$ . Hence, by Lemma 4 again,  $\phi$  restricted to  $\sum B_i$  is pure onto  $K$ . Since  $\phi$  restricted to  $\sum A_i$  factors through a direct sum of cyclics ( $\phi$  is small on  $A_i$ ), and  $\phi \mid \sum B_i$  is pure onto  $K$ , we can factor  $\phi \mid \sum A_i$  through  $\phi \mid \sum B_i$ . Lemma 5 then implies that  $\ker \phi \cong \sum A_i \oplus C$  where  $C$  is a subgroup of  $\sum B_i$  and is therefore a direct sum of cyclic groups.

### REFERENCES

1. J. M. Irwin and F. Richman, *Direct sums of countable groups and related concepts*, J. of Alg. **2** (1965), 443-450.
2. J. M. Irwin, F. Richman and E. A. Walker, *Countable direct sums of torsion complete groups*, Proc. Amer. Math. Soc. **17**, (1966), 763-766.
3. C. Megibben, *Large subgroups and small homomorphisms*, Michigan Math. J. **13** (1966), 153-160.

4. R. J. Nunke, "Purity and subfunctors of the identity," in *Topics in Abelian Groups*, edited by J. M. Irwin and E. A. Walker, Scott, Foresman, 1963.
5. R. S. Pierce, "Homomorphisms of primary Abelian Groups," in *Topics in Abelian Groups*, edited by J. M. Irwin and E. A. Walker, Scott, Foresman, 1963.
6. F. Richman, C. Walker and E. A. Walker, *Projective classes of Abelian Groups* (to appear in Proc. Colloq. Abelian Groups, Montpellier, 1967).

Received September 21, 1967. This research supported by NSF-GP-6564.

NEW MEXICO STATE UNIVERSITY

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. ROYDEN  
Stanford University  
Stanford, California

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. R. PHELPS  
University of Washington  
Seattle, Washington 98105

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL WEAPONS CENTER

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California 90024.

Each author of each article receives 50 reprints free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners of publishers and have no responsibility for its content or policies.

# Pacific Journal of Mathematics

Vol. 27, No. 3

March, 1968

Charles A. Akemann, <i>Invariant subspaces of <math>C(G)</math></i> . . . . .	421
Dan Amir and Zvi Ziegler, <i>Generalized convexity cones and their duals</i> . . . .	425
Raymond Balbes, <i>On <math>(J, M, m)</math>-extensions of order sums of distributive lattices</i> . . . . .	441
Jan-Erik Björk, <i>Extensions of the maximal ideal space of a function algebra</i> . . . . .	453
Frank Castagna, <i>Sums of automorphisms of a primary abelian group</i> . . . . .	463
Theodore Seio Chihara, <i>On determinate Hamburger moment problems</i> . . . . .	475
Zeev Ditzian, <i>Convolution transforms whose inversion function has complex roots in a wide angle</i> . . . . .	485
Myron Goldstein, <i>On a paper of Rao</i> . . . . .	497
Velmer B. Headley and Charles Andrew Swanson, <i>Oscillation criteria for elliptic equations</i> . . . . .	501
John Willard Heidel, <i>Qualitative behavior of solutions of a third order nonlinear differential equation</i> . . . . .	507
Alan Carleton Hindmarsh, <i>Pick's conditions and analyticity</i> . . . . .	527
Bruce Ansgar Jensen and Donald Wright Miller, <i>Commutative semigroups which are almost finite</i> . . . . .	533
Lynn Clifford Kurtz and Don Harrell Tucker, <i>An extended form of the mean-ergodic theorem</i> . . . . .	539
S. P. Lloyd, <i>Feller boundary induced by a transition operator</i> . . . . .	547
Henry B. Mann, Josephine Mitchell and Lowell Schoenfeld, <i>A new proof of the maximum principle for doubly-harmonic functions</i> . . . . .	567
Robert Einsohn Mosher, <i>The product formula for the third obstruction</i> . . . . .	573
Sam Bernard Nadler, Jr., <i>Sequences of contractions and fixed points</i> . . . . .	579
Eric Albert Nordgren, <i>Invariant subspaces of a direct sum of weighted shifts</i> . . . . .	587
Fred Richman, <i>Thin abelian <math>p</math>-groups</i> . . . . .	599
Jordan Tobias Rosenbaum, <i>Simultaneous interpolation in <math>H_2</math>. II</i> . . . . .	607
Charles Thomas Scarborough, <i>Minimal Urysohn spaces</i> . . . . .	611
Malcolm Jay Sherman, <i>Disjoint invariant subspaces</i> . . . . .	619
Joel John Westman, <i>Harmonic analysis on groupoids</i> . . . . .	621
William Jennings Wickless, <i>Quasi-isomorphism and TFM rings</i> . . . . .	633
Minoru Hasegawa, <i>Correction to "On the convergence of resolvents of operators"</i> . . . . .	641