

Pacific Journal of Mathematics

A THEOREM ON SEQUENTIAL CONVERGENCE OF MEASURES AND SOME APPLICATIONS

JOHN BLIGH CONWAY

A THEOREM ON SEQUENTIAL CONVERGENCE OF MEASURES AND SOME APPLICATIONS

JOHN B. CONWAY

If S is a locally compact Hausdorff space, let βS be its Stone-Cech compactification and let $M(S)$ be the space of all finite complex valued regular Borel measures on S . In this paper we will prove that whenever S is paracompact and $\{\mu_n\}$ is a sequence in $M(\beta S)$ which converges to zero in the weak star topology, then $\lim \int_S f d\mu_n = 0$ for every continuous function f , and $\{\mu_n\}$ satisfies a certain uniformity condition on S . This generalizes a result of R. S. Phillips on weak star sequential convergence in the dual of l^∞ . Moreover, by using our theorem we can obtain many previously known results whose proofs, though all similar, were apparently independent.

Indeed, the motivation for undertaking the research which led to this paper came from the similarity of the proofs of a number of theorems. Among these is the following.

THEOREM A. *If S is paracompact and H is a subset of $M(S)$ which is countably compact in the $C(S)$ topology on $M(S)$ then for every $\varepsilon > 0$ there is a compact set $K_\varepsilon \subset S$ such that*

$$|\mu|(S - K_\varepsilon) < \varepsilon$$

for every μ in H .

This theorem was first proved by LeCam [15] and then, independently, by the present author [4]. Another result which utilizes the same method of proof referred to above is

THEOREM B. *If S is paracompact and A is a closed linear subspace of $C(\beta S)$ such that A interpolates $\beta S - S$ then there is a closed neighborhood V of $\beta S - S$ such that A interpolates V .*

NOTE. To say that a closed linear subspace A of $C(\beta S)$ interpolates a closed set $K \subset \beta S$ means that any f in $C(K)$ has an extension F to βS such that F is in A .

Theorem B is due to Bade [1] for σ -compact spaces. In addition to these two results there are still others which fit into this same category. These will be presented in §3 together with the proofs using the main theorem of this paper. However, we have not yet been able to discover a proof of Theorem A based on this result.

Theorem B can be proved by means of Theorem 2.2, but we have postponed this to a later paper.

1. Preliminaries. The original proofs of the two theorems above, as well as those of some of the theorems in § 3, employ the following approach: Using some previous result; an embedding of l^∞ into $C(S)$, the space of bounded continuous complex valued functions on S , is constructed. The adjoint of this map is examined and a result of R. S. Phillips is applied to achieve the desired conclusion. This result of Phillips is the following (see [5, p. 32] for a proof).

THEOREM 1.1. *If $\{\mu_n\}$ is a sequence in $M(\beta N)$ which converges to zero in the weak * topology of $M(\beta N)$ then*

$$\lim_{m \rightarrow \infty} \sum_{k=m} |\mu_n(\{k\})| = 0$$

uniformly in n .

Here we have taken liberty to identify l^∞ with $C(N)$ and $(l^\infty)^*$ with $M(\beta N)$; N is the set of positive integers with the discrete topology. The proof of our theorem will also use this result.

The notation and terminology used here will be standard as is found in [9], [12], and [14]; however, there will be a few minor deviations. We will denote by $C_0(S)$ those functions in $C(S)$ which vanish at infinity, and $C_{00}(S)$ will be the space of continuous functions having compact support. The term "weak * topology" will be restricted to mean only the $\sigma(M(\beta S), C(\beta S))$ topology on $M(\beta S)$. The $\sigma(M(S), C_0(S))$ and $\sigma(M(S), C(S))$ topologies on $M(S)$ will be referred to as the C_0 -weak * and C -weak * topologies respectively. Notice that if we consider $M(S)$ as a subset of $M(\beta S)$ then the C -weak * topology is exactly the relativisation of the weak * topology to $M(S)$.

If μ is in $M(S)$ and B is a Borel subset of S then μ_B denotes the restriction of μ to B . That is, $\mu_B(A) = \mu(B \cap A)$ for every Borel set A . To simplify our notation, we will not use any symbol to distinguish between a function in $C(S)$ and its extension to βS .

Finally, we wish to state the characterization of paracompact locally compact spaces needed below. The proof may be found in [9, p. 241].

THEOREM 1.2. *A locally compact space S is paracompact if and only if $S = \bigcup_\alpha S_\alpha$ where the S_α are pairwise disjoint open and closed σ -compact subsets of S .*

2. The main theorem. If H is a subset of $M(S)$ then we say that H is *tight* if and only if H is uniformly bounded and for every $\varepsilon > 0$ there is a compact set $K_\varepsilon \subset S$ such that $|\mu|(S - K_\varepsilon) < \varepsilon$ for every μ in H . (In [4] the author showed a relationship between tightness and the strict topology on $C(S)$). With this definition we state a well-known lemma whose proof will not be given.

LEMMA 2.1. *Let $\{\mu_n\}$ be a sequence in $M(S)$ which is tight and let μ be in $M(S)$ such that*

$$\lim_{n \rightarrow \infty} \int_S \varphi d\mu_n = \int_S \varphi d\mu$$

for each φ in $C_{00}(S)$. Then $\{\mu_n\}$ converges to μ C -weak $*$.

We now give the main result.

THEOREM 2.2. *If S is paracompact and $\{\mu_n\}$ is a sequence in $M(\beta S)$ which converges weak $*$ to μ in $M(\beta S)$ then $\{\mu_{n_S}\}$ is tight and converges C -weak $*$ to μ_S .*

REMARKS. It is easily seen that $\lim \int_S \varphi d\mu_n = \int_S \varphi d\mu$ for each φ in $C_{00}(S)$. Hence, by Lemma 2.1, the important part of the result is that $\{\mu_{n_S}\}$ is tight. A moments reflection will show that tightness in $M(N) = l^1$ is exactly the uniform limit condition in Phillips' theorem. Therefore this is a generalization of Theorem 1.1.

Proof. As was noted in the above remarks we need only show that $\{\mu_{n_S}\}$ is tight. Since this proof is similar to that of another result of the author [4, p. 478], we will omit many of the details. As in [4] we will only consider the case where S is σ -compact, the general theorem being accomplished by means of Theorem 1.2. Therefore, let $S = \bigcup_{n=1}^\infty D_n$ where each D_n is compact and D_n is contained in $\text{int } D_{n+1}$ (the interior of D_{n+1}).

Suppose $\{\mu_{n_S}\}$ is not tight; since $\{\mu_n\}$ is uniformly bounded there is an $\varepsilon > 0$ such that for every $k \geq 1$ there is a μ_{n_k} , subsets K_k and U_k of S , and a function φ_k having the properties:

- (a) K_k is compact, U_k is open in S with U_k^- compact and $U_k^- \cap K_k$ empty;
- (b) $D_k \cup K_k \cup U_k^- \subset \text{int } K_{k+1}$;
- (c) $|\mu_{n_k}(U_k)| > \varepsilon/4$;
- (d) φ_k is in $C_{00}(S)$, $\|\varphi_k\|_\infty = 1$, the support of φ_k is contained in U_k and $|\mu_{n_k}|(U_k) < \left| \int_S \varphi_k d\mu_{n_k} \right| + \varepsilon/8$. If for every $\xi = \{x_k\}$ in

l^∞ and $s \in S$ we define

$$T(\xi)(s) = \sum_{k=1}^{\infty} x_k \varphi_k(s)$$

then $T(\xi)$ is in $C(S)$ and $\|T(\xi)\|_\infty = \|\xi\|_\infty$. It follows that T is an isometry of l^∞ into $C(S)$. Therefore $T^*: M(\beta S) \rightarrow (l^\infty)^* = M(\beta N)$ exists and is weak * continuous. From our hypothesis on $\{\mu_n\}$ it follows that $\{T^*(\mu_{n_k})\}$ converges to $T^*(\mu)$ weak * in $M(\beta N)$. It is an easy matter to check that

$$T^*(\mu_{n_k})(\{j\}) = \int_{\beta S} \varphi_j d\mu_{n_k} = \int_S \varphi_j d\mu_{n_k}.$$

Applying Theorem 1.1 we get that

$$\lim_{m \rightarrow \infty} \sum_{j=m}^{\infty} \left| \int_S \varphi_j d\mu_{n_k} \right| = 0$$

uniformly in k . In particular, we have that

$$\lim_{k \rightarrow \infty} \int_S \varphi_k d\mu_{n_k} = 0$$

which, when combined with condition (d) above, yields a contradiction to (c). This completes the proof of the theorem.

3. Applications. The applications here are to sequential convergence in $M(S)$. For further results and references concerning this topic the reader is referred to the papers of Varadarajan [16] and Dudley [8].

The first result is due to Diendoné [7]. It should be noted that his original proof was a generalization of a proof of a theorem of Schür. However, it is an immediate consequence of Theorem 2.2.

THEOREM 3.1. *If S is paracompact and $\{\mu_n\}$ is a C -weak * convergent sequence in $M(S)$ then $\{\mu_n\}$ is tight.*

The next theorem was proved independently by Varadarajan [16, p. 195] and Collins and Dorroh [2]. Both have different proofs; Varadarajan uses a method of LeCam [15, p. 218] and Collins and Dorroh use the method of Theorem 2.2.

THEOREM 3.2. *If S is paracompact then $M(S)$ is C -weak * sequentially complete. Moreover, if $\{\mu_n\}$ is a C -weak * Cauchy sequence then $\{\mu_n\}$ is tight.*

Proof. If $\{\mu_n\}$ is a C -weak * Cauchy sequence in $M(S)$ then

consider each μ_n as an element in $M(\beta S)$. Since $\sup \{ \|\mu_n\| : n \geq 1 \} < \infty$, Alagolu's theorem implies that there is a μ in $M(\beta S)$ which is a weak * cluster point of $\{\mu_n\}$. But for each f in $C(S)$,

$$\lim_{n \rightarrow \infty} \int_{\beta S} f d\mu_n = \lim_{n \rightarrow \infty} \int_S f d\mu_n$$

exists and so $\{\mu_n\}$ converges to μ weak * in $M(\beta S)$. Therefore, Theorem 2.2 implies that $\{\mu_n\}$ converges to μ_S C -weak * in $M(S)$ and $\{\mu_n\}$ is tight. This completes the proof.

The following theorem of Grothendieck [11] characterizes the weakly convergent sequences in $M(S)$ (i.e., the sequences which converge for the $\sigma(M(S), M(S)^*)$ topology).

THEOREM 3.3. *If S is an arbitrary locally compact space and μ, μ_1, μ_2, \dots are elements in $M(S)$, then the following are mutually equivalent:*

- (a) $\{\mu_n\}$ converges to μ weakly in $M(S)$;
- (b) $\{\mu_n\}$ is uniformly bounded and $\mu(U) = \lim \mu_n(U)$ for every open F_σ set $U \subset S$;
- (c) $\lim \int_S f d\mu_n = \int_S f d\mu$ for every bounded lower semicontinuous function f ;
- (d) (i) $\{\mu_n\}$ is tight, (ii) $\{\mu_n\}$ converges to μ C -weak *, and (iii) for every $\varepsilon > 0$ and every compact set $K \subset S$ there is an open set $V_\varepsilon \supset K$ such that $|\mu_n|(V_\varepsilon - K) < \varepsilon$ for every $n \geq 1$.

Proof. It is trivial to see that (a) implies (b) and that (b) and (c) are equivalent. Therefore, assume that (c) holds and let us prove (d). Since we are only dealing with a countable family of regular measures we can find an open σ -compact set $S_1 \subset S$ such that

$$|\mu|(S - S_1) = |\mu_n|(S - S_1) = 0$$

for every $n \geq 1$. If f is a nonnegative function of S which is continuous on S_1 and vanishes on $S - S_1$, then f is lower semicontinuous. From this we have that $\{\mu_n\}$ converges to μ C -weak * in $M(S_1)$. Since S_1 is σ -compact, Theorem 2.2 implies $\{\mu_n\}$ is tight in $M(S_1)$ and hence in $M(S)$.

Since (ii) is clearly true it remains only to prove (iii). By means of (i), the proof of the general case requires only minor technical changes in the proof of the case when S is compact. Hence, let us make this assumption. Let $\varepsilon > 0$ and let K be a given compact subset of S . By passing to a larger compact set with the same measure as K , we may assume that K is a G_δ set. But then $W = S - K$ is an

open σ -compact subset of S . As above, it follows that $\{\mu_{n_W}\}$ converges to μ_W C-weak $*$ in $M(W)$. Hence $\{\mu_{n_W}\}$ is tight in $M(W)$; that is, there is a compact set $D \subset W$ such that $|\mu_n|(W - D) < \varepsilon$ for every n . But if $V = S - D$, V is open, $K \subset V$ and $|\mu_n|(V - K) = |\mu_n|(W - D) < \varepsilon$ for each n .

Now let us assume that (d) is true and prove (a). As before we will only consider the case where S is compact; we will further assume that $\mu = 0$. Observe that with S compact condition (iii) is equivalent to the following:

(iii)' for every $\varepsilon > 0$ and every open set U there is a compact set $K \subset U$ such that $|\mu_n|(U - K) < \varepsilon$ for each n .

Using (iii)', Urysohn's Lemma, and property (ii) we get that $\lim \mu_n(U) = 0$ for every open set U . It remains to show that this implies that $\lim \mu_n(A) = 0$ for each Borel set A in S (see [10, p. 308]). But again we need only show this when $A = \cup K_n$, where each K_n is compact and $K_n \subset K_{n+1}$. Let $\varepsilon > 0$ and for each n choose an open set $V_n \supset K_n$ such that for every $k \geq 1$

$$|\mu_k|(V_n - K_n) < \varepsilon \left(\frac{1}{2}\right)^{n+1}.$$

Let $V = \bigcup_{n=1}^{\infty} V_n$; then V is open, $A \subset V$ and for every k

$$\begin{aligned} |\mu_k|(V_n - A) &= |\mu_k|\left(\bigcap_{j=1}^{\infty} (V_n - K_j)\right) \\ &\leq |\mu_k|(V_n - K_n) \\ &\leq \varepsilon \left(\frac{1}{2}\right)^{n+1}. \end{aligned}$$

Thus,

$$|\mu_k|(V - A) = |\mu_k|\left(\bigcup_{n=1}^{\infty} (V_n - A)\right) \leq \frac{1}{2}\varepsilon$$

for all k . But $\lim \mu_n(V) = 0$ and so $|\mu_n(A)| < \varepsilon$ for sufficiently large n . This completes the proof of the theorem.

REMARKS. The proof that (d) implies (a) does not employ Theorem 2.2, but was given here for completeness. Dieudonné [6] showed that (c) and (d) are equivalent for S compact and that (a) and (b) are equivalent when S is a compact metric space. In fact, when S is metrisable he showed that the condition of uniform boundedness in (b) can be dropped.

A compact space S is said to be a σ -Stonian space if and only

if every bounded sequence of continuous real valued functions has a supremum. This is equivalent to the proposition that the closure of every open F_σ set is open. One interesting property of a σ -Stonian space is that if U is an open F_σ -set then βU is exactly U^- , the closure of U . Using this fact and our Theorem 2.2 we can prove the following result of Grothendieck [11, p. 168] and Isbell and Semadeni [13, p. 46].

THEOREM 3.4. *If the compact space S is a σ -Stonian space and $\{\mu_n\}$ is a sequence in $M(S)$ which converges weak $*$ then $\{\mu_n\}$ converges weakly.*

Proof. Let μ be the weak $*$ limit of $\{\mu_n\}$. By the preceding theorem we must show that $\mu(U) = \lim \mu_n(U)$ whenever U is an open F_σ set. But then $\beta U = U^-$ and U^- is open in S . Hence, if ν_n (respectively, ν) is the restriction of μ_n (respectively, μ) to U^- we have that $\{\nu_n\}$ converges to ν weak $*$ in $M(\beta U)$. Therefore, by Theorem 2.2 we have that $\{\nu_{nU}\}$ converges to ν_U C -weak $*$ in $M(U)$. In particular, we get that $\mu(U) = \lim \mu_n(U)$. This completes the proof of the theorem.

REMARKS. Grothendieck proved this for S a Stonian space and Isbell and Semadeni showed it to be true in the more general σ -Stonian spaces.

As a final application we give a generalization of Phillips' theorem that c_0 is not complemented in l^∞ .

THEOREM 3.5. *If S is paracompact and not compact then there is no bounded projection from $C(S) = C(\beta S)$ onto $C_0(S)$.*

Proof. Suppose P is a bounded projection from $C(\beta S)$ onto $C_0(S)$. Then $P^*: M(S) \rightarrow M(\beta S)$ is a bounded linear map which is C_0 -weak $*$ — weak $*$ continuous. Also, if φ is in $C_0(S)$ and μ is in $M(S)$ we have

$$\int_{\beta S} \varphi dP^*(\mu) = \int_S \varphi d\mu .$$

Therefore $P^*(\mu)_S = \mu$ for all μ in $M(S)$. But since S is paracompact and not compact we can, by Theorem 1.2, choose a sequence $\{s_n\}$ in S which is eventually in the complement of every compact subset of S . Hence, $\lim \varphi(s_n) = 0$ for every φ in $C_0(S)$. If μ_n is the unit point mass at s_n then $\{\mu_n\}$ converges C_0 -weak $*$ to zero in $M(S)$. Thus, $\{P^*(\mu_n)\}$ converges to zero weak $*$ in $M(\beta S)$. By Theorem 2.2 we have that $\{\mu_n\} = \{P^*(\mu_n)_S\}$ converges to zero C -weak $*$ in $M(S)$ and is tight. Clearly this is a contradiction and the theorem is proved.

REMARKS. This proof is the analogue of Phillips' proof that c_0 is not complemented in l^∞ , where Theorem 2.2 is used in place of Phillips' Theorem 1.1. In [3] the present author has proved a more general result than Theorem 3.5. Also Theorem 3.5 can be obtained as a corollary to Bade's Theorem B.

BIBLIOGRAPHY

1. W. G. Bade, *Extension of interpolating sets*, Functional Analysis—Proceedings of a conference held at the University of California, Irvine; edited by B. R. Gelbaum, Thompson Book Company, Washington, D.C. 1967.
2. H. S. Collins and J. R. Dorroh, *Remarks on certain function spaces*, Math. Ann. **176** (19768), 87-95.
3. J. B. Conway, *Projection and retractions*, Proc. Amer. Math. Soc. **17** (1966), 843-847.
4. ———, *The strict topology and compactness in the space of measures II*, Trans. Amer. Math. Soc. **126** (1967), 474-486.
5. M. M. Day, *Normed linear spaces*, Springer-Verlag, Berlin, 1958.
6. J. Dieudonné, *Sur la convergence des suites mesures de Radon*, Anais da Acad. Bras. di Ci. **23** (1951), 21-38 and 277-282.
7. ———, *Sur le produit de composition*, Composition Math. **12** (1954), 17-34.
8. R. M. Dudley, *Convergence of Baire measures*, Studia Math. **27** (1966), 251-268.
9. J. Dugundi, *Topology*, Allyn and Bacon, Inc., Boston, 1966.
10. N. Dunford and J. Schwartz, *Linear operators*, Part I. Interscience, New York, 1958.
11. A. Grothendieck, *Sur les applications linéaires faiblement compactes d'espaces du type $C(K)$* , Canad. J. Math. **5** (1953), 129-173.
12. E. Hewitt and K. Stromberg, *Real and abstract analysis*, Springer-Verlag, New York, 1965.
13. J. Isbell and Z. Semadeni, *Projection constants and spaces of continuous functions*, Trans. Amer. Math. Soc. **107** (1963), 38-48.
14. J. L. Kelley, I. Namioka, et al, *Linear topological spaces*, D. Van Nostrand Co., Inc., Princeton, 1963.
15. L. LeCam, *Convergence in distribution of stochastic processes*, Univ. of California Publications in Statistics, **2** (1957), 207-236.
16. V. S. Varadarajan, *Measures on topological spaces*, Amer. Math. Soc. Translations, Series 2, **48** (1965), 161-228.

Received December 11, 1967. This research was done while the author held a Summer Faculty Fellowship from Indiana University

INDIANA UNIVERSITY
BLOOMINGTON, INDIANA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R. PHELPS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. Yosida

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Patrick Robert Ahern, <i>On the geometry of the unit ball in the space of real annihilating measures</i>	1
Kirby Alan Baker, <i>Equational classes of modular lattices</i>	9
E. F. Beckenbach and Gerald Andrew Hutchison, <i>Meromorphic minimal surfaces</i>	17
Tae Ho Choe, <i>Intrinsic topologies in a topological lattice</i>	49
John Bligh Conway, <i>A theorem on sequential convergence of measures and some applications</i>	53
Roger Cuppens, <i>On the decomposition of infinitely divisible probability laws without normal factor</i>	61
Lynn Harry Erbe, <i>Nonoscillatory solutions of second order nonlinear differential equations</i>	77
Burton I. Fein, <i>The Schur index for projective representations of finite groups</i>	87
Stanley P. Gudder, <i>A note on proposition observables</i>	101
Kenneth Kapp, <i>On Croisot's theory of decompositions</i>	105
Robert P. Kaufman, <i>Gap series and an example to Malliavin's theorem</i>	117
E. J. McShane, Robert Breckenridge Warfield, Jr. and V. M. Warfield, <i>Invariant extensions of linear functionals, with applications to measures and stochastic processes</i>	121
Marvin Victor Mielke, <i>Rearrangement of spherical modifications</i>	143
Akio Osada, <i>On unicity of capacity functions</i>	151
Donald Steven Passman, <i>Some $5/2$ transitive permutation groups</i>	157
Harold L. Peterson, Jr., <i>Regular and irregular measures on groups and dyadic spaces</i>	173
Habib Salehi, <i>On interpolation of q-variate stationary stochastic processes</i>	183
Michael Samuel Skaff, <i>Vector valued Orlicz spaces generalized N-functions. I</i>	193
A. J. Ward, <i>On H-equivalence of uniformities. II</i>	207
Thomas Paul Whaley, <i>Algebras satisfying the descending chain condition for subalgebras</i>	217
G. K. White, <i>On subgroups of fixed index</i>	225
Martin Michael Zuckerman, <i>A unifying condition for implications among the axioms of choice for finite sets</i>	233