

Pacific Journal of Mathematics

FIXED-POINT-FREE OPERATOR GROUPS OF ORDER 8

FLETCHER GROSS

FIXED-POINT-FREE OPERATOR GROUPS OF ORDER 8

FLETCHER GROSS

Let A be a group of order 2^n which acts as a fixed-point-free group of operators on the finite solvable group G . If no additional assumptions are made concerning G , then "reasonable" upper bounds on the nilpotent length, $l(G)$, of G have been obtained only when A is cyclic [Gross] or elementary abelian [Shult]. As a small step in extending the class of 2-groups A for which such bounds exist, it is shown in the present paper that if $|A| = 8$, then $l(G) \leq 3$ if A is elementary abelian or quaternion and $l(G) \leq 4$ otherwise.

Unfortunately, the author was unable to generalize his methods of proof to a wider class of groups.

The notation used in this paper agrees with that of [1] with two additions: (1) If G is a linear group operating on V and U is a G -invariant subspace, then $\{G|U\}$ denotes the restriction of G to U ; and (2) $F_0(G) = 1$ and $F_{n+1}(G)/F_n(G)$ is the greatest normal nilpotent subgroup of $G/F_n(G)$.

THEOREM 1. *Let $G = NQ$ be a finite solvable linear group over a field K whose characteristic is not 2 and does not divide $|F_1(N)|$. Assume that N is a normal 2-complement of G and Q is a group of order 8 containing an element x of order 4. If, in addition, $C_N(Q) = 1$ and $\sum_{g \in Q} g = 0$, then it must follow that*

$$[x^2, F_2(N)/F_1(N)] = 1.$$

Proof. According to the hypothesis Q can be any group of order 8 except an elementary abelian group. If Q is cyclic, this theorem is a special case of [4, Th. 1.2], and if Q is a quaternion group, then a stronger result is possible. Thus the main interest in the theorem is when Q is either dihedral or is the direct product of cyclic groups of orders 4 and 2.

To prove the theorem we first notice that extending K affects neither hypothesis or conclusion. Thus we may as well assume that K is algebraically closed. We now assure that G is a minimal counterexample to the theorem and let V be the space on which G operates.

Choose S to be a subgroup of $F_2(N)$ such that Q normalizes S , $[x^2, S] \not\leq F_1(N)$, and S is minimal with respect to the above properties. S must be a p -group for some prime p . Now Q normalizes $[x^2, S]$, and $[x^2, [x^2, S]] = [x^2, S]$ [2]. Due to the minimality of S , this implies that $[x^2, S] = S$.

Now $C_S(O_p(F_1(N))) = S \cap F_1(N)$. Thus there is an r -group R for some prime $r \neq p$ such that QS normalizes R , $R \leq F_1(N)$, $[S, R] \neq 1$, and R is minimal with respect to the above properties. R must be a special r -group, and R/R' must be transformed irreducibly by QS .

Since the characteristic of K does not divide $|F_1(N)|$, V is a completely reducible $K - R$ module. From this and the fact that $[S, R] \triangleleft QSR$, it follows that V contains a maximal $K - QSR$ submodule M such that $[S, R]$ is not the identity on V/M . Now let H be the kernel of the representation of QSR afforded by V/M .

Since $\langle x \rangle$ must be faithfully represented on V/M , we have that either $Q \cap H = 1$ or $Q/Q \cap H$ is cyclic of order 4. But Q has no non-zero fixed vector in V and so certainly has none in V/M . Thus if $Q/Q \cap H$ is cyclic of order 4, then it follows from [4] that $[x^2, S, R] = 1$. Hence we must have $Q \cap H = 1$. This implies that QSR/H acting as a linear group on V/M satisfies the hypothesis but not the conclusion of the theorem. Therefore, in proving the theorem we may as well assume that $G = QSR$ and that V is an irreducible $K - G$ module.

Clifford's theorem now implies that V is a completely reducible $K - SR$ module and $V = V_1 \oplus V_2 \oplus \dots \oplus V_t$ where the V_i are the homogeneous $K - SR$ modules. Q must permute the V_i transitively, and, since $[S, R] \triangleleft QSR$, it must be that $\{[S, R] | V_i\} \neq 1$ for all i .

We now proceed to prove that $t = 1$, or, in other words, that V is a homogeneous $K - SR$ module. For this purpose let

$$Q_i = \{g | g \in Q, V_i g = V_i\}$$

and

$$C_i = \{g | g \in Q_i, \{[g, SR] | V_i\} = 1\} .$$

Then Q_i and Q_j as well as C_i and C_j are conjugate in Q for all i and j . $[Q: Q_i] = t$, V_i is an irreducible $K - Q_i SR$ module, and $\{\sum_{g \in Q_i} g | V_i\} = 0$ for all i . The last fact implies that $Q_i \neq 1$. Since $\{[x^2, S] | V_i\} \neq 1$, x^2 cannot belong to C_i .

LEMMA. $C_i = 1$ for all i .

Proof. Suppose $C_i \neq 1$. Since $\langle x \rangle \cap C_i = 1$, it follows that C_i is cyclic of order 2 generated by an element y_i . Now $C_{RS}(x)$ is normalized by Q . It follows from this and the fact that conjugation by x transitively permutes the y_i that $[u, y_i] = [u, y_j]$ for all i and j and all $u \in C_{RS}(x)$. Since $[u, y_i]$ is represented by the identity on V_i , this all implies that $[C_{RS}(x), y_i] = 1$ for all i . Since x and y_i generate Q , we obtain that $C_{RS}(x) = C_{RS}(Q) = 1$. Hence x acts as a fixed-point-free automorphism on RS . From this follows $[x^2, S, R] = 1$ [3] which is a contradiction.

LEMMA. $Q_i = Q$ and $t = 1$.

Proof. If Q_i is elementary abelian, it follows from [7, Th. 4.1] that $C_i \neq 1$. Thus, since $Q_i \neq 1$, we must have either $Q_i = Q$ or Q_i is cyclic of order 4 generated by an element y . If Q_i is cyclic of order 4 we must have $y^2 = x^2$ because Q only has 8 elements. Now Q_i can have no nonzero fixed vector in V_i . Theorem 1.2 of [4] now yields that $[x^2, S, R]$ is represented by the identity on V_i . Since this is impossible, Q_i must be Q . Then $t = [Q:Q_i] = 1$ and so V is a homogeneous $K - SR$ module.

COROLLARY. $Z(SR) = R' = 1$.

Proof. $Z(SR)$ is represented by scalar matrices and so Q must centralize $Z(SR)$. Thus $Z(SR) \leq C_{RS}(Q) = 1$. Now R' is normalized by QS and so, due to the minimality of R , we must have $[S, R'] = 1$. Therefore $R' \leq Z(SR)$.

Now let $V = U_1 \oplus U_2 \oplus \dots \oplus U_s$ where the U_i are the homogeneous $K - R$ submodules of V . Let $H_i = \{g \mid g \in QS, U_i g = U_i\}$ and $S_i = H_i \cap S$. Now SQ must permute the U_i transitively since V is an irreducible $K - QSR$ module. Thus $s = [QS:H_i]$ for all i . But V is a homogeneous $K - SR$ module. This implies that $(U_i S)Q = U_i S$. Hence $U_i S = V$ for all i . Therefore $s = [S:S_i] = [QS:H_i]$ which means that H_i must contain a Sylow 2-subgroup of SQ . Since the H_i are all conjugate in QS , this implies that $Q \leq H_i$ for some i , $i = 1$ say. Then Q fixes U_1 . Let R_1 be the kernel of the representation of R afforded by U_1 . Clearly R_1 is normalized by Q . But R is abelian and so R is represented by scalar matrices on U_1 . It now follows that $[R/R_1, Q] = 1$. Since $C_N(Q) = 1$, this implies that $R_1 = R$. But, since V is an irreducible $K - QSR$ module and $R \triangleleft QSR$, this is impossible. This contradiction proves the theorem.

THEOREM 2. Let $G = NQ$ be a finite solvable linear group over a field K whose characteristic does not divide $|F_1(N)|$. Assume that N is a normal 2-complement of G and Q is an ordinary quaternion group. If, in addition, $C_N(Q) = 1$ and $\sum_{g \in Q} g = 0$, then it must follow that $[Q', F_1(N)] = 1$.

Proof. Extending K affects neither hypothesis nor conclusion. Thus we assume that K is algebraically closed. If $[Q', F_1(N)] \neq 1$, then there is a subgroup P of $F_1(N)$ such that Q normalizes P , Q' does not centralize P , and P is minimal with respect to the above properties. Then P is a special p -group for some prime p and P/P' is transformed faithfully and irreducibly by Q . This implies that

$|P/P'| = p^2$, and so P is either elementary abelian of order p^2 or extra-special of order p^3 and exponent p .

If V is the vector space on which G operates, then

$$V = V_1 \oplus V_2 \oplus \dots$$

where the V_i are the homogeneous $K - P$ modules. By renumbering, we may assume that $[Q', P]$ is not the identity on V_1 . Now if Q , as a permutation group on the V_i , had an orbit of length 8, then $\sum_{g \in Q} g$ would not be 0. This implies that Q' must fix V_1 .

If $\{P|V_1\}$ is abelian, then P is represented by scalar matrices on V_1 and so we would have $\{[Q', P]|V_1\} = 1$. Thus $\{P|V_1\}$ is not abelian. This implies that $P = \{P|V_1\}$ is an extra-special p -group of order p^3 and exponent p .

Now let $H = \{g | g \in Q, V_1 g = V_1\}$. In order that $\sum_{g \in Q} g = 0$, we must have $\{\sum_{g \in H} g | V_1\} = 0$. Now a faithful irreducible K -representation of P is uniquely determined by the representation of P' [6]. It follows from this that $H = C_Q(P')$. Since $C_P(Q) = 1, H \neq P$. But the automorphism group of P' is cyclic. Thus Q/H is cyclic. This implies that H is cyclic of order 4. Let x generate H and let y be an element of Q not contained in H .

Case 1. $p \equiv 1 \pmod{4}$.

Suppose first that $\text{char}(K) \neq 2$. Then Theorem 3.1 of [7] implies that $\{[x^2, P]|V_1\} = 1$, which is a contradiction. If $\text{char}(K) = 2$, then Theorem B of [6] leads to $\{x^3 + x^2 + x + 1 | V_1\} \neq 0$, also a contradiction.

Case 2. $p \equiv 3 \pmod{4}$.

In this case $GF(p)$ does not contain a primitive 4th root of unity. Since Q faithfully transforms P/P' , it follows that there elements a, b generating P such that

$$a^y \equiv b, b^y \equiv a^{-1} \pmod{P'}$$

But this implies that $[a, b]^y = [b, a^{-1}] = [a, b]$, contrary to $y \in C_Q(P')$.

THEOREM 3. *Let Q be a group of order 8 which acts as a fixed-point-free group of automorphisms of the finite group G . Then G is solvable and $l(G) \leq 3$ if Q is either elementary abelian or a quaternion group and $l(G) \leq 4$ otherwise. The upper bound in the case when Q is elementary abelian or a quaternion group is best-possible.*

Proof. If G admits a 2-group as a fixed-point-free operator group,

then G must have odd order and so G must be solvable from the Feit-Thompson Theorem [1]. If Q is elementary abelian, the result follows from Theorem 4.3 of [7]. Therefore assume that Q has an element x of order 4. We now use induction on $|G|$.

If H_1, H_2 are distinct minimal Q -admissible normal subgroups, then $l(G) \leq l[(G/H) \times (G/H_2)] = \text{Max} \{l(G/H_1), l(G/H_2)\}$. Thus in proving the theorem we may assume that G has only one minimal Q -admissible normal subgroup. Hence $F_1(G)$ is a p -group for some prime p . Now let $N = G/F_1(G)$ and consider NQ as a linear group acting on V where V is $F_1(G)/D(F_1(G))$ written additively. Theorems 1 and 2 imply that $[x^2, F_k(N)/F_{k-1}(N)] = 1$ where $k = 1$ if Q is a quaternion group and $k = 2$ otherwise. It follows from this that $[x^2, N/F_{k-1}(N)] = 1$. But then $N/F_{k-1}(N)$ admits a fixed-point-free operator group of order 4. This implies that $l(N/F_{k-1}(N)) \leq 2$. We now have that

$$l(G) = 1 + l(N) = 1 + (k - 1) + l(N/F_{k-1}(N)) \leq k + 2.$$

Finally, the claim of best-possible in the statement of the theorem is justified by [5].

REFERENCES

1. W. Feit and J. Thompson, *Solvability of groups of odd order*, Pacific J. Math. **13** (1963), 775-1029.
2. G. Glauberman, *Fixed points in groups with operator groups*, Math. Z. **84** (1964), 120-125.
3. D. Gorenstein and I. Herstein, *Finite groups admitting a fixed-point-free automorphism of order 4*, Amer. J. Math. **83** (1961), 71-78.
4. F. Gross, *Groups admitting a fixed-point-free automorphism of order 2^n* , Pacific J. Math. (in press)
5. ———, *A note on fixed-point-free solvable operator groups*, Proc. Amer. Math. Soc. (in press).
6. P. Hall and G. Higman, *On the p -length of p -soluble groups and reduction theorems for Burnside's problem*, Proc. London Math. Soc. (3) **6** (1956), 1-42.
7. E. Shult, *On groups admitting fixed-point-free abelian groups*, Illinois J. Math. **9** (1965), 701-720.

Received March 19, 1968.

UNIVERSITY OF UTAH
SALT LAKE CITY, UTAH

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R. PHELPS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics

Vol. 28, No. 2

April, 1969

Richard Arens and Donald George Babbitt, <i>The geometry of relativistic n-particle interactions</i>	243
Kirby Alan Baker, <i>Hypotopological spaces and their embeddings in lattices with Birkhoff interval topology</i>	275
J. Lennart (John) Berggren, <i>Finite groups in which every element is conjugate to its inverse</i>	289
Beverly L. Brechner, <i>Homeomorphism groups of dendrons</i>	295
Robert Ray Colby and Edgar Andrews Rutter, <i>QF - 3 rings with zero singular ideal</i>	303
Stephen Daniel Comer, <i>Classes without the amalgamation property</i>	309
Stephen D. Fisher, <i>Bounded approximation by rational functions</i>	319
Robert Gaines, <i>Continuous dependence for two-point boundary value problems</i>	327
Bernard Russel Gelbaum, <i>Banach algebra bundles</i>	337
Moses Glasner and Richard Emanuel Katz, <i>Function-theoretic degeneracy criteria for Riemannian manifolds</i>	351
Fletcher Gross, <i>Fixed-point-free operator groups of order 8</i>	357
Sav Roman Harasymiv, <i>On approximation by dilations of distributions</i>	363
Cheong Seng Hoo, <i>Nilpotency class of a map and Stasheff's criterion</i>	375
Richard Emanuel Katz, <i>A note on extremal length and modulus</i>	381
H. L. Krall and I. M. Sheffer, <i>Difference equations for some orthogonal polynomials</i>	383
Yu-Lee Lee, <i>On the construction of lower radical properties</i>	393
Robert Phillips, <i>Liouville's theorem</i>	397
Yum-Tong Siu, <i>Analytic sheaf cohomology groups of dimension n of n-dimensional noncompact complex manifolds</i>	407
Michael Samuel Skaff, <i>Vector valued Orlicz spaces. II</i>	413
James DeWitt Stein, <i>Homomorphisms of B^*-algebras</i>	431
Mark Lawrence Teply, <i>Torsionfree injective modules</i>	441
Richard R. Tucker, <i>The δ^2-process and related topics. II</i>	455
David William Walkup and Roger Jean-Baptiste Robert Wets, <i>Lifting projections of convex polyhedra</i>	465
Thomas Paul Whaley, <i>Large sublattices of a lattice</i>	477