

Pacific Journal of Mathematics

DIFFERENCE EQUATIONS FOR SOME ORTHOGONAL POLYNOMIALS

H. L. KRALL AND I. M. SHEFFER

DIFFERENCE EQUATIONS FOR SOME ORTHOGONAL POLYNOMIALS

H. L. KRALL AND I. M. SHEFFER

It is well-known that every orthogonal polynomial set $\{P_n(x)\}$ satisfies a 3-term recurrence relation of the form

$$(1.1) \quad P_{n+1}(x) = (a_n x + b_n)P_n(x) + c_n P_{n-1}(x) \quad (n = 1, 2, \dots).$$

Some orthogonal sets (polynomials of Jacobi, Hermite and so on) are solutions of differential equations. It will be shown that there exist orthogonal polynomial sets that satisfy 3-term difference equations of the form

$$(1.2) \quad A(x)y(x + \alpha) + B(x)y(x - \alpha) + C(x)y(x) = \lambda y(x)$$

where A, B, C are polynomials of degree ≤ 2 and λ is a parameter.

Consider the difference equation

$$(1.3) \quad A(x)y(x + \alpha) + B(x)y(x + \beta) + C(x)y(x) = \lambda y(x)$$

where A, B, C are real polynomials, λ is a parameter, and $\alpha, \beta, 0$ are distinct and real. We examine two cases, according as A, B, C are of degree ≤ 1 :

(a) $A(x) = a_1 x + a_0, B(x) = b_1 x + b_0, C(x) = c_1 x,$
 or are of degree ≤ 2 :

(b) $A(x) = a_2 x^2 + a_1 x + a_0, B(x) = b_2 x^2 + b_1 x + b_0, C(x) = c_2 x^2 + c_1 x$
 (a_2, b_2, c_2 not all zero). We shall use the notation (1.3a), (1.3b) to denote equation (1.3) for the respective conditions (a), (b).

Equation (1.3) will be termed *admissible* if there exists a real sequence $\{\lambda_n\}$ ($n = 0, 1, \dots$) such that for $\lambda = \lambda_n$ there is a polynomial solution $y_n(x)$, unique to within a multiplicative constant, and $y_n(x)$ is of degree (exactly) n . It follows that admissibility implies that

$$(1.4) \quad \lambda_m \neq \lambda_n (m \neq n).$$

LEMMA 1.1. *Equation (1.3a) is admissible if and only if*

$$(1.5) \quad a_1 + b_1 + c_1 = 0, \quad \delta \equiv a_1 \alpha + b_1 \beta \neq 0.$$

And in this case we have

$$(1.6) \quad \lambda_n = (a_0 + b_0) + n(a_1 \alpha + b_1 \beta) \quad (n = 0, 1, \dots).$$

Proof. Let n be arbitrary. If we substitute the n th degree polynomial

$$(1.7) \quad y(x) = x^n + \sum_{j=0}^{n-1} p_j x^j$$

into (1.3a), a necessary and sufficient condition that $y(x)$ be a solution is that coefficients of x^{n+1} , x^n , \dots agree on both sides. The coefficient of x^{n+1} yields the first of (1.5), that of x^n gives $\lambda = \lambda_n$ as in (1.6); and those of x^{n-1} , \dots , x^0 give successive equations for p_{n-1} , \dots , p_0 . In these equations the coefficients of p_{n-1} , \dots , p_0 are respectively $\lambda_n - \lambda_{n-1}$, $\lambda_n - \lambda_{n-2}$, \dots , $\lambda_n - \lambda_0$, so there is one and only one choice of the p_j 's if and only if $\lambda_n \neq \lambda_j$ ($j \leq n-1$). This condition is equivalent to the second part of (1.5); and the lemma is established.

LEMMA 1.2. *Equation (1.3b) is admissible if and only if*

$$(1.8) \quad a_2 + b_2 + c_2 = 0, \quad a_1 + b_1 + c_1 = 0, \quad a_2\alpha + b_2\beta = 0;$$

$$(1.9) \quad 2(a_1\alpha + b_1\beta) + n(a_2\alpha^2 + b_2\beta^2) \neq 0 \quad (n = 0, 1, \dots).$$

And in this case λ_n is given by

$$(1.10) \quad \lambda_n = (a_0 + b_0) + n(a_1\alpha + b_1\beta) + n(n-1)(a_2\alpha^2 + b_2\beta^2)/2 \\ (n = 0, 1, \dots).$$

Proof. Substituting (1.7) into (1.3b) and equating like terms (as a necessary and sufficient condition for a solution) we find that the terms in x^{n+2} , x^{n+1} give (1.8), the x^n term gives $\lambda = \lambda_n$ as in (1.10), and p_{n-1} , \dots , p_0 again are uniquely determined if and only if $\lambda_n \neq \lambda_j$ ($j \leq n-1$). Now the condition $\lambda_m \neq \lambda_n$ ($m \neq n$) is seen to reduce to (1.9); so the lemma is proved.

In the proofs of Lemmas 1.1, 1.2 it was seen that if a polynomial $y(x)$ of degree n satisfies (1.3a or b) then the corresponding value of λ is λ_n as given by (1.6) or (1.10); so we have the

COROLLARY. *If (1.3a) or (1.3b) is admissible then for each $\lambda \neq \lambda_n$ ($n = 0, 1, \dots$) the only polynomial solution is $y(x) \equiv 0$.*

Let (1.3a) or (1.3b) be admissible. In both cases the solution for $n = 1$ is

$$(1.11) \quad y_1(x) = x + (a_0\alpha + b_0\beta)\delta^{-1}$$

where δ is given in (1.5). If we set

$$x + d = x^*, \quad z(x^*) = y(x^* - d)$$

with

$$d = (a_0\alpha + b_0\beta)\delta^{-1},$$

the equation in $z(x^*)$ will also be admissible and will have the form (1.3a) or (1.3b) after the constant term in $C(x^*)$ has been absorbed into the λ . Moreover, for $n = 1$ we have

$$z_1(x^*) = x^* .$$

An admissible equation (1.3a) or (1.3b) in which for $n = 1$ the solution contains no constant term will be called *canonical*. It is no restriction to limit ourselves to canonical equations.

From (1.11) we obtain

LEMMA 1.3. *The admissible equation (1.3a) or (1.3b) is canonical if and only if*

$$(1.12) \quad a_0\alpha + b_0\beta = 0 .$$

2. **Orthogonality for case (1.3a).** We consider the problem of determining those canonical equations (1.3a) [(1.3b) in § 3] whose polynomial solutions form an orthogonal set. For all polynomials $y(x)$ we have

$$(2.1) \quad y(x + u) = \sum_{k=0}^{\infty} y^{(k)}(x)u^k/k !$$

so (1.3a) is equivalent, with respect to polynomial solutions, to the differential equation of infinite order

$$(2.2) \quad xy'(x) + \sum_{k=2}^{\infty} H_k(x)y^{(k)}(x)/k! = \sigma y(x)$$

where

$$(2.3) \quad H_k(x) = r_k + s_kx = (a_0\alpha^k + b_0\beta^k)\delta^{-1} + (a_1\alpha^k + b_1\beta^k)\delta^{-1}x$$

($k = 1, 2, \dots$) with $\sigma = \{\lambda - (a_0 + b_0)\}\delta^{-1}$. Using (1.6) we find that the sequence $\{\sigma_n\}$ for which there are polynomial solutions is given by $\sigma_n = n$.

Equation (2.2) is identical with equation (3.1) of [1]. In Remark (i) ([1], p. 151) it is shown that if $r_2 = 0$ the polynomial solutions do not form an orthogonal set. We therefore assume $r_2 \neq 0$. In this case, Theorem 3.1 ([1], p. 151) states that the solutions of (our present) equation (2.2), hence of cononical equation (1.3a), form a weak orthogonal set if and only if

$$(2.4) \quad \begin{aligned} r_{2p+1} &= 0, & s_{2p+1} &= s_3^p, \\ r_{2p+2} &= r_2s_3^p, & s_{2p+2} &= s_2s_3^p \quad (p = 0, 1, \dots) . \end{aligned}$$

Moreover the weak orthogonal set is an orthogonal set when and only when one of the following two relations holds:

$$(2.5_1) \quad s_2^2 - s_3 = 0 ;$$

$$(2.5_2) \quad s_2^2 - s_3 \neq 0 \quad \text{and} \quad 2r_2(s_2^2 - s_3)^{-1} \neq 0, 1, 2, \dots .$$

The condition $r_{2p+1} = 0$ is

$$(2.6) \quad a_0\alpha^{2p+1} + b_0\beta^{2p+1} = 0 \quad (p = 0, 1, \dots) .$$

If $a_0 = 0$ or $b_0 = 0$ then both are zero since $\alpha\beta \neq 0$. But then $r_2 = 0$, contrary to assumption. So $a_0b_0 \neq 0$. Taking $p = 0, 1$ in (2.6) we then get $\beta^2 = \alpha^2$. Since α, β are distinct, then $\beta = -\alpha$; and again from (2.6) with $p = 0$: $a_0 = b_0$. Thus, if $r_2 \neq 0$ then $r_{2p+1} = 0$ ($p = 0, 1, \dots$) if and only if

$$(2.7) \quad \beta = -\alpha, a_0 = b_0 \neq 0 .$$

With (2.7) holding then

$$\delta = \alpha(a_1 - b_1) \neq 0 ,$$

so

$$(2.8) \quad \begin{aligned} r_{2p+1} &= 0, s_{2p+1} = \alpha^{2p}, r_{2p+2} = 2a_0(a_1 - b_1)^{-1}\alpha^{2p+1}, \\ s_{2p+2} &= (a_1 + b_1)(a_1 - b_1)^{-1}\alpha^{2p+1} . \end{aligned}$$

Conditions (2.4) are seen to be satisfied. And (2.5₁), (2.5₂) become respectively:

$$(2.9_1) \quad a_1b_1 = 0 ;$$

$$(2.9_2) \quad a_1b_1 \neq 0, a_0(a_1 - b_1)(\alpha a_1 b_1)^{-1} \neq 0, 1, \dots .$$

To sum up:

THEOREM 2.1. *Let equation (1.3a) be canonical. Then its polynomial solutions form an orthogonal set if and only if (2.7) holds and one of (2.9₁), (2.9₂) holds.*

REMARKS. (i) If (1.3a) is canonical its polynomial solutions form an orthogonal set if and only if it is of the form

$$(2.10) \quad \begin{aligned} (a_1x + a_0)y(x + \alpha) + (b_1x + a_0)y(x - \alpha) \\ - (a_1 + b_1)xy(x) = \lambda y(x) , \end{aligned}$$

with $a_0 \neq 0, a_1 \neq b_1, \alpha \neq 0$, and either (2.9₁) or (2.9₂) holding.

(ii) In (2.10) make the variable changes $x = \alpha x^*, z(x^*) = y(\alpha x^*)$. There results a similar difference equation in $z(x^*)$, in which α is replaced by 1. This equation has an orthogonal set of solutions when (2.10) does. It may be termed a *standard* canonical equation. After

dividing by a_0 this equation has the form (dropping asterisks)

$$(2.11) \quad \begin{aligned} (c_1x + 1)z(x + 1) + (d_1x + 1)z(x - 1) \\ - (c_1 + d_1)xz(x) = \mu z(x), \end{aligned}$$

with $c_1 - d_1 \neq 0$ and either $c_1d_1 = 0$ or

$$c_1d_1 \neq 0, (c_1 - d_1)(c_1d_1)^{-1} \neq 0, 1, 2, \dots$$

3. Orthogonality for case (1.3b). Let equation (1.3b) be canonical, so that (1.12) holds. Putting (2.1) into (1.3b) we get an infinite order differential equation with polynomial coefficients of degree ≤ 2 , which is equivalent to (1.3b) at least for polynomial solutions:

$$(3.1) \quad xy'(x) + \sum_{k=2}^{\infty} T_k(x)y^{(k)}(x)/k! = \sigma y(x),$$

where

$$(3.2) \quad \begin{aligned} T_k(x) = r_k + s_kx + t_kx^2 = (a_0\alpha^k + b_0\beta^k)\delta^{-1} + (a_1\alpha^k + b_1\beta^k)\delta^{-1}x \\ + (a_2\alpha^k + b_2\beta^k)\delta^{-1}x^2 \quad (k = 2, 3, \dots) \end{aligned}$$

and $\sigma = \{\lambda - (a_0 + b_0)\}\delta^{-1}$ and δ is given by (1.5). From (1.10) we see that $\{\sigma_n\}$ is given by

$$\sigma_n = n + n(n - 1)t_2/2.$$

Equations of the form (3.1), that is, with $\max_k \{\text{degree } T_k(x)\} = 2$ were considered in [1], but the results obtained were not as complete as for the case where the coefficients are of degree ≤ 1 . We must therefore proceed differently. We first show that if canonical equation (1.3b), hence also (3.1), has an orthogonal set of solutions then $\beta = -\alpha$.

For suppose not. Then $|\alpha| \neq |\beta|$, since α, β are distinct. We may assume that $|\alpha| > |\beta|$. By Theorem 2.2 ([1], p. 148) there is a sequence of constants $\{\alpha_n\}$ (the moments of the weight function corresponding to the orthogonal set), with $\alpha_0 \neq 0$, that satisfies the system of equations

$$(3.3) \quad d_{p+k}^p = 0, D_{p+k}^p = 0 \quad (p, k = 0, 1, \dots)$$

where (in our present case, as seen in [1], p. 153)

$$(3.4) \quad \begin{aligned} d_{p+k}^p = \sum_{i=k}^{2k+2} \alpha_i \left[\binom{k}{i-k} r_{2p+2k+1-i} + \binom{k}{i-k-1} s_{2p+2k+2-i} \right. \\ \left. + \binom{k}{i-k-2} t_{2p+2k+3-i} \right], \end{aligned}$$

$$\begin{aligned}
 D_{p+k}^p &= \sum_{i=k}^{2k+3} \alpha_i \left[\frac{i+1}{k+1} \binom{k+1}{i-k} r_{2p+2k+2-i} \right. \\
 (3.5) \quad &+ \frac{i}{k+1} \binom{k+1}{i-k-1} s_{2p+2k+3-i} \\
 &\left. + \frac{i-1}{k+1} \binom{k+1}{i-k-2} t_{2p+2k+4-i} \right].
 \end{aligned}$$

Here the convention is made that $\binom{m}{q} = 0$ for $q < 0$, and $r_j = s_j = t_j = 0$ for $j \leq 0$ and $r_1 = t_1 = 0, s_1 = 1$.

Putting the values of r_k, s_k, t_k from (3.2) into (3.3) we get

$$\begin{cases} \alpha^{2p+2k+1} U_k + \beta^{2p+2k+1} V_k = 0 \\ \alpha^{2p+2k+2} W_k + \beta^{2p+2k+2} X_k = 0 \end{cases} \quad (p, k = 0, 1, \dots)$$

where

$$\begin{aligned}
 U_k &= \sum_{i=k}^{2k+2} \alpha_i \left[\binom{k}{i-k} a_0 \alpha^{-i} + \binom{k}{i-k-1} a_1 \alpha^{-i+1} \right. \\
 (3.7) \quad &\left. + \binom{k}{i-k-2} a_2 \alpha^{-i+2} \right], \\
 W_k &= \sum_{i=k}^{2k+3} \alpha_i \left[\frac{i+1}{k+1} \binom{k+1}{i-k} a_0 \alpha^{-i} + \frac{i}{k+1} \binom{k+1}{i-k-1} a_1 \alpha^{-i+1} \right. \\
 &\left. + \frac{i-1}{k+1} \binom{k+1}{i-k-2} a_2 \alpha^{-i+2} \right],
 \end{aligned}$$

and V_k, X_k are obtained from U_k, W_k by replacing

$$a_0, a_1, a_2, \alpha \quad \text{by} \quad b_0, b_1, b_2, \beta.$$

Let k be arbitrary but fixed. If we divide (3.6) by $\alpha^{2p+2k+1}, \alpha^{2p+2k+2}$ respectively and let $p \rightarrow \infty$, then since $|\beta/\alpha| < 1$ we get

$$(3.8) \quad U_k = 0, W_k = 0 \quad (k = 0, 1, \dots).$$

And from (3.6) we then have

$$(3.9) \quad V_k = 0, X_k = 0 \quad (k = 0, 1, \dots).$$

For $k = 0$, (3.8), (3.9) reduce to

$$\begin{aligned}
 (3.10) \quad &\alpha_0 a_0 + \alpha_1 a_1 + \alpha_2 a_2 = 0, \alpha_1 a_0 + \alpha_2 a_1 + \alpha_3 a_2 = 0, \\
 &\alpha_0 b_0 + \alpha_1 b_1 + \alpha_2 b_2 = 0, \alpha_1 b_0 + \alpha_2 b_1 + \alpha_3 b_2 = 0.
 \end{aligned}$$

Now from (3.3) with $p = k = 0$ we have

$$\alpha_0 r_1 + \alpha_1 s_1 + \alpha_2 t_1 = 0.$$

But $r_1 = t_1 = 0, s_1 = 1$; hence

$$\alpha_1 = 0 .$$

So (3.10) becomes

$$(3.11) \quad \begin{aligned} \alpha_0 a_0 + \alpha_2 a_2 = 0, \alpha_2 a_1 + \alpha_3 a_2 = 0, \\ \alpha_0 b_0 + \alpha_2 b_2 = 0, \alpha_2 b_1 + \alpha_3 b_2 = 0 . \end{aligned}$$

Now $a_2 b_2 \neq 0$. For if a_2 or b_2 is zero then from $a_2 \alpha + b_2 \beta = 0$ (in (1.8)) and $\alpha \beta \neq 0$ we get $a_2 = b_2 = 0$. Hence (again from (1.8)) $c_2 = 0$; so all coefficients in (1.3b) are of degree < 2 , contrary to assumption. Again, $a_0 b_0 \neq 0$. For if a_0 or b_0 is zero then (3.11) implies that $\alpha_2 = 0$. Since we already have $\alpha_1 = 0$, then $\Delta_1 = \begin{vmatrix} \alpha_0 \alpha_1 \\ \alpha_1 \alpha_2 \end{vmatrix} = 0$. But for the moments $\{\alpha_n\}$ corresponding to an orthogonal set it is known [2] that

$$\Delta_n \equiv \begin{vmatrix} \alpha_0 \alpha_1 \cdots \alpha_n \\ \alpha_1 \alpha_2 \cdots \alpha_{n+1} \\ \dots \dots \dots \\ \alpha_n \alpha_{n+1} \cdots \alpha_{2n} \end{vmatrix} \neq 0 \quad (n = 0, 1, \dots) ;$$

so we have a contradiction. Thus,

$$(3.12) \quad a_2 b_2 \neq 0, a_0 b_0 \neq 0, \alpha_2 \neq 0 .$$

The right hand equations in (3.11) give us

$$-b_1 a_2 + a_1 b_2 = 0 .$$

This with

$$\alpha a_2 + \beta b_2 = 0$$

from (1.8) implies

$$\alpha a_1 + \beta b_1 = 0 ,$$

contrary to (1.9) for $n = 0$. So the assumption $\beta \neq -\alpha$ leads to a contradiction, and we have

$$(3.13) \quad \beta = -\alpha .$$

Then from (1.12):

$$(3.14) \quad a_0 = b_0 .$$

In (3.2) we now have

$$(3.15) \quad \begin{cases} r_{2p} = 2a_0 \delta^{-1} \alpha^{2p}, s_{2p} = (a_1 + b_1) \delta^{-1} \alpha^{2p}, t_{2p} = (a_2 + b_2) \delta^{-1} \alpha^{2p} \\ r_{2p+1} = 0, s_{2p+1} = (a_1 - b_1) \delta^{-1} \alpha^{2p+1}, t_{2p+1} = (a_2 - b_2) \delta^{-1} \alpha^{2p+1} \end{cases}$$

($p = 1, 2, \dots$), with $r_1 = t_1 = 0, s_1 = 1$. (3.12) and (3.15) show that

$$r_2 \neq 0.$$

Let

$$(3.16) \quad u_p = s_{2p+1}, v_p = t_{2p+1}, w_p = t_{2p+2}.$$

From $a_2\alpha + b_2\beta = 0, \beta = -\alpha \neq 0$ we get

$$(3.17) \quad a_2 = b_2.$$

It is then readily seen that

$$(3.18) \quad v_p = 0, w_p - t_2 u_p = 0 \quad (p = 0, 1, 2, \dots)$$

Choose r_2, s_2, t_2, s_3 to satisfy the conditions

$$(3.19) \quad r_2 \neq 0, 2 + kt_2 \neq 0 (k = 0, 1, \dots), \Delta_2 \neq 0,$$

where $\alpha_1 = 0, \alpha_2, \alpha_3, \alpha_4$ are obtained from the equations

$$(3.20) \quad D_0^0 = 0, d_1^0 = 0, D_1^0 = 0.$$

(3.18)–(3.20) make Theorem 4.2 ([1], p. 158) applicable, so that the solutions of (1.3b) form a weak orthogonal set if and only if

$$(3.21) \quad \begin{cases} s_{2p+1} = s_3^p, t_{2p+1} = 0, r_{2p+1} = 0 \\ s_{2p+2} = s_2 s_3^p, t_{2p+2} = t_2 s_3^p, r_{2p+2} = r_2 s_3^p \end{cases} \quad (p = 0, 1, \dots).$$

Now these conditions do hold in view of (3.14).

The first two conditions of (3.19) become

$$(3.22) \quad a_0 \neq 0; (a_1 - b_1) + k\alpha a_2 \neq 0 \quad (k = 0, 1, 2, \dots).$$

Finally, for weak orthogonality to imply orthogonality it is necessary and sufficient ([1], pp. 161–162) that $t_2 \notin S(r_2, s_2, s_3)$ where $S(r_2, s_2, s_3)$ is the set of all real values of t_2 for which $\pi_n(r_2, s_2, s_3, t_2) = 0$ for some $n > 1$. The expression for π_n is lengthy, and we do not reproduce it here. We merely observe that for given r_2, s_2, s_3 the set $S(r_2, s_2, s_3)$ is at most denumerable.

To sum up:

THEOREM 3.1. *Let the admissible equation (1.3b) be canonical. Its solutions form an orthogonal polynomial set if and only if:*

- (i) (3.12), (3.13), (3.14), (3.17), (3.19) hold.
- (ii) $t_2 \notin S(r_2, s_2, s_3)$.

REMARKS. (a) If the canonical equation (1.3b) has an orthogonal polynomial set of solutions then it has the form

$$(3.23) \quad (a_2x^2 + a_1x + a_0)y(x + \alpha) + (a_2x^2 + b_1x + a_0)y(x - \alpha) - [2a_2x^2 + (a_1 + b_1)x]y(x) = \lambda y(x) ,$$

with

$$(3.24) \quad a_0a_2(a_1 - b_1)\alpha \neq 0; (a_1 - b_1) + k\alpha a_2 \neq 0 \quad (k = 0, 1, \dots) .$$

(b) As in § 2 the transformation $x = \alpha x^*, z(x^*) = y(\alpha x^*)$ carries (3.24) into a similar equation with α replaced by 1.

4. **Two examples.** If an orthogonal polynomial set $\{P_n(x)\}$ satisfies (2.10) with $\lambda = \lambda_n$ for $y = P_n(x)$ then from (1.6) we have

$$(4.1) \quad \lambda_n = 2a_0 + n\alpha(a_1 - b_1) \quad (n = 0, 1, \dots) .$$

Let $\{P_n(x)\}, \{Q_n(x)\}$ be polynomial sets defined by the respective generating functions

$$(4.2) \quad e^{ct}(1 - t)^{x+c} = \sum_{n=0}^{\infty} P_n(x)t^n \quad (c \neq 0) ,$$

$$(4.3) \quad (1 - t)^{x-bd} \cdot (1 - bt)^{-x+d} = \sum_{n=0}^{\infty} Q_n(x)t^n \quad (b \neq 0, 1) .$$

We shall show that these sets are orthogonal and satisfy an equation of the form (2.10).

Denote the left side of (2.10) by $L[y]$. If $G(x, t)$ is the generating function in (4.2) then

$$(4.4) \quad L[G] = G\{(a_1x + a_0)(1 - t)^\alpha + (b_1x + a_0)(1 - t)^{-\alpha} - (a_1 + b_1)x\} .$$

Also,

$$(4.5) \quad \sum_{n=0}^{\infty} \lambda_n P_n(x)t^n = 2a_0G + \alpha(a_1 - b_1)t \partial G/\partial t = G\{2a_0 + \alpha(a_1 - b_1)t[c - (x + c)(1 - t)^{-1}]\} .$$

$\{P_n(x)\}$ will satisfy (2.10) if (4.4) and (4.5) are identical. It is a straightforward computation to show that they are identical if

$$(4.6) \quad \alpha = 1; a_1 = 0; b_1 = a_0/c .$$

Hence $\{P_n(x)\}$ is an orthogonal set which satisfies the equation

$$(4.7) \quad P_n(x + 1) + (x + c)P_n(x - 1) - xP_n(x) = (2c - n)P_n(x) .$$

In the same way it is found that $\{Q_n(x)\}$ is an orthogonal set that is a solution of (2.10) for

$$(4.8) \quad \alpha = 1; a_1 = bb_1; a_0 = -bdb_1 .$$

The equation reduces to

$$(4.9) \quad \begin{aligned} b(x-d)Q_n(x+1) + (x-bd)Q_n(x-1) - (b+1)xQ_n(x) \\ = \{-2bd + n(b-1)\}Q_n(x). \end{aligned}$$

In the case of (4.9) the condition (2.9₂) is to hold. It reduces to

$$(4.10) \quad -d(b-1) \neq 0, 1, 2, \dots$$

REFERENCES

1. H. L. Krall and I. M. Sheffer, *Differential equations of infinite order for orthogonal polynomials*, *Annali di Matematica* (4) **74** (1966), 135-172.
2. G. Szegő, *Orthogonal polynomials*, American Mathematical Society Colloquium Publications, vol. 23, 1959.

Received March 12, 1968.

PENNSYLVANIA STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R. PHELPS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Richard Arens and Donald George Babbitt, <i>The geometry of relativistic n-particle interactions</i>	243
Kirby Alan Baker, <i>Hypotopological spaces and their embeddings in lattices with Birkhoff interval topology</i>	275
J. Lennart (John) Berggren, <i>Finite groups in which every element is conjugate to its inverse</i>	289
Beverly L. Brechner, <i>Homeomorphism groups of dendrons</i>	295
Robert Ray Colby and Edgar Andrews Rutter, <i>QF – 3 rings with zero singular ideal</i>	303
Stephen Daniel Comer, <i>Classes without the amalgamation property</i>	309
Stephen D. Fisher, <i>Bounded approximation by rational functions</i>	319
Robert Gaines, <i>Continuous dependence for two-point boundary value problems</i>	327
Bernard Russel Gelbaum, <i>Banach algebra bundles</i>	337
Moses Glasner and Richard Emanuel Katz, <i>Function-theoretic degeneracy criteria for Riemannian manifolds</i>	351
Fletcher Gross, <i>Fixed-point-free operator groups of order 8</i>	357
Sav Roman Harasymiv, <i>On approximation by dilations of distributions</i>	363
Cheong Seng Hoo, <i>Nilpotency class of a map and Stasheff's criterion</i>	375
Richard Emanuel Katz, <i>A note on extremal length and modulus</i>	381
H. L. Krall and I. M. Sheffer, <i>Difference equations for some orthogonal polynomials</i>	383
Yu-Lee Lee, <i>On the construction of lower radical properties</i>	393
Robert Phillips, <i>Liouville's theorem</i>	397
Yum-Tong Siu, <i>Analytic sheaf cohomology groups of dimension n of n-dimensional noncompact complex manifolds</i>	407
Michael Samuel Skaff, <i>Vector valued Orlicz spaces. II</i>	413
James DeWitt Stein, <i>Homomorphisms of B^*-algebras</i>	431
Mark Lawrence Teply, <i>Torsionfree injective modules</i>	441
Richard R. Tucker, <i>The δ^2-process and related topics. II</i>	455
David William Walkup and Roger Jean-Baptiste Robert Wets, <i>Lifting projections of convex polyhedra</i>	465
Thomas Paul Whaley, <i>Large sublattices of a lattice</i>	477