

# Pacific Journal of Mathematics

**AUTOMORPHISMS OF GROUPS OF SIMILITUDES OVER  $F_3$**

JON F. CARLSON

## AUTOMORPHISMS OF GROUPS OF SIMILITUDES OVER $F_3$

J. F. CARLSON

**Let  $Q(x)$  be a quadratic form defined on a vector space  $M$  of dimension  $n = 2m > 4$  ( $n \neq 8$ ) over the field with three elements. The purpose of this paper is to show that any automorphism of the projective group of similitudes or of the projective group of proper similitudes can not take the coset of an  $(n - 2, 2)$  involution into the coset of an  $(n - p, p)$  involution or into the coset of a similitude  $T$  of ratio  $\rho$  where  $T^2 = \rho_L$  (left multiplication by  $\rho$ ) and where  $\rho$  is not a square in  $F_3$ . This result, together with some results of Wonenburger, shows that any such automorphism is induced by an automorphism of the group of similitudes.**

In [6] Wonenburger proved the above statements except when the field  $K$  had 3 or 5 elements. Only the corollary to Lemma 3 of [6] needs to be verified to complete this result for all fields of characteristic not 2. The case in which  $K = F_5$  can be proved by a slight alteration of Wonenburger's argument, so it will not be considered in this paper.

A linear transformation  $T$  of a vector space  $M$  of dimension  $n = 2m$  is called a similitude of ratio  $\rho$  ( $0 \neq \rho \in K$ ), of the quadratic form  $Q$ , if for every  $x \in M$ ,  $Q(xT) = \rho Q(x)$ . The similitudes of ratio 1 make up the orthogonal group  $O(Q)$ . A similitude  $T$  of ratio  $\rho$  is called a proper similitude if the determinant of the matrix of  $T$ ,  $\text{Det } T = \rho^m$ , and improper if  $\text{Det } T = -\rho^m$ . Let  $S(Q)$  ( $S^+(Q)$ ) denote the group of similitudes (proper similitudes). In the projective groups  $PS(Q) = S(Q)/K^*$  and  $PS^+(Q) = S^+(Q)/K^*$ , we will use the symbol  $\bar{T}$  to denote the coset of the similitude  $T$ .

A projective involution is a similitude  $T$  such that  $\bar{T}^2 = \bar{1}$ , the coset of the identity. There are three types of projective involutions (see [6, p. 608]). First there are scalar multiples of orthogonal involutions. With respect to an orthogonal involution  $T$ , we can decompose  $M$  into a "minus space"  $M^-$  of dimension  $p$  such that  $xT = -x$  for all  $x \in M^-$ , and a "plus space"  $M^+$  of dimension  $n - p$  such that  $xT = x$  for all  $x \in M^+$ . Such a transformation will be called an  $(n - p, p)$  involution and its image in  $PS(Q)$  or in  $PS^+(Q)$  will be called an  $(n - p, p)$  coset. A  $P$ -involution is a similitude  $T$  of ratio  $\rho$  with  $T^2 = -\rho_L$ , left multiplication by  $-\rho$ . All  $P$ -involutions are proper similitudes (see [3, p. 64]). A  $P'$ -involution is a similitude  $T$  of ratio  $\rho$  such that  $T^2 = \rho_L$ ,  $\rho$  not a square in  $K$ . If  $K = F_3$ , it can be easi-

ly shown that the determinant of the matrix of a  $P'$ -involution is 1. Therefore a  $P'$ -involution is proper if and only if the dimension of  $M$  is divisible by 4; i.e., if and only if  $(-1)^m = 1$ .

**THEOREM.** *Let  $M$  be a vector space of dimension  $n = 2m > 4$  over the field  $K = F_3$ , and let  $Q$  be a nondegenerate quadratic form on  $M$ . No automorphism of  $PS(Q)$  or of  $PS^+(Q)$  can take an  $(n - 2, 2)$  coset into an  $(n - p, p)$  coset,  $p \neq 2, n - 2$ , or into the coset of a  $P'$ -involution.*

The proof is by comparing the orders of the centralizers of these projective involutions. First we need two lemmas. It shall be assumed that the hypotheses of the theorem hold.

**LEMMA 1.** *If  $\bar{T}$  is an  $(n - p, p)$  coset and  $p = 2r$ , then the highest power of the prime 3 dividing the order of the centralizer of  $\bar{T}$  in  $PS(Q)$  or in  $PS^+(Q)$  is*

$$(p(p - 2) + (n - p)(n - p - 2))/4 .$$

*Proof.* The centralizer of  $\bar{T}$  in  $PS^+(Q)$ , which we shall denote by  $C_{PS^+}(\bar{T})$ , contains all cosets of elements of  $0^+(Q^+) \times 0^+(Q^-)$ , where  $0^+(Q^+)(0^+(Q^-))$  is the group of all proper orthogonal transformations of the plus (minus) space of  $T$ . Since  $F_3$  has only two nonzero elements, the index  $[S^+(Q^+):0^+(Q^+)] = 2$ . Hence the order

$$[PS^+(Q^+):1] = [0^+(Q^+):1] .$$

An element  $\bar{U}$  will be contained in  $C_{PS^+}(\bar{T})$  if and only if  $U$  leaves invariant both the plus and minus spaces of  $T$ , except in the case in which  $p = n - p$  and the discriminant of  $Q^+$ , the restriction of  $Q$  to the plus space of  $T$ , equals the discriminant of  $Q^-$ . In this latter case there will exist similitudes which will interchange the plus and minus spaces of  $T$ . The number of these elements will equal the number of similitudes which leave invariant both the plus and minus spaces. Therefore we have  $[C_{PS}(\bar{T}):1] = 2[C_{PS^+}(\bar{T}):1] = k[0^+(Q^+):1] \cdot [0^+(Q^-):1]$  where  $k$  is a power of 2. But by [1, p. 147] we have that

$$[0^+(Q^-):1] = 3^{p(p-2)/4}(3^{p/2} - \varepsilon) \prod_{i=1}^{(p-2)/2} (3^{2i} - 1)$$

where  $\varepsilon$  is always either  $+1$  or  $-1$ . Replacing  $p$  by  $n - p$  gives us an expression for the order of  $0^+(Q^+)$ . By factoring out the powers of 3 we get the desired result.

**LEMMA 2.** *Let the dimension of  $M = n = 4r > 4$ . The highest*

power of the prime 3 dividing the order of the centralizer of the coset of a  $P'$ -involution  $T$  in  $PS^+(Q)$  is  $n(n - 4)/8$ .

*Proof.* Let  $F = K[i] = K[x]/(x^2 + 1)$  where  $i^2 = -1$ . Then  $F$  is the field with  $3^2 = 9$  elements. Let  $M_F = M \otimes_K F$ . This is a vector space of dimension  $n$  over  $F$ . The extension  $Q_F$  of  $Q$  to  $M_F$  is non-degenerate since  $Q$  is (see [5]). The  $P'$ -involution  $T$  can be extended to a transformation  $T_F$  on  $M_F$ , and  $T_F$  is a scalar multiple of an  $(m, m)$  involution  $T_F^*$ . In [5] it was shown that the centralizer of  $T$  in  $O^+(Q)$  is isomorphic to the orthogonal group  $O(Q_F^+)$  of the plus space of  $T_F^*$ . Hence by an argument similar to the one in the last lemma we get that  $[C_{PS^+}(\bar{T}):1] = k[O^+(Q_F^+):1]$  where  $k$  is a power of 2. Also

$$[O^+(Q_F^+):1] = 9^{m(m-2)/4}(9^{m/2} - \varepsilon) \prod_{i=1}^{(m-2)/2} (9^{2i} - 1),$$

where  $m = n/2$  and  $\varepsilon$  is either 1 or  $-1$ . The power of 3 dividing this expression is  $n(n - 4)/8$ .

*Proof of theorem.* Since the commutator subgroup  $(PS(Q))' = PO^+(Q)$  (see [3, p. 58]), and since

$$PS^+(Q) \cap PO(Q) = PO^+(Q),$$

no automorphism of  $PS(Q)$  or of  $PS^+(Q)$  can take an  $(n - 2, 2)$  coset into an  $(n - p, p)$  coset for  $p$  an odd integer.

Suppose  $\varphi$  is an automorphism of  $PS^+(Q)$ . Then  $C_{PS^+}(\varphi(\bar{T})) = \varphi(C_{PS^+}(\bar{T}))$  and the order of the centralizer of  $\bar{T}$  is the same as the order of the centralizer of  $\varphi(\bar{T})$ . In particular if  $\bar{T}$  is an  $(n - 2, 2)$  coset, the power of 3 dividing the order of  $C_{PS^+}(\bar{T})$  must be the same as that dividing  $C_{PS^+}(\varphi(\bar{T}))$ . Suppose  $\varphi(\bar{T})$  is an  $(n - p, p)$  coset. Then  $(n - 2)(n - 4)/4 = p(p - 2)/4 + (n - p)(n - p - 2)/4$  and  $p = 2$  or  $p = n - 2$ . The same holds if  $\varphi$  is an automorphism of  $PS(Q)$ .

Suppose  $\varphi(\bar{T})$  is the coset of a  $P'$ -involution. Then the dimension of  $M$  must be divisible by 4, since otherwise  $\varphi(\bar{T})$  would be an improper similitude. By Lemma 2 we have  $(n - 2)(n - 4)/4 = n(n - 4)/8$  and  $n = 4$  which is a contradiction.

An automorphism of  $PS(Q)$  can not take an  $(n - 2, 2)$  coset into the coset of a  $P'$ -involution since it must leave  $(PS(Q))' = PO^+(Q)$  invariant.

#### REFERENCES

1. E. Artin, *Geometric algebra*, Interscience Publishers, New York, 1957.
2. O. T. O'Meara, *Introduction to quadratic forms*, Academic Press, New York, 1963.
3. M. J. Wonenburger, *The Clifford algebra and the group of similitudes*, *Canad. J. Math.* **14** (1962), 45-59.

4. ———, *Study of certain similitudes*, *Canad. J. Math.* **14** (1962), 60-68.
5. ———, *Study of a semi-involutive similitude*, *Rev. Mat. Hisp.-Amer.* **20** (1960), 34-45.
6. ———, *The automorphisms of the group of similitudes and some related groups*, *Amer. J. Math.* **84** (1962), 600-614.

Received March 5, 1968. This paper is part of the author's Ph. D. dissertation written at the University of Virginia under the direction of Dr. E. C. Paige.

UNIVERSITY OF VIRGINIA  
AND  
UNIVERSITY OF GEORGIA

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. ROYDEN  
Stanford University  
Stanford, California

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. R. PHELPS  
University of Washington  
Seattle, Washington 98105

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL WEAPONS CENTER

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

# Pacific Journal of Mathematics

Vol. 28, No. 3

May, 1969

Jon F. Carlson, <i>Automorphisms of groups of similitudes over <math>F_3</math></i> .....	485
W. Wistar (William) Comfort, Neil Hindman and Stelios A. Negrepointis, <i><math>F'</math>-spaces and their product with <math>P</math>-spaces</i> .....	489
Archie Gail Gibson, <i>Triples of operator-valued functions related to the unit circle</i> .....	503
David Saul Gillman, <i>Free curves in <math>E^3</math></i> .....	533
E. A. Heard and James Howard Wells, <i>An interpolation problem for subalgebras of <math>H^\infty</math></i> .....	543
Albert Emerson Hurd, <i>A uniqueness theorem for weak solutions of symmetric quasilinear hyperbolic systems</i> .....	555
E. W. Johnson and J. P. Lediaev, <i>Representable distributive Noether lattices</i> .....	561
David G. Kendall, <i>Incidence matrices, interval graphs and seriation in archeology</i> .....	565
Robert Leroy Kruse, <i>On the join of subnormal elements in a lattice</i> .....	571
D. B. Lahiri, <i>Some restricted partition functions; Congruences modulo 3</i> ....	575
Norman D. Lane and Kamla Devi Singh, <i>Strong cyclic, parabolic and conical differentiability</i> .....	583
William Franklin Lucas, <i>Games with unique solutions that are nonconvex</i> .....	599
Eugene A. Maier, <i>Representation of real numbers by generalized geometric series</i> .....	603
Daniel Paul Maki, <i>A note on recursively defined orthogonal polynomials</i> ....	611
Mark Mandelker, <i><math>F'</math>-spaces and <math>z</math>-embedded subspaces</i> .....	615
James R. McLaughlin and Justin Jesse Price, <i>Comparison of Haar series with gaps with trigonometric series</i> .....	623
Ernest A. Michael and A. H. Stone, <i>Quotients of the space of irrationals</i> ....	629
William H. Mills and Neal Zierler, <i>On a conjecture of Golomb</i> .....	635
J. N. Pandey, <i>An extension of Haimo's form of Hankel convolutions</i> .....	641
Terence John Reed, <i>On the boundary correspondence of quasiconformal mappings of domains bounded by quasicircles</i> .....	653
Haskell Paul Rosenthal, <i>A characterization of the linear sets satisfying Herz's criterion</i> .....	663
George Thomas Sallee, <i>The maximal set of constant width in a lattice</i> .....	669
I. H. Sheth, <i>On normaloid operators</i> .....	675
James D. Stasheff, <i>Torsion in BBSO</i> .....	677
Billy Joe Thorne, <i><math>A - P</math> congruences on Baer semigroups</i> .....	681
Robert Breckenridge Warfield, Jr., <i>Purity and algebraic compactness for modules</i> .....	699
Joseph Zaks, <i>On minimal complexes</i> .....	721