

Pacific Journal of Mathematics

GAMES WITH UNIQUE SOLUTIONS THAT ARE NONCONVEX

WILLIAM FRANKLIN LUCAS

GAMES WITH UNIQUE SOLUTIONS THAT ARE NONCONVEX

W. F. LUCAS

In 1944 von Neumann and Morgenstern introduced a theory of solutions (stable sets) for n -person games in characteristic function form. This paper describes an eight-person game in their model which has a unique solution that is nonconvex. Former results in solution theory had not indicated that the set of all solutions for a game should be of this nature.

First, the essential definitions for an n -person game will be stated. Then, a particular eight-person game is described. Finally, there is a brief discussion on how to construct additional games with unique and nonconvex solutions.

The author [2] has subsequently used some variations of the techniques described in this paper to find a ten-person game which has no solution; thus providing a counterexample to the conjecture that every n -person game has a solution in the sense of von Neumann and Morgenstern.

2. Definitions. An n -person game is a pair (N, v) where $N = \{1, 2, \dots, n\}$ and v is a real valued characteristic function on 2^N , that is, v assigns the real number $v(S)$ to each subset S of N and $v(\varnothing) = 0$. The set of all *imputations* is

$$A = \left\{ x: \sum_{i \in N} x_i = v(N) \text{ and } x_i \geq v(\{i\}) \text{ for all } i \in N \right\}$$

where $x = (x_1, x_2, \dots, x_n)$ is a vector with real components. If x and y are in A and S is a nonempty subset of N , then $x \text{ dom}_S y$ means $\sum_{i \in S} x_i \leq v(S)$ and $x_i > y_i$ for all $i \in S$. For $B \subset A$ let $\text{Dom}_S B = \{y \in A: \text{there exists } x \in B \text{ such that } x \text{ dom}_S y\}$ and let $\text{Dom } B = \bigcup_{S \subset N} \text{Dom}_S B$. A subset K of A is a *solution* if $K \cap \text{Dom } K = \varnothing$ and $K \cup \text{Dom } K = A$. The *core* of a game is

$$C = \left\{ x \in A: \sum_{i \in S} x_i \geq v(S) \text{ for all } S \subset N \right\}.$$

The core consists of those imputations which are maximal with respect to all of the relations dom_S , and hence it is contained in every solution.

3. Example. Consider the game (N, v) where $N = \{1, 2, 3, 4, 5, 6, 7, 8\}$ and where v is given by: $v(N) = 4$, $v(\{1, 4, 6, 7\}) = 2$, $v(\{1, 2\}) =$

$v(\{3, 4\}) = v(\{5, 6\}) = v(\{7, 8\}) = 1$, and $v(S) = 0$ for all other $S \subset N$. For this game

$$A = \left\{ x: \sum_{i \in N} x_i = 4 \text{ and } x_i \geq 0 \text{ for all } i \in N \right\}$$

and

$$C = \{x \in A: x_1 + x_2 = x_3 + x_4 = x_5 + x_6 = x_7 + x_8 = 1 \text{ and } x_1 + x_4 + x_6 + x_7 \geq 2\}.$$




FIG. 1. Traces in H of L , C and $K-C$

Also define the four-dimensional hypercube

$$H = \{x \in A: x_1 + x_2 = x_3 + x_4 = x_5 + x_6 = x_7 + x_8 = 1\} .$$

Three traces of H as well as its 16 vertices are pictured in Fig. 1.

The unique solution for this game is

$$K = C \cup F_1 \cup F_4 \cup F_6 \cup F_7,$$

where the cube F_i is the face of H given by

$$F_i = H \cap \{x: x_i = 1\} \quad i = 1, 4, 6, 7 .$$

Each $F_i - C$ is a tetrahedron with one face meeting C . In the three traces of H illustrated in Fig. 1, the traces of C are shown in heavy solid lines and the traces of the $F_i - C$ are shown in heavy broken lines.

The proof that K is the unique solution follows readily from two observations. First, K is just those imputations in H which are maximal in H with respect to the relation $\text{dom}_{\{1,4,6,7\}}$. Second, the closed line segment L joining the imputations $(0, 1, 0, 1, 0, 1, 0, 1)$ and $(1, 0, 1, 0, 1, 0, 1, 0)$ has the properties $L \subset C$ and $\bigcup_S \text{Dom}_S L = A - H$ when $S = \{1, 2\}, \{3, 4\}, \{5, 6\},$ and $\{7, 8\}$.

To see that K is nonconvex, note the lower trace

$$F_8 = H \cap \{x: x_8 = 1\}$$

in Fig. 1. The heavy lines (solid and broken) in this trace show $K \cap F_8$, which is clearly not convex. For example, the imputation

$$\begin{aligned} \frac{1}{3}(1, 2, 2, 1, 2, 1, 0, 3) &= \frac{1}{3}(0, 1, 1, 0, 0, 1, 0, 1) \\ &+ \frac{1}{3}(0, 1, 0, 1, 1, 0, 0, 1) + \frac{1}{3}(1, 0, 1, 0, 1, 0, 0, 1) \end{aligned}$$

is a linear combination of points in K , but it is not itself in K .

4. Remarks. The original von Neumann-Morgenstern theory [3] assumed that the characteristic function of a game is superadditive, that is, $v(S_1 \cup S_2) \geq v(S_1) + v(S_2)$ whenever S_1 and $S_2 \subset N$ and $S_1 \cap S_2 = \varphi$. Using the method of Gillies [1, p. 68] this example can be made into a game with a superadditive characteristic function without changing A, C , or the unique solution K .

The essential idea in the example above is that $\bigcup_S \text{Dom}_S L = A - H$ where $S = \{1, 2\}, \{3, 4\}, \{5, 6\},$ and $\{7, 8\}$. One can generalize this relation in various ways to obtain many games in other dimensions which have a similar property. He can then introduce into these games additional $S \subset N$ with $v(S) > 0$, but in such a way as to maintain the corresponding L as a subset of the core. As a result he will

obtain large classes of interesting solutions, many of which are unique and nonconvex.

REFERENCES

1. D. B. Gillies, *Solutions to general non-zero-sum games*, Annals of Mathematics Studies, No. 40, A. W. Tucker and R. D. Luce (editors), Princeton University Press, Princeton, 1959.
2. W. F. Lucas, *A Game with no solution*, RAND Memorandum RM-5518-PR, The RAND Corporation, Santa Monica, November 1967.
3. J. von Neumann and O. Morgenstern, *Theory of games and economic behavior*, Princeton University Press, Princeton, (1944).

Received June 26, 1967.

THE RAND CORPORATION
SANTA MONICA, CALIFORNIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R. PHELPS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

| | |
|------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| Jon F. Carlson, <i>Automorphisms of groups of similitudes over F_3</i> | 485 |
| W. Wistar (William) Comfort, Neil Hindman and Stelios A. Negrepointis, <i>F'-spaces and their product with P-spaces</i> | 489 |
| Archie Gail Gibson, <i>Triples of operator-valued functions related to the unit circle</i> | 503 |
| David Saul Gillman, <i>Free curves in E^3</i> | 533 |
| E. A. Heard and James Howard Wells, <i>An interpolation problem for subalgebras of H^∞</i> | 543 |
| Albert Emerson Hurd, <i>A uniqueness theorem for weak solutions of symmetric quasilinear hyperbolic systems</i> | 555 |
| E. W. Johnson and J. P. Lediaev, <i>Representable distributive Noether lattices</i> | 561 |
| David G. Kendall, <i>Incidence matrices, interval graphs and seriation in archeology</i> | 565 |
| Robert Leroy Kruse, <i>On the join of subnormal elements in a lattice</i> | 571 |
| D. B. Lahiri, <i>Some restricted partition functions; Congruences modulo 3</i> | 575 |
| Norman D. Lane and Kamla Devi Singh, <i>Strong cyclic, parabolic and conical differentiability</i> | 583 |
| William Franklin Lucas, <i>Games with unique solutions that are nonconvex</i> | 599 |
| Eugene A. Maier, <i>Representation of real numbers by generalized geometric series</i> | 603 |
| Daniel Paul Maki, <i>A note on recursively defined orthogonal polynomials</i> | 611 |
| Mark Mandelker, <i>F'-spaces and z-embedded subspaces</i> | 615 |
| James R. McLaughlin and Justin Jesse Price, <i>Comparison of Haar series with gaps with trigonometric series</i> | 623 |
| Ernest A. Michael and A. H. Stone, <i>Quotients of the space of irrationals</i> | 629 |
| William H. Mills and Neal Zierler, <i>On a conjecture of Golomb</i> | 635 |
| J. N. Pandey, <i>An extension of Haimo's form of Hankel convolutions</i> | 641 |
| Terence John Reed, <i>On the boundary correspondence of quasiconformal mappings of domains bounded by quasicircles</i> | 653 |
| Haskell Paul Rosenthal, <i>A characterization of the linear sets satisfying Herz's criterion</i> | 663 |
| George Thomas Sallee, <i>The maximal set of constant width in a lattice</i> | 669 |
| I. H. Sheth, <i>On normaloid operators</i> | 675 |
| James D. Stasheff, <i>Torsion in BBSO</i> | 677 |
| Billy Joe Thorne, <i>$A - P$ congruences on Baer semigroups</i> | 681 |
| Robert Breckenridge Warfield, Jr., <i>Purity and algebraic compactness for modules</i> | 699 |
| Joseph Zaks, <i>On minimal complexes</i> | 721 |