

Pacific Journal of Mathematics

TORSION IN BBSO

JAMES D. STASHEFF

TORSION IN BBSO

JAMES D. STASHEFF

The cohomology of BBSO, the classifying space for the stable Grassmanian BSO, is shown to have torsion of order precisely 2^r for each natural number r . Moreover, the elements of order 2^r appear in a pattern of striking simplicity.

Many of the stable Lie groups and homogeneous spaces have torsion at most of order 2 [1, 3, 5]. There is one such space, however, with interesting torsion of higher order. This is $BBSO = SU/\text{Spin}$ which is of interest in connection with Bott periodicity and in connection with the J-homomorphism [4, 7]. By the notation SU/Spin we mean that $BBSO$ can be regarded as the fibre of $B \text{ Spin} \rightarrow BSU$ or that, up to homotopy, there is a fibration

$$SU \rightarrow BBSO \rightarrow B \text{ Spin}$$

induced from the universal SU bundle by $B \text{ Spin} \rightarrow BSU$. The mod 2 cohomology $H^*(BBSO; Z_2)$ has been computed by Clough [4]. The purpose of this paper is to compute enough of $H^*(BBSO; Z)$ to obtain the mod 2 Bockstein spectral sequence [2] of $BBSO$.

Given a ring R , we shall denote by $R[x_i | i \in I]$ the polynomial ring on generators x_i indexed by elements of a set I . The set I will often be described by an equation or inequality in which case i is to be understood to be a natural number. Similarly $E(x_i | i \in I)$ will denote the exterior algebra on generators x_i . In this case, we will need only $R = Z_2$.

Let us recall the results on $B \text{ Spin}$ as given by Thomas [6] and on $BBSO$ as given by Clough [4].

$$H^*(B \text{ Spin}; Z_2) \approx Z_2[w_i | i \neq 2^j + 1]$$

where w_i is (the image of) the Stiefel-Whitney class w_i .

$$H^*(B \text{ Spin}; Z) \approx Z[Q_i | i > 0] \oplus \hat{T}$$

where $2\hat{T} = 0$ and $Q_i \in H^{4i}$.

$$H^*(BBSO; Z_2) \approx E(e_i | i \geq 3)$$

where $e_i \in H^i$ and is the image of w_i if $i \neq 2^j + 1$ while e_{2^j+1} maps to an indecomposable element in $H^*(SU; Z_2)$.

Now let ${}_\beta E_r$ denote the mod 2 Bockstein spectral sequence of $BBSO$ [2]. In particular, ${}_\beta E_2 = \text{Ker } Sq^1 / \text{Im } Sq^1$. Now $Sq^1 w_{2i} = w_{2i+1}$ in BSO and $Sq^1 w_{2i+1} = 0$ while $Sq^1 e_{2^j} = 0$ in $B \text{ Spin}$. We will see that

e_{2^j+1} can be chosen to have $Sq^1 e_{2^j+1} = 0$ except for $Sq^1 e_3 = e_4$. Thus

$$\beta E_2 = E(e_3 e_4, e_{2^2+i}, v_{4i+1} \mid i > 0)$$

where $v_{4i+1} = e_{2i} e_{2i+1}$ except $v_{2^j+1} = e_{2^j+1}; j > 1$.

THEOREM 1.

$$\beta E_r \approx E(e_3 e_4 \cdots e_{2^r}, e_{2^r+i}, v_{4i+1} \mid i > 0)$$

and $d_r(e_3 \cdots e_{2^r}) = e_{2^r+1}$ modulo decomposable elements.

To prove Theorem 1, we will exhibit torsion of order 2^r for all r .

THEOREM 2. *In $H^*(BBSO; Z)$, we have*

$$2^r Q_{2^r} \neq 0 \quad \text{and} \quad 2^{r+1} Q_{2^r} = 0 .$$

$H^*(BBSO; Z_2)$. We recall some of Clough's observations on $H^*(BBSO; Z_2)$. We know $H^*(SU; Z_2) = E(y_i \mid i > 1)$ where $y_i \in H^{2i+1}$ transgresses universally to the mod 2 reduction of the Chern class c_i and hence to the image of w_i^2 in $B \text{ Spin}$. Thus $w_i^2 = 0$ in $BBSO$ for $i \neq 2^j + 1$ and y_{2^j} is the restriction of a class e_{2^j+1+1} . In particular since $Sq^{2^j}(w_{2^j-1+1})^2 = (w_{2^j+1})^2$ we can take e_{2^j+1} to be $Sq^{2^j-1} Sq^{2^j-2} \cdots Sq^4 Sq^2 e_3$. The class e_3 is uniquely determined ($H^3(BBSO; Z_2) \approx Z_2$) and this definition of e_{2^j+1} implies $Sq^1 e_{2^j+1+1} = (e_{2^i+1})^2 = 0$ if $e_3^2 = 0$. The only alternative to $e_3^2 = 0$ is $e_3^2 = e_6$; there is no other class in this dimension. Since $Sq^1 w_6 = w_7$ in $B \text{ Spin}$ and w_6, w_7 map to e_6, e_7 , we have $Sq^1 e_6 = e_7$ but $Sq^1(e_3)^2 = 0$; therefore e_3^2 must be zero.

$H^*(BBSO, Z)$. Consider $BBSO$ as the fibre of $B \text{ Spin} \rightarrow BSU$. The latter map factors: $B \text{ Spin} \xrightarrow{\pi} BSO \rightarrow BSU$. Recall that

$$H^*(BSU; Z) = Z[c_i \mid i > 1] \quad \text{and} \quad H^*(BSO; Z) = Z[P_i] \oplus T$$

where T is the torsion ideal, $2T = 0$, c_{2i+1} maps into T and c_{2i} maps to P_i . To determine $\text{Im}(H^*(B \text{ Spin}))$ in $H^*(BBSO)$, we need to know $\pi^*[P_i]$ in $H^*(B \text{ Spin})$.

THEOREM 3 (Thomas [6]). *If i is not a power of 2, $\pi^* P_i = Q_i$. If $j = 2^r, r > 0, \pi^* P_{2^j} = 2Q_{2^j} + Q_j^2 - \pi^* \Phi_{2^j}$. $\pi^* P_1 = 2Q_1$.*

LEMMA. $\pi^* \Phi_{2^j}$ maps into $\text{Im } T \subset H^*(BBSO)$.

Proof. $H^*(BSO; Z)$ maps onto $\text{Im } T$ in $H^*(BBSO)$ since $H^*(BSU)$ maps onto the $Z[P_i]$ part.

Since $\pi^* P_j$ goes to zero in $BBSO$, we have in $H^*(BBSO; Z)$

$$2Q_{2^j} = -Q_j^2 + t \quad \text{where } 2t = 0 \quad \text{and } j = 2^r .$$

$$2Q_1 = 0 .$$

By iteration we find

$$2^{r+1}Q_{2^r} = \pm 2Q_{2^r}Q_{2^{r-1}} \cdots Q_2(Q_1)^2 = 0 .$$

To determine the order of Q_{2^i} in $BBSO$, consider $\Gamma(u \mid 2u = 0)$, a divided polynomial algebra on a single generator u of dimension 4 and order 2; i.e., additively Γ has generators $\gamma_i(u)$ in dimension $4i$ and the multiplication table is $\gamma_i(u)\gamma_j(u) = (i, j)\gamma_{i+j}(u)$ where (i, j) is the binomial coefficient $\{(i + j)!/i!j!\}$.

In particular $i!\gamma_i(u) = u^i$.

We construct a map f from $\text{Im}(H^*(B\text{Spin}; Z) \rightarrow H^*(BBSO; Z))$ to Γ by mapping \hat{T} to zero, Q_i to zero for $i \neq 2^j$ and Q_{2^j} to $-\gamma_2(f(Q_{2^{j-1}}))$ with $f(Q_1) = u$. Since $2Q_{2^j} = -Q_{2^{j-1}}^2 + \pi^*\Phi_{2^j}$, and Φ_{2^j} goes into $\text{Im } \hat{T}$ in $BBSO$, the map f is well defined. Since for any x , the order of $\gamma_2(x)$ is twice the order of x , we have

$$\text{ord } f(Q_{2^j}) = 2 \text{ord } f(Q_{2^{j-1}}) = 2^j \text{ord } f(Q_1) = 2^{j+1} .$$

Thus the order of Q_{2^j} is at least 2^{j+1} and that $2^{j+1}Q_{2^j}$ is in fact zero we have already seen.

Thus we have 2^r torsion for each r . From the exact cohomology sequence derived from $0 \rightarrow Z \xrightarrow{2^r} Z \rightarrow Z_{2^r} \rightarrow 0$, we see that $Q_{2^{r-1}} = \beta_{2^r}^\infty x_r$ for some class $x_r \in H^*(BBSO; Z_{2^r})$, where $\beta_{2^r}^\infty$ is the connecting homomorphism $H^*(; Z_{2^r}) \rightarrow H^{*+1}(; Z)$.

LEMMA. $(\beta_{2^r}^\infty x_r)_2 = d_r(x_r)_2$ where $()_2$ means reduction mod 2.

Proof. Recall how d_r is defined: $d_r(x) = (\beta_2^\infty(x)/2^{r-1})_2$. From the commutativity of the diagram

$$\begin{array}{ccccc} Z & \xrightarrow{2^r} & Z & \longrightarrow & Z_{2^r} \\ & & \downarrow 2^{r-1} & \parallel & \downarrow \\ Z & \xrightarrow{2} & Z & \longrightarrow & Z_2 \end{array}$$

it follows that $\beta_2^\infty = 2^{r-1}\beta_{2^r}^\infty$. In particular, $d_r(x_r)_2 = (Q_{2^{r-1}})_2$. According to Thomas, $(Q_{2^{r-1}})_2 = \pi^*(w_{2^r+1} + \psi_{2^r+1})$ where ψ_{2^r+1} is decomposable. In particular, $(Q_1)_2 = W_4$.

We prove Theorem 2 by induction. Since

$$Sq^1 w_{2i} = w_{2i+1} \quad \text{and} \quad Sq^1 w_{2i+1} = 0 ,$$

we know $Sq^1 e_{2i} = e_{2i+1}$ and $Sq^1 e_{2i+1} = 0$ unless $i = 2^j$. Since we have chosen $e_{2^{j+1}} = Sq^{2^j-1} \cdots Sq^2 e_3$, we have $Sq^1 e_{2^{j+1}} = (e_{2^j-1+1})^2 = 0$ for

$j \geq 2$. For $j = 1$, we have $Sq^1 e_3 = e_4$ because $e_4 = (Q_1)_2$ which is in the image of Sq^1 since $2Q_1 = 0$.

Thus

$$\begin{aligned} \beta E_2 &= \text{Ker } Sq^1 / \text{Im } Sq^1 \\ &= E(e_3 e_4) \otimes E(e_{2i} e_{2i+1} \mid 2 < i \neq 2^j) \otimes E(e_{2^j+1}, e_{2^j+1} \mid j \geq 2). \end{aligned}$$

Since $d_2(x_2)_2 = (Q_2)_2 = e_3$, we must have $x_2 = e_3 e_4$.

In general $d_r(x_r)_2 = (Q_{2^{r-1}})_2 = e_{2^{r+1}}$ modulo decomposables. Now consider $H^*(BBSO; Q)$. Since $H^*(BSO; Q) = Q[P_i]$ with the usual diagonal $m^*(P_i) = \sum_{j+k=i} P_j \otimes P_k$, we have $H^*(BBSO; Q) = E(R_i)$ where $\dim R_i \in H^{4j+1}$. Thus $\beta E_\infty = E(S_{4i+1})$ and the only possibility is

$$\begin{aligned} S_{4i+1} &= e_{2i} e_{2i+1} \quad i \neq 2^j, \\ S_{2^i+1} &= e_{2^i+1} \end{aligned}$$

modulo terms decomposable in terms of the S_{4i+1} . This leaves $e_3 e_4 \cdots e_{2^r}$ as the only possibility for x_r , i.e., $d_r(e_3 e_4 \cdots e_{2^r}) = e_{2^{r+1}}$ mod decomposables as claimed.

BIBLIOGRAPHY

1. A. Borel, *La cohomologie mod 2 des certains espaces homogènes*, Comm. Math. Helv. **27** (1953), 165-197.
2. W. Browder, *Torsion in H-spaces*, Ann. of Math. **74** (1961), 24-51.
3. H. Cartan, *Séminaire—1959/60*, Exposés 3, 17, Paris, 1961.
4. R. R. Clough, *Calculation of $H^*(B_{\text{Im}(j)}; Z_2)$* (to appear).
5. E. Dyer and R. K. Lashof, *A topological proof of the Bott periodicity theorem*, Ann. Mat. Pura Appl. (4) **54** (1961), 231-254.
6. E. Thomas, *On the cohomology groups of the classifying space for the stable spinor group*, Bol. Soc. Mat. Mex. (1962), 57-69.
7. G. W. Whitehead, *On the homotopy groups of spheres and rotation groups*, Ann. of Math. **43** (1942), 634-640.

Received May 3, 1968. Research supported in part by NSF Grant GP-6101. The author is an Alfred P. Sloan Fellow.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R. PHELPS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics

Vol. 28, No. 3

May, 1969

Jon F. Carlson, <i>Automorphisms of groups of similitudes over F_3</i>	485
W. Wistar (William) Comfort, Neil Hindman and Stelios A. Negrepointis, <i>F'-spaces and their product with P-spaces</i>	489
Archie Gail Gibson, <i>Triples of operator-valued functions related to the unit circle</i>	503
David Saul Gillman, <i>Free curves in E^3</i>	533
E. A. Heard and James Howard Wells, <i>An interpolation problem for subalgebras of H^∞</i>	543
Albert Emerson Hurd, <i>A uniqueness theorem for weak solutions of symmetric quasilinear hyperbolic systems</i>	555
E. W. Johnson and J. P. Lediaev, <i>Representable distributive Noether lattices</i>	561
David G. Kendall, <i>Incidence matrices, interval graphs and seriation in archeology</i>	565
Robert Leroy Kruse, <i>On the join of subnormal elements in a lattice</i>	571
D. B. Lahiri, <i>Some restricted partition functions; Congruences modulo 3</i>	575
Norman D. Lane and Kamla Devi Singh, <i>Strong cyclic, parabolic and conical differentiability</i>	583
William Franklin Lucas, <i>Games with unique solutions that are nonconvex</i>	599
Eugene A. Maier, <i>Representation of real numbers by generalized geometric series</i>	603
Daniel Paul Maki, <i>A note on recursively defined orthogonal polynomials</i>	611
Mark Mandelker, <i>F'-spaces and z-embedded subspaces</i>	615
James R. McLaughlin and Justin Jesse Price, <i>Comparison of Haar series with gaps with trigonometric series</i>	623
Ernest A. Michael and A. H. Stone, <i>Quotients of the space of irrationals</i>	629
William H. Mills and Neal Zierler, <i>On a conjecture of Golomb</i>	635
J. N. Pandey, <i>An extension of Haimo's form of Hankel convolutions</i>	641
Terence John Reed, <i>On the boundary correspondence of quasiconformal mappings of domains bounded by quasicircles</i>	653
Haskell Paul Rosenthal, <i>A characterization of the linear sets satisfying Herz's criterion</i>	663
George Thomas Sallee, <i>The maximal set of constant width in a lattice</i>	669
I. H. Sheth, <i>On normaloid operators</i>	675
James D. Stasheff, <i>Torsion in BBSO</i>	677
Billy Joe Thorne, <i>$A - P$ congruences on Baer semigroups</i>	681
Robert Breckenridge Warfield, Jr., <i>Purity and algebraic compactness for modules</i>	699
Joseph Zaks, <i>On minimal complexes</i>	721