

Pacific Journal of Mathematics

THE LATTICE OF PRETOPOLOGIES ON AN ARBITRARY SET
 S

ALLAN MATLOCK WEBER CARSTENS

THE LATTICE OF PRETOPOLOGIES ON AN ARBITRARY SET S

ALLAN M. CARSTENS

The structure of the lattice of pretopologies on the set S , unlike that of the lattice of topologies on S (a proper sublattice of the former), has not been closely examined. We establish that pretopologies may be identified with products of certain filters in a natural way. From this identification, we are able to determine much of the structure of this lattice.

We show that $(p(S), \leq)$, the lattice of pretopologies (pretopologies in the sense of Kent [2; p. 126]) on the set S , is order isomorphic to a sublattice of filters on S^S (using Bourbaki's [1; p. 61-63] approach to filters). From this, we deduce that $(p(S), \leq)$ is complete, atomic, coatomic, modular, distributive, and compactly generated; S being finite is both necessary and sufficient for the lattice to be co-compactly generated and complemented (in which case it has a unique complement). It is infinitely distributive only in the trivial case of S being finite. (The lattice terminology is that of Szász [3] with the exception of coatomic which we use rather than dually atomic and co-compactly generated which is used for the notion dual to that of compactly generated.)

1. The isomorphism φ . A pretopology p on a set S is completely determined by a specification of the neighborhood filter $\eta_p(x)$ of each x in S . These neighborhoods necessarily satisfy $\eta_p(x) \leq \overline{x}$, where \overline{x} is the principal filter generated by $\{x\}$. For each $x \in S$, let $\underline{F}(x) = \{\mathfrak{F}: \mathfrak{F} \geq \overline{x}, \mathfrak{F} \text{ a filter on } S\}$, and let $\underline{F} = \prod_{x \in S} \underline{F}(x)$ (Bourbaki [1; p. 69-70]); both ordered by $\mathfrak{F} \leq \mathfrak{G}$ if and only if $\mathfrak{F} \subset \mathfrak{G}$. Then \underline{F} is a subset of the set of filters on S^S . Indeed, it is easily seen that \underline{F} , with this ordering, is a sublattice of the lattice of filters on S^S . For given $\mathfrak{F}, \mathfrak{G} \in \underline{F}$, we have $\mathfrak{F} \wedge \mathfrak{G} = \{F \cup G: F \in \mathfrak{F}, G \in \mathfrak{G}\}$ and $\mathfrak{F} \vee \mathfrak{G} = \{F \cap G: F \in \mathfrak{F}, G \in \mathfrak{G}\}$ ($F \cap G \neq \phi$ since $\prod_{x \in S} \{x\} \in F, G$).

Given a pretopology p , we define φ by $\varphi(p) = \prod_{x \in S} \eta_p(x)$. Then φ is easily seen to be a one-to-one mapping from the pretopologies on S onto \underline{F} . Furthermore, if p, q are pretopologies on S , the following will be equivalent:

- (1) $p \leq q$;
- (2) $\eta_p(x) \leq \eta_q(x)$ for all x in S ; and
- (3) $\prod_{x \in S} \eta_p(x) \leq \prod_{x \in S} \eta_q(x)$.

Thus we have

THEOREM 1. φ is an order isomorphism from the lattice of pretopologies on S onto the sublattice (\underline{F}, \leq) of filters on S^s .

2. The structure of $(\underline{F}(x), \leq)$. We shall deduce the structure of (\underline{F}, \leq) from an examination of the structures of the lattices $(\underline{F}(x), \leq)$, for each x .

It follows readily from the definition of \wedge and \vee in $(\underline{F}(x), \leq)$ that this lattice is complete, modular, and distributive. The remaining propositions of this section further describe its structure.

PROPOSITION 1. $(\underline{F}(x), \leq)$ is atomic. Its atoms are precisely those elements of the form $\overline{S \setminus \{a\}}$ for $a \neq x$. (\overline{A} denotes the filter of all super-sets of A in S).

Proof. Given $\mathfrak{F} \neq 0 \equiv \overline{S}$ in $\underline{F}(x)$, select $A \in \mathfrak{F}$, $A \neq S$. Then there exists an $a \neq x$ in $S \setminus A$, and $\overline{S \setminus \{a\}} \leq \mathfrak{F}$.

To show that $\overline{S \setminus \{a\}}$ is an atom of $(\underline{F}(x), \leq)$ for $a \neq x$, let $\mathfrak{F} < \overline{S \setminus \{a\}}$. Then $S \setminus \{a\} \subset F$ for all $F \in \mathfrak{F}$, and $F \subset S \setminus \{a\}$ for no $F \in \mathfrak{F}$. Thus $\mathfrak{F} = \overline{S} = 0$.

PROPOSITION 2. $(\underline{F}(x), \leq)$ is coatomic. Its coatoms are precisely those $\mathfrak{F} = \overline{x} \wedge \mathfrak{U}$ where $\mathfrak{U} \neq \overline{x}$ is an ultrafilter.

Proof. Let $\mathfrak{F} \in \underline{F}(x)$ be distinct from $1 \equiv \overline{x}$. Then since \mathfrak{F} is not an ultrafilter, there must be at least two ultrafilters above \mathfrak{F} . One of these, say \mathfrak{U} , must be distinct from \overline{x} . Then $\mathfrak{F} \leq \overline{x} \wedge \mathfrak{U}$.

To show that $\overline{x} \wedge \mathfrak{U}$ is a coatom of $(\underline{F}(x), \leq)$, assume there is an $\mathfrak{G} \in \underline{F}(x)$ with $\overline{x} \wedge \mathfrak{U} < \mathfrak{G} < \overline{x}$. Since $\mathfrak{F} < \overline{x}$, $\{F \setminus \{x\} : F \in \mathfrak{F}\}$ is a base for some filter \mathfrak{G} . Clearly $\overline{x} \wedge \mathfrak{G} = \mathfrak{F}$. Now, for each $U \in \mathfrak{U}$, there exists $F \in \mathfrak{F}$ such that $F \subset U \subset \{x\}$, since $\overline{x} \wedge \mathfrak{U} < \mathfrak{F}$. Thus $F \setminus \{x\} \subset U$. Hence $\mathfrak{G} \geq \mathfrak{U}$. But \mathfrak{U} is an ultrafilter. Consequently $\mathfrak{G} = \mathfrak{U}$. Thus, we must conclude that $\overline{x} \wedge \mathfrak{U} = \overline{x} \wedge \mathfrak{G} = \mathfrak{F}$, a contradiction.

PROPOSITION 3. $\mathfrak{F} \in \underline{F}(x)$ is compact if and only if $\mathfrak{F} = \overline{A}$ for some $A \subset S$ with $x \in A$. Consequently $(\underline{F}(x), \leq)$ is compactly generated.

Proof. Let $\mathfrak{F} \in \underline{F}(x)$ be compact. Observe that $\mathfrak{F} = \bigvee \{\overline{F} : F \in \mathfrak{F}\}$. Thus $\mathfrak{F} \leq \bigvee_{i=1}^n \overline{F}_i$ for some choice of n and $F_i \in \mathfrak{F}$ ($i = 1, \dots, n$). But since filters include finite intersections of their members, $\mathfrak{F} \geq \overline{\bigcap_{i=1}^n F_i} = \bigvee_{i=1}^n \overline{F}_i$. Thus $\mathfrak{F} \equiv \overline{\bigcap_{i=1}^n F_i}$.

Conversely, let $\mathfrak{F} = \overline{A}$ and let $\mathfrak{F} \leq \bigvee_{\gamma \in \Gamma} \mathfrak{F}_\gamma$. Then since $A \in \mathfrak{F}$, there exists $\Gamma_0 \subset \Gamma$ (Γ_0 finite), and $F_\gamma \in \mathfrak{F}_\gamma$ such that $\bigcap_{\gamma \in \Gamma_0} F_\gamma \subset A$.

Thus $\mathfrak{F} \leq \bigvee_{r \in r_0} F_r$.

For any $\mathfrak{G} \in \underline{F}(x)$, we have

$$\begin{aligned} \mathbf{V}\{\mathfrak{F}: \mathfrak{F} \leq \mathfrak{G}, \mathfrak{F} \text{ compact}\} &\leq \mathfrak{G} = \mathbf{V}\{\overline{G}: G \in \mathfrak{G}\} \\ &\leq \mathbf{V}\{\mathfrak{F}: \mathfrak{F} \leq \mathfrak{G}, \mathfrak{F} \text{ compact}\} \end{aligned}$$

thus $\mathfrak{G} = \mathbf{V}\{\mathfrak{F}: \mathfrak{F} \leq \mathfrak{G}, \mathfrak{F} \text{ compact}\}$ and $(\underline{F}(x), \leq)$ is compactly generated.

PROPOSITION 4. $\mathfrak{F} \in \underline{F}(x)$ is co-compact if and only if $\mathfrak{F} = \overline{A}$ where A is some finite subset of S containing x . Consequently $(\underline{F}(x), \leq)$ is co-compactly generated if and only if S is finite.

Proof. Let $\mathfrak{F} \in \underline{F}(x)$ be co-compact and let $T = S \setminus \{x\}$. Observe that $\mathfrak{F} \geq \overline{S} \equiv \bigwedge_{a \in T} \overline{\{x, a\}}$. Consequently for some n and $a_i \in T$ ($i = 1, \dots, n$), $\mathfrak{F} \geq \bigwedge_{i=1}^n \overline{\{x, a_i\}} = \overline{\bigcup_{i=1}^n \{x, a_i\}}$. Thus $\overline{\bigcup_{i=1}^n \{x, a_i\}} \in \mathfrak{F}$. But any filter containing a finite set B can be expressed as \overline{A} for some $A \subseteq B$. Thus $\mathfrak{F} = \overline{A}$ for some $A \subseteq \bigcup_{i=1}^n \{x, a_i\}$.

Conversely, let $\mathfrak{F} = \overline{A}$ where $A = \{x, a_1, a_2, \dots, a_n\}$, and suppose that $\mathfrak{F} \geq \bigwedge_{r \in r} \mathfrak{F}_r$. Then we may select $F_r \in \mathfrak{F}_r$ such that $\bigcup_{r \in r} F_r \supseteq A$. Select γ_i so that $a_i \in F_{\gamma_i}$. Then $\mathfrak{F} \geq \bigwedge_{i=1}^n \mathfrak{F}_{\gamma_i}$.

If S is finite, each $\mathfrak{F} \in \underline{F}(x)$ is of the form \overline{A} with A finite, so $(\underline{F}(x), \leq)$ will consist only of co-compact elements and hence be co-compactly generated. Observe however, that $\bigwedge_{r \in r} \overline{A}_r = \overline{\bigcup_{r \in r} A_r}$ for arbitrary filters. Thus, in particular, the only elements of $(\underline{F}(x), \leq)$ which will be co-compactly generated are the principal filters. Consequently $(\underline{F}(x), \leq)$ is not co-compactly generated when S is infinite.

PROPOSITION 5. $\mathfrak{F} \in \underline{F}(x)$ has a complement \mathfrak{G} if and only if $\mathfrak{F} = \overline{A}$. In this case \mathfrak{G} is unique and $\mathfrak{G} = \overline{(S \setminus A) \cup \{x\}}$. Consequently $(\underline{F}(x), \leq)$ is complemented if and only if S is finite.

Proof. Let $\mathfrak{F} = \overline{A}$. If $\mathfrak{G} = \overline{(S \setminus A) \cup \{x\}}$, then $\mathfrak{F} \wedge \mathfrak{G} = \overline{S} \equiv 0$ and $\mathfrak{F} \vee \mathfrak{G} = \overline{x} \equiv 1$. Thus \mathfrak{G} is a complement of \mathfrak{F} . Let \mathfrak{G}' be any complement of \mathfrak{F} . Then since $\mathfrak{F} \wedge \mathfrak{G}' = \overline{S}$, $(S \setminus A) \cup \{x\}$ must be in \mathfrak{G}' . But $\mathfrak{F} \vee \mathfrak{G}' = \overline{x}$, so no proper subset of $(S \setminus A) \cup \{x\}$ may be in \mathfrak{G}' . Consequently $\mathfrak{G}' = \overline{(S \setminus A) \cup \{x\}} = \mathfrak{G}$.

Suppose on the other hand, that \mathfrak{F} is not principal. Let $A = \bigcap \mathfrak{F}$. Then $A \neq \phi$ since $x \in A$. Suppose that \mathfrak{G} is a complement of \mathfrak{F} . Then for each $F \in \mathfrak{F}$, $G \in \mathfrak{G}$, we have $F \cup G = S$, since $\mathfrak{F} \wedge \mathfrak{G} = \overline{S}$. Thus $B = (S \setminus A) \cup \{x\}$ must be a subset of every G in \mathfrak{G} . Observe that any F in \mathfrak{F} will contain A as a proper subset since \mathfrak{F} is nonprincipal. Thus any $F \in \mathfrak{F}$ will include points of B distinct from x . Hence for each $G \in \mathfrak{G}$, $F \vee G$ will contain at least two points. But this violates

the requirement that $\mathfrak{F} \vee \mathfrak{G} = \overline{x}$. Therefore \mathfrak{F} can not have a complement.

We conclude this section with a discussion of infinite distributivity. Let $\mathfrak{F}, \mathfrak{F}_\gamma \in \underline{F}(x)$ ($\gamma \in \Gamma$) be arbitrary. Then, since filter joins are given by finite intersections, we have $\mathfrak{F} \wedge (\bigvee_{\gamma \in \Gamma} \mathfrak{F}_\gamma) = \bigvee_{\gamma \in \Gamma} (\mathfrak{F} \wedge \mathfrak{F}_\gamma)$. We also have $\mathfrak{F} \vee (\bigwedge_{\gamma \in \Gamma} \mathfrak{F}_\gamma) \leq \bigwedge_{\gamma \in \Gamma} (\mathfrak{F} \vee \mathfrak{F}_\gamma)$. However, if \underline{S} is not finite, we need not have equality. A particular example can be found by letting $\Gamma = S$, $\mathfrak{F}_\gamma = \overline{\{\gamma, x\}}$, and $\mathfrak{F} = \{A: x \in A, A \text{ cofinite}\}$. For in this case $\{s\} \in \bigwedge_{\gamma \in \Gamma} (\mathfrak{F} \vee \mathfrak{F}_\gamma)$ is not cofinite. Thus $(\underline{F}(x), \leq)$ is distributive only in the trivial case where S , and consequently $\underline{F}(x)$, is finite.

3. Structure of (\underline{F}, \leq) and $(\underline{p}(S), \leq)$. The results of § 2 carry directly over to the lattice (\underline{F}, \leq) . For letting $\mathfrak{F} = \prod_{x \in S} \mathfrak{F}_x$, $\mathfrak{G} = \prod_{x \in S} \mathfrak{G}_x$ with $\mathfrak{F}_x, \mathfrak{G}_x \in \underline{F}(x)$ for each x , we see that $\mathfrak{F} \leq \mathfrak{G}$ if and only if $\mathfrak{F}_x \leq \mathfrak{G}_x$ in $(\underline{F}(x), \leq)$ for each x , while $\mathfrak{F} \wedge \mathfrak{G} = \prod_{x \in S} (\mathfrak{F}_x \wedge \mathfrak{G}_x)$ and $\mathfrak{F} \vee \mathfrak{G} = \prod_{x \in S} (\mathfrak{F}_x \vee \mathfrak{G}_x)$. We summarize these results in the following proposition. Each of its components follows from the corresponding result in § 2.

PROPOSITION 6. 1. (\underline{F}, \leq) is complete, modular, and distributive. It is infinitely distributive only in the trivial case of S , and consequently \underline{F} , being finite.

2. (\underline{F}, \leq) is atomic (coatomic). $\mathfrak{F} = \prod_{x \in S} \mathfrak{F}_x \in \underline{F}$ is an atom (coatom) if and only if $\mathfrak{F}_x = \overline{S}$ for $x \neq s$ ($\mathfrak{F}_x = \overline{x}$ for $x \neq s$) and \mathfrak{F}_s is an atom of $\underline{F}(s)$ (a coatom of $\underline{F}(s)$).

3. $\mathfrak{F} \in \underline{F}$ is compact (co-compact) if and only if \mathfrak{F}_x is compact (co-compact) for each $x \in \overline{S}$ and $\mathfrak{F}_x = \overline{S}$ ($\mathfrak{F}_x = \overline{x}$) except for most a finite number of the $x \in S$.

4. (\underline{F}, \leq) is compactly generated.

5. (\underline{F}, \leq) is co-compactly generated if and only if S is finite.

6. \mathfrak{F} has a complement $\mathfrak{G} = \prod_{x \in S} \mathfrak{G}_x$ if and only if \mathfrak{F}_x and \mathfrak{G}_x are complements for each $x \in S$.

7. (\underline{F}, \leq) is complemented if and only if S is finite. In this case complements will be unique.

Using the isomorphism φ , these results immediately carry over to $(\underline{p}(S), \leq)$. Thus we have

THEOREM 2. $(\underline{p}(S), \leq)$ is always complete, modular, distributive, atomic, coatomic, and compactly generated. It is complemented (and has unique complements), co-compactly generated, and infinitely distributive if and only if S is finite.

REFERENCES

1. N. Bourbaki, *Topologie Générale*, Hermann, Paris 1965.
2. D. Kent, *Convergence functions and their related topologies*, *Fund. Math.* **54** (1964).
3. G. Szász, *Introduction to lattice theory*, Academic Press, New York, 1963.

Received March 31, 1968, and in revised form June 17, 1968.

WASHINGTON STATE UNIVERSITY

Pacific Journal of Mathematics

Vol. 29, No. 1

May, 1969

Jorge Alvarez de Araya, <i>A Radon-Nikodým theorem for vector and operator valued measures</i>	1
Deane Eugene Arganbright, <i>The power-commutator structure of finite p-groups</i>	11
Richard Eugene Barlow, Albert W. Marshall and Frank Proschan, <i>Some inequalities for starshaped and convex functions</i>	19
David Clarence Barnes, <i>Some isoperimetric inequalities for the eigenvalues of vibrating strings</i>	43
David Hilding Carlson, <i>Critical points on rim-compact spaces</i>	63
Allan Matlock Weber Carstens, <i>The lattice of pretopologies on an arbitrary set S</i>	67
S. K. Chatterjea, <i>A bilateral generating function for the ultraspherical polynomials</i>	73
Ronald J. Ensey, <i>Primary Abelian groups modulo finite groups</i>	77
Harley M. Flanders, <i>Relations on minimal hypersurfaces</i>	83
Allen Roy Freedman, <i>On asymptotic density in n-dimensions</i>	95
Kent Ralph Fuller, <i>On indecomposable injectives over artinian rings</i>	115
George Isaac Glauberman, <i>Normalizers of p-subgroups in finite groups</i>	137
William James Heinzer, <i>On Krull overrings of an affine ring</i>	145
John McCormick Irwin and Takashi Ito, <i>A quasi-decomposable abelian group without proper isomorphic quotient groups and proper isomorphic subgroups</i>	151
Allan Morton Krall, <i>Boundary value problems with interior point boundary conditions</i>	161
John S. Lowndes, <i>Triple series equations involving Laguerre polynomials</i>	167
Philip Olin, <i>Indefinability in the arithmetic isolic integers</i>	175
Ki-Choul Oum, <i>Bounds for the number of deficient values of entire functions whose zeros have angular densities</i>	187
R. D. Schafer, <i>Standard algebras</i>	203
Wolfgang M. Schmidt, <i>Irregularities of distribution. III</i>	225
Richard Alfred Tapia, <i>An application of a Newton-like method to the Euler-Lagrange equation</i>	235