

Pacific Journal of Mathematics

GEOMETRIES ON SURFACES

HELMUT R. SALZMANN

GEOMETRIES ON SURFACES

HELMUT SALZMANN

Among the topological geometries, two classes have so far attracted special attention, namely the locally compact, 2-dimensional projective and affine planes. Such a plane has a pointset which is homeomorphic to the pointset of the real arguesian plane, hence is a 2-dimensional manifold. In this paper, all the 2-manifolds that can carry topological geometries with connected lines will be determined.

THEOREM. *Let M be a surface carrying a topological geometry such that any two distinct points are on a unique line. Then either*

- (1) *M is an open disk, and each line is open, i.e., homeomorphic to the real line R , or*
- (2) *M is a compact surface of characteristic 1, and each line is closed, i.e., homeomorphic to a circle, or*
- (3) *M is a Moebius strip, and through each point there pass closed lines and at least one open line.*

The proof follows immediately by combining statements 1.10 and 2.3, 6, 9, 10, 13 below. From 2.9 and 2.11 we get

COROLLARY. *M is orientable if and only if the space \mathfrak{L} of lines is a Moebius strip, and M is nonorientable if and only if \mathfrak{L} is compact.*

1. Let \mathfrak{L} be a family of subsets of a nontrivial topological space M , and assume that each two distinct "points" $p, q \in M$ are joined by a unique "line" $L = p \cup q \in \mathfrak{L}$. The system $E = (M, \mathfrak{L})$ is called [3] a "plane" whenever \mathfrak{L} can be provided with a topology (necessarily unique) for which E becomes a topological geometry in the following sense:

- (a) $p \cup q$ depends continuously on (p, q) , and
- (b) the set \mathfrak{D} of pairs of intersecting lines is open in $\mathfrak{L} \times \mathfrak{L}$, and intersection is a continuous map from \mathfrak{D} onto M .

Condition (b) is equivalent to

- (b') If $U \subseteq M$ is open, then $\{(K, L); K \cap L \in U\}$ is open in $\mathfrak{L} \times \mathfrak{L}$.

We shall be concerned with "flat" planes only, i.e., with those planes in which the underlying space M is a 2-dimensional manifold or "surface". The line space \mathfrak{L} of a flat plane is also a surface, and the incident point-line pairs or "flags" form a 3-dimensional closed submanifold F of $M \times \mathfrak{L}$; each line is closed in M and is locally homeomorphic to the real line [3, 2.3].

If B is a connected open subset of M , then the system

$$E_B = (B, \mathfrak{L}_B = \{L \cap B \neq \emptyset; L \in \mathfrak{L}\})$$

is again a flat plane. This shows that, in general, the lines of a geometry will not be connected sets; it is therefore reasonable to determine only those flat planes that have connected lines, i.e. planes satisfying

(c) every line is homeomorphic to the real line or to the circle. Each line can then be provided with two opposite orientations, and the space \mathfrak{L}^+ of oriented lines becomes a two-fold covering of \mathfrak{L} . The lines homeomorphic to \mathbb{R} will be called "open", and the others compact or "closed".

LEMMA 1.1. *A compact disk $D \subseteq M$ cannot contain any line.*

Proof. A line $K \subseteq D$ would have to be a closed line. Hence K would separate some points $a \in D$ and $b \in M$, and then $a \cup b$ would intersect K twice.

DEFINITION 1.2. A set $C \subseteq M$ is convex if $L \cap C$ is connected for every line L . If the boundary $\text{bd } C$ consists of finitely many, resp. of 4 or 3 line segments, C will be called a polygon resp. a quadrangle or a triangle.

LEMMA 1.3. *Each point $p \in M$ has a neighborhood basis consisting of quadrangles (triangles).*

Proof. Let W be a neighborhood of p . Choose points a_1, a_2 such that $p \notin a_1 \cup a_2$, and let J_k be a sufficiently small connected neighborhood of p on $a_k \cup p$. Then

$$U = \{(x_1 \cup a_2) \cap (x_2 \cap a_1); x_k \in J_k\} \subseteq W$$

is a neighborhood of p homeomorphic to $J_1 \times J_2$, which is easily seen to be a convex quadrangle.

COROLLARY 1.4. *The space \mathfrak{L}_p^+ of oriented lines through p is homeomorphic to a circle.*

Convergence in M and in \mathfrak{L} are related by

LEMMA 1.5. $L_n \xrightarrow[n]{} L$ if and only if $L = \{x; \vee x_n \in L_n x_n \xrightarrow[n]{} x\}$.

This follows immediately from the continuity of join and intersection, (a) and (b).

A set is termed "bounded" if it has compact closure.

LEMMA 1.6. *A set \mathfrak{S} of lines is bounded in \mathfrak{L} if and only if there is a bounded set $C \subseteq M$ intersecting every line $S \in \mathfrak{S}$.*

Proof. Let $S_n \in \mathfrak{S}$ and $x_n \in S_n \cap C$. We may assume $x_n \xrightarrow{n} x$. Choose a triangle with boundary T containing x in its interior; then for each sufficiently large n there exists a point $t_n \in S_n \cap T$. The t_n accumulate at some $t \in T$, and a subsequence of the S_n converges to $t \cup x$. For the converse note that M is a union of a countable family of compact subsets.

LEMMA 1.7. *Let K be a closed line, and $B \subseteq M$ a bounded open set containing K . Then $\mathfrak{B} = \{L \in \mathfrak{L}; L \subseteq B\}$ is an open neighborhood of K in \mathfrak{L} consisting of closed lines only.*

Proof. Since bounded lines are closed, \mathfrak{B} cannot contain any open line. If \mathfrak{B} were not a neighborhood of K , we would find a sequence of lines $L_n \xrightarrow{n} K$ with $L_n \cap B \neq \emptyset$ and $L_n \notin \mathfrak{B}$. By (c) each line L_n would contain a point x_n of the compact boundary $\text{bd } B$, in contradiction to 1.5.

COROLLARY 1.8. *The space \mathfrak{R} of closed lines is open in \mathfrak{L} .*

LEMMA 1.9. *A closed line K intersects every line $L \in \mathfrak{L}$.*

Proof. For $a \in K$ the map $\alpha = (x \mapsto a \cup x)$ maps K into the set \mathfrak{L}_a of lines through a . Since \mathfrak{L}_a is homeomorphic to a circle by 1.4, and α is one-to-one, α is even a bijection.

COROLLARY 1.10. *If M is a compact surface, then (M, \mathfrak{L}) is a projective plane, and M is homeomorphic to the real projective plane.*

Proof. Any two different lines intersect by 1.9, and this is the first assertion. The second statement follows now from [2].

PROPOSITION 1.11. *If a geometry contains a closed line K , then each point lies on a closed line.*

Proof. Assume that each line through s is homeomorphic to R , and let $s \cup z^+$ denote the line $s \cup x$ oriented from s to x . Then $(x \mapsto s \cup x^+): K \rightarrow \mathfrak{L}_s^+$ would be a homeomorphism from K onto a set of oriented lines through s containing exactly one of any two lines of opposite orientation. This contradicts 1.4.

2. From now on we shall assume for the sake of simplicity that

M is a surface of finite connectivity, i.e., that M can be embedded in a compact surface \bar{M} by adding a finite number of "endpoints" e_1, \dots, e_r .

LEMMA 2.1. *Let $g: R \rightarrow M$ represent an oriented open line $L = g(R)$. Then L has unique endpoints*

$$e^\pm(L) = \lim_{t \rightarrow \pm\infty} g(t)$$

in \bar{M} .

Proof. Choose small disjoint open neighborhoods U_k of e_k in \bar{M} . Since each line is closed in M , there exists an $s \in R$ such that the connected set $\{g(t); t \geq s\}$ is contained in $\bigcup_{k=1}^r U_k$ and hence in one of the U_k .

LEMMA 2.2. *The oriented lines L having a given endpoint $e^+(L) = e$ form a closed subset of \mathfrak{L}^+ .*

Proof. Let $L_n \xrightarrow{n} L$ with $e^+(L_n) = e$, and choose a small neighborhood U of e in \bar{M} . Then all the L_n intersect the compact boundary $\text{bd } U \subseteq M$. Hence $L \cap \text{bd } U \neq \emptyset$ by 1.5, and $e^+(L) = e$.

LEMMA 2.3. *For each $a \in M$ and each endpoint e there is an oriented line $L \in \mathfrak{L}_a^+$ with $e^+(L) = e$.*

Proof. Let $b_n \xrightarrow{n} e$ in \bar{M} , and $a \cup b_n \xrightarrow{n} L$.

COROLLARY 2.4. *Let (M, \mathfrak{L}) be a flat plane with connected lines. Then M is compact if and only if each line is closed.*

PROPOSITION 2.5. *A surface M carrying a plane geometry with connected lines has at most one endpoint.*

Proof. $\mathfrak{L}_{ik} = \{L \in \mathfrak{L}^+; e^-(L) = e_i, e^+(L) = e_k\}$ is closed in \mathfrak{L}^+ by 2.2. Assume $r > 1$, choose small disjoint neighborhoods U_k of e_k in \bar{M} , and let $b_{kn} \xrightarrow{n} e_k$ in U_k . For $i \neq k$ the lines $b_{in} \cup b_{kn}$ meet the compact complement C of $\bigcup_{k=1}^r U_k$, and hence accumulate at a line in \mathfrak{L}_{ik} . Since \mathfrak{L}^+ is connected, the boundary $\text{bd } \mathfrak{L}_{ik} \neq \emptyset$, and there is a sequence of closed lines K_n converging to a line $L \in \mathfrak{L}_{ik}$. For large n a line K_n will contain two arcs joining U_i and U_k ; thus a line H intersecting L at a point of C would meet some line K_n twice, a contradiction.

PROPOSITION 2.6. *A plane geometry on an orientable surface M*

has no compact lines.

Proof. A compact line K on an orientable surface M would have a bounded neighborhood $B \subseteq M$ which is homeomorphic to a cylinder. By 1.7 the set \mathfrak{B} of all lines $L \subseteq B$ is an open neighborhood of K in \mathfrak{L} , and each $L \in \mathfrak{B}$ is closed. Moreover, it follows from 1.1 and 1.9 that a line $L \subseteq B$ decomposes the cylinder and intersects K . But then K and L would have to intersect twice.

PROPOSITION 2.7. The complement of an open line L is homeomorphic to an open subset of R^2 .

Proof. For $a, b \in L$ consider

$$(x \mapsto (a \cup x, b \cup x)): M - L \rightarrow (\mathfrak{L}_a - \{L\}) \times (\mathfrak{L}_b - \{L\}) .$$

PROPOSITION 2.8. If M is orientable, then any line L decomposes M into two open half-planes, each of which is homeomorphic to an open disk. Consequently \bar{M} is homeomorphic to a 2-sphere.

Proof. Let $L = a \cup b$, and choose homeomorphisms

$$\alpha: \mathfrak{L}_a - \{L\} \rightarrow (0, 1) , \quad \beta: \mathfrak{L}_b - \{L\} \rightarrow (0, 1)$$

corresponding to the same orientation of M . For $\xi = A^\alpha$ let

$$\delta(\xi) = \inf \{B^\beta; A \cap B = \emptyset\} .$$

Then δ is a monotonically increasing, left-continuous mapping. According to the orientation of M one of the half-planes defined by L will be mapped by

$$x \mapsto ((a \cup x)^\alpha, (b \cup x)^\beta)$$

onto the Jordan domain

$$D = \{(\xi, \eta) \in R^2; 0 < \xi < 1, 0 < \eta < \delta(\xi)\} .$$

COROLLARY 2.9. Let (M, \mathfrak{L}) be a flat plane with connected lines. If M is orientable, then M is homeomorphic to R^2 , and \mathfrak{L} is a Moebius strip.

For the last assertion see [1, § 3(6)] or [3, 2.14].

An argument completely analogous to the proof of 2.8 shows

COROLLARY 2.10. Each point of a plane geometry on a non-

orientable surface is incident with a compact line.

THEOREM 2.11. *The line space of a plane geometry (M, \mathfrak{L}) on a nonorientable surface is a compact surface of characteristic 1; the space of oriented lines is homeomorphic to a 2-sphere.*

Proof. If H, K are closed lines through a , then $\mathfrak{L} - \mathfrak{L}_a$ is homeomorphic to $(H - \{a\}) \times (K - \{a\})$.

LEMMA 2.12. *In a plane geometry (M, \mathfrak{L}) on a nonorientable surface the set of closed lines through a point a is connected.*

Proof. Because of 2.5 we may assume that M has exactly one endpoint e . Choose an open line L and two closed lines H and K through a . We have to show that no open line through a is separated from L by H, K . By 2.7 the Jordan curve theorem holds in $M - L$. Hence the union $H \cup K$ decomposes $M - L$ into two connected subsets, only one of which contains points of a small neighborhood U of e . Therefore, a line G which is separated from L by H, K must be disjoint from U and hence is compact.

THEOREM 2.13. *An open nonorientable surface M carrying a plane geometry with connected lines is a Moebius strip.*

Proof. The dual (\mathfrak{L}, M) of the geometry defined by the system \mathfrak{R} of closed lines on M is a flat plane by 2.10, 1.8, 1.9, and 2.12. The theorem follows now from 2.9 and 2.11.

Note that in general the lines of a topological geometry on a Moebius strip need not be connected.

The author is indebted to James Dugundji for very stimulating discussions and valuable comments.

REFERENCES

1. H. Busemann and H. Salzmann, *Metric collineations and inverse problems*, Math. Zeit. **87** (1965), 214-240.
2. H. Salzmann, *Topologische Struktur zweidimensionaler projektiver Ebenen*, Math. Zeit. **71** (1959), 408-413.
3. ———, *Topological planes*, Advances in Math. **2** (1967), 1-60.

Received March 5, 1968.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R. PHELPS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics

Vol. 29, No. 2

June, 1969

Bruce Langworthy Chalmers, <i>On boundary behavior of the Bergman kernel function and related domain functionals</i>	243
William Eugene Coppage, <i>Peirce decomposition in simple Lie-admissible power-associative rings</i>	251
Edwin Duda, <i>Compactness of mappings</i>	259
Earl F. Ecklund Jr., <i>On prime divisors of the binomial coefficient</i>	267
Don E. Edmondson, <i>A modular topological lattice</i>	271
Phillip Alan Griffith, <i>A note on a theorem of Hill</i>	279
Marcel Herzog, <i>On finite groups with independent cyclic Sylow subgroups</i>	285
James A. Huckaba, <i>Extensions of pseudo-valuations</i>	295
S. A. Huq, <i>Semivarieties and subfunctors of the identity functor</i>	303
I. Martin (Irving) Isaacs and Donald Steven Passman, <i>Finite groups with small character degrees and large prime divisors. II</i>	311
Carl Kallina, <i>A Green's function approach to perturbations of periodic solutions</i>	325
Al (Allen Frederick) Kelley, Jr., <i>Analytic two-dimensional subcenter manifolds for systems with an integral</i>	335
Alistair H. Lachlan, <i>Initial segments of one-one degrees</i>	351
Marion-Josephine Lim, <i>Rank k Grassmann products</i>	367
Raymond J. McGivney and William Henry Ruckle, <i>Multiplier algebras of biorthogonal systems</i>	375
J. K. Oddson, <i>On the rate of decay of solutions of parabolic differential equations</i>	389
Helmut R. Salzmann, <i>Geometries on surfaces</i>	397
Annemarie Schlette, <i>Artinian, almost abelian groups and their groups of automorphisms</i>	403
Edgar Lee Stout, <i>Additional results on modules over polydisc algebras</i>	427
Lajos Tamássy, <i>A characteristic property of the sphere</i>	439
Mark Lawrence Teply, <i>Some aspects of Goldie's torsion theory</i>	447
Freddie Eugene Tidmore, <i>Extremal structure of star-shaped sets</i>	461
Leon Jarome Weill, <i>Unconditional and shrinking bases in locally convex spaces</i>	467