

# Pacific Journal of Mathematics

**CONDITIONS FOR A MAPPING TO HAVE THE SLICING  
STRUCTURE PROPERTY**

GERALD S. UNGAR

## CONDITIONS FOR A MAPPING TO HAVE THE SLICING STRUCTURE PROPERTY

GERALD S. UNGAR

Let  $p: E \rightarrow B$  be a fibering in the sense of Serre. As is well known the fibering need not be a fibering in any stronger sense. However it is expected that if certain conditions are placed on  $E, p$  or  $B$  then  $p$  might be a fibration in a stronger sense. This paper gives such conditions.

The main result of this paper is:

**THEOREM 1.** Let  $p$  be an  $n$ -regular perfect map from a complete metric space  $(E, d)$  onto a locally equiconnected space  $B$ . If  $\dim E \times B \leq n$  then  $p$  has the slicing structure property (in particular  $p$  is a Hurewicz fibration).

The following definitions will be needed.

**DEFINITION 1.** A space  $X$  is *locally equiconnected* if for each point  $x$ , there exists a neighborhood  $U_x$  of  $x$  and a map

$$N: U_x \times U_x \times I \rightarrow X$$

satisfying  $N(a, b, 0) = a$ ,  $N(a, b, 1) = b$ , and  $N(a, a, t) = a$ .

**DEFINITION 2.** A map  $p$  from  $E$  to  $B$  is  $n$ -regular if it is open and satisfies the following property: given any  $x$  in  $E$  and any neighborhood  $U$  of  $x$  there exists a neighborhood  $V$  of  $x$  such that if  $f: S^m \rightarrow V \cap p^{-1}(y)$  for some  $y \in B$  ( $m \leq n$ ) then there exists

$$F: B^{m+1} \rightarrow U \cap p^{-1}(y)$$

which is an extension of  $f$ .

**DEFINITION 3.** A family  $\mathcal{S}$  of sets of  $Y$  is *equi- $LC^n$*  if for every  $y \in S \in \mathcal{S}$  and every neighborhood  $U$  of  $y$  in  $Y$  there exists a neighborhood  $V$  of  $y$  such that for every  $S \in \mathcal{S}$ , every continuous image of an  $m$ -sphere ( $m \leq n$ ) in  $S \cap V$  is contractible in  $S \cap U$ .

*Note 1.* If  $p: E \rightarrow B$  is  $n$ -regular then the collection  $\{p^{-1}(b) \mid b \in B\}$  is *equi- $LC^n$* .

**DEFINITION 4.** A family  $\mathcal{S}$  of sets of a metric space  $(Y, d)$  is *uniformly equi- $LC^n$*  with respect to  $d$  if given  $\varepsilon > 0$  there exists  $\delta > 0$  such that if  $f: S^m \rightarrow S \cap N(x, \delta)$  ( $m \leq n$  and  $S \in \mathcal{S}$ ) then there exists  $F: B^{m+1} \rightarrow S \cap N(x, \varepsilon)$  which is an extension of  $f$ .

**DEFINITION 5.** A map  $p: E \rightarrow B$  has the *covering homotopy pro-*

property for a class of spaces if given any space  $X$  in the class and maps  $F: X \times I \rightarrow B$  and  $g: X \rightarrow E$  such that  $F(x, 0) = pg(x)$  then there exists a map:  $G: X \times I \rightarrow E$  such that  $pG = F$  and  $G(x, 0) = g(x)$ .

DEFINITION 6. A map  $p: E \rightarrow B$  is a *Serre fibration* if  $p$  has the covering homotopy property for the class of polyhedra. It is a *Hurewicz fibration* if it has the covering homotopy property for all spaces.

DEFINITION 7. A map  $p: E \rightarrow B$  has the *slicing structure property* (SSP) if for each point  $b \in B$  there exists a neighborhood  $U_b$  of  $b$  and a map  $\psi_b: p^{-1}(U_b) \times U_b \rightarrow p^{-1}(U_b)$  such that (1)  $\psi_b(e, p(e)) = e$  and (2)  $p\psi_b = \pi_2$  (the projection onto  $U_b$ ).

DEFINITION 8. A function  $\varphi: X \rightarrow 2^Y$  ( $Y$  metric) is continuous if given  $\varepsilon > 0$ ; every  $x_0 \in x$  has a neighborhood  $U$  such that for every  $x \in U$ ,  $\varphi(x_0) \subset N_\varepsilon(\varphi(x))$  and  $\varphi(x) \subset N_\varepsilon(\varphi(x_0))$ .

DEFINITION 9. A selection for a function  $\varphi: X \rightarrow 2^Y$  is a map  $g: X \rightarrow Y$  such that  $g(x) \in \varphi(x)$ .

A mapping is a continuous function. All spaces will be Hausdorff. The  $n$ -dimensional sphere will be denoted by  $S^n$  and the ball which it bounds  $B^{n+1}$ . If  $f$  is a mapping  $\text{Gr}(f)$  will denote the graph of  $f$ .

The following theorem of Michael will be needed:

THEOREM M. Let  $Z$  be paracompact, let  $X = Z \times I$  and let  $Y$  be a complete metric space with metric  $\rho$ . Let  $\mathcal{S} \subset 2^Y$  be uniformly equi-LC<sup>n</sup> with respect to  $\rho$  and let  $\varphi: X \rightarrow \mathcal{S}$  be continuous with respect to  $\rho$ . Let  $\dim Z \leq n$  and let  $A = (Z \times 0) \cup (C \times I)$  where  $C$  is closed in  $Z$ . Then every selection for  $\varphi|_A$  can be extended to a selection for  $\varphi$ .

## 2. Proof of Theorem 1 and its consequences.

*Proof.* Let  $b_0 \in B$ . Since  $B$  is locally equiconnected at  $b_0$  there exists a neighborhood  $U$  of  $b_0$  and a map  $N_U: U \times U \times I \rightarrow B$  such that  $N_U(x, y, 0) = x$ ,  $N_U(x, y, 1) = y$ , and  $N_U(x, x, t) = x$ . Let  $P_U = p|_{p^{-1}(U)}$  and define  $g: \text{Gr}(p_U) \rightarrow p^{-1}(U)$  by  $g(e, p(e)) = e$ . Also define  $F: p^{-1}(U) \times B \rightarrow B$  by  $F(e, b) = b$  and

$$H: p^{-1}(U) \times U \times I \rightarrow p^{-1}(U) \times B$$

by  $H(e, b, t) = (e, N_U(p(e), b, t))$ . Note  $H(e, b, 0) = (e, N_U(p(e), b, 0)) = (e, p(e))$  and  $H(e, b, 1) = (e, N_U(p(e), b, 1)) = (e, b)$ . Further define


$$g': (p^{-1}(U) \times U \times 0) \cup (\text{Gr}(P_U) \times I) \rightarrow E$$

by  $g'(e, b, t) = e$  and  $K: p^{-1}(U) \times U \times I \rightarrow B$  by

$$K(e, b, t) = F(H(e, b, t))$$

and note that  $pg' = K|_{(p^{-1}(U) \times U \times 0) \cup \text{Gr}(P_U) \times I}$ .

Therefore we have the following commutative diagram.



Now Theorem  $M$  will be applied. Let  $Z = p^{-1}(U) \times U$ ,  $Y = E$ , and  $\varphi: Z \times I \rightarrow \mathcal{S} \subset 2^Y$  be defined by  $\varphi(z, t) = p^{-1}K(z, t)$  and let  $C = \text{Gr}(P_U)$ . Note  $Z$  is paracompact and  $\varphi$  is continuous since  $p$  is perfect. Since  $p$  is  $n$ -regular  $\{p^{-1}(b)\}$  in equi- $LC^n$  and by Proposition 2.1 [3] there exists a metric  $\sigma$  on  $E$  agreeing with the topology such that  $\sigma \geq d$  and  $\{p^{-1}(b)\}$  is uniformly equi- $LC^n$ . Since  $\sigma \geq d$ ,  $(E, \sigma)$  is a complete metric space. It should also be noted that  $\dim Z \leq n$  and that  $g'$  is a selection for  $\varphi|_{(Z \times 0) \cup (C \times I)}$ . Hence by Theorem  $M$ ,  $g'$  could be extended to a selection  $G$  for  $\varphi$  (i.e.,

$$G: p^{-1}(U) \times U \times I \rightarrow E$$

in such a way that the above diagram will still be commutative with the addition of  $G$ ).

Define  $\varphi_U: p^{-1}(U) \times U \rightarrow p^{-1}(U)$  by  $\varphi_U(e, b) = G(e, b, 1)$ . Note if  $(e, b) \in p^{-1}(U) \times U$  then

$$\begin{aligned} G(e, b, 1) &\in p^{-1}K(e, b, 1) = p^{-1}FH(e, b, 1) = p^{-1}F(e, N_U(p(e), b, 1)) \\ &= p^{-1}F(e, b) = p^{-1}(b) \in p^{-1}(U) . \end{aligned}$$

Hence the range of  $\varphi_U$  is as stated. It is now easy to see that  $\varphi_U$  satisfies the conditions to be a slicing function. This completes the proof.

*Note 2.* The hypothesis that  $p$  be perfect was used only to show that  $\{p^{-1}(b) | b \in B\}$  is a continuous collection and that  $B$  is paracom-

pact. Hence if this could be shown some other way a stronger theorem will be obtained.

**COROLLARY 1.** *If  $p: E \rightarrow B$  is a Serre fibration and  $E$  and  $B$  are finite dimensional compact ANR's then  $p$  has the SSP.*

*Proof.* It is well known that ANR's are locally equiconnected.

It also follows from [2] that  $p$  is  $n$ -regular for all  $n$ . Hence the proof follows from Theorem 1.

Theorem 1 and Corollary 1 allow us to get the following generalizations of Raymond's results in [5].

**COROLLARY 2.** *Let  $p: E \rightarrow B$  be a Serre fibration of a connected compact metric finite dimensional ANR onto a compact metric finite dimensional ANR. Suppose that  $E$  is an  $n$ -gm over  $L$  (a field or the integers). Then:*

- (a) *each fiber  $F_b$  is a  $k$ -gm over  $L$*
- (b)  *$B$  is an  $(n-k) - gm$  over  $L$ .*

**COROLLARY 3.** *Let  $p: E \rightarrow B$  is a Serre fibration of a connected compact metric finite dimensional ANR onto a compact metric finite dimensional ANR base  $B$ . Suppose that  $E$  is a (generalized) manifold (over a principal ideal domain) and some fiber has a component of dimension  $\leq 2$ . Then  $p$  is locally trivial.*

Another theorem which follows from Michael's Theorem 1.2 [3] is the following:

**THEOREM 2.** *Let  $p: E \rightarrow B$  be an  $n$ -regular map from a complete metric space  $E$  onto a paracompact space  $B$ . Assume that*

$$\dim E \times B \leq n + 1$$

*and  $p^{-1}(b)$  is  $C^n$  for every  $b \in B$ . Then  $p$  has the SSP and the slicing structure could be chosen with only one slicing function.*

*Proof.* Define  $g: \text{Gr}(p) \rightarrow E$  by  $g(e, p(e)) = e$  and  $F: E \times B \rightarrow B$  by  $F(e, b) = e$ . The  $\varphi(e, b) = p^{-1}F(e, b)$  is a carrier and  $g$  is a selection for  $\varphi | \text{Gr}(p)$ . Hence by Theorem 1.2 [3]  $g$  could be extended to a selection  $G$  for  $\varphi$ . It is easily seen that  $G$  is the desired slicing function.

*Note 3.* Theorem 2 has corollaries similar to those of Theorem 1 and the author leaves them to the reader to develop.

## REFERENCES

1. S. T. Hu, *Homotopy theory*, Academic Press, New York, 1959.
2. L. McAuley and P. Tulley, *Fiber spaces and N-regularity*, Topology Seminar Wisconsin 1965, Edited by R. Bing and R. Bean, Ann. of Math Studies, No. 60, Princeton University Press, Princeton, N. J., 1966.
3. E. Michael, *Continuous selections II*, Ann. of Math. **64** (1956), 567-580.
4. ———, *Continuous selections III*, Ann. of Math. **65** (1957), 375-390.
5. F. Raymond *Local triviality for Hurewicz fiberings of manifolds*, Topology **3** (1956), 43-57.
6. G. S. Ungar, *Light fiber maps*, Fund. Math. **62** (1968), 31-45.

Received October 30, 1968.

CASE WESTERN RESERVE UNIVERSITY  
CLEVELAND, OHIO



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. ROYDEN  
Stanford University  
Stanford, California

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. R. PHELPS  
University of Washington  
Seattle, Washington 98105

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL WEAPONS CENTER

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.



# Pacific Journal of Mathematics

Vol. 30, No. 2

October, 1969

|  |     |
|--|-----|
| Gregory Frank Bachelis, <i>Homomorphisms of annihilator Banach algebras. II</i> .....  | 283 |
| Leon Bernstein and Helmut Hasse, <i>An explicit formula for the units of an algebraic number field of degree <math>n \geq 2</math></i> .....                               | 293 |
| David W. Boyd, <i>Best constants in a class of integral inequalities</i> .....   | 367 |
| Paul F. Conrad and John Dauns, <i>An embedding theorem for lattice-ordered fields</i> .....  | 385 |
| H. P. Dikshit, <i>Summability of Fourier series by triangular matrix transformations</i> .....   | 399 |
| Dragomir Z. Djokovic, <i>Linear transformations of tensor products preserving a fixed rank</i> .....   | 411 |
| John J. F. Fournier, <i>Extensions of a Fourier multiplier theorem of Paley</i> ....   | 415 |
| Robert Paul Kopp, <i>A subcollection of algebras in a collection of Banach spaces</i> .....  | 433 |
| Lawrence Louis Larmore, <i>Twisted cohomology and enumeration of vector bundles</i> .....  | 437 |
| William Grenfell Leavitt and Yu-Lee Lee, <i>A radical coinciding with the lower radical in associative and alternative rings</i> .....                                     | 459 |
| Samuel Merrill and Nand Lal, <i>Characterization of certain invariant subspaces of <math>H^p</math> and <math>L^p</math> spaces derived from logmodular algebras</i> ..... | 463 |
| Sam Bernard Nadler, Jr., <i>Multi-valued contraction mappings</i> .....  | 475 |
| T. V. Panchapagesan, <i>Semi-groups of scalar type operators in Banach spaces</i> .....  | 489 |
| J. W. Spellmann, <i>Concerning the infinite differentiability of semigroup motions</i> .....   | 519 |
| H. M. (Hari Mohan) Srivastava, <i>A note on certain dual series equations involving Laguerre polynomials</i> .....   | 525 |
| Ernest Lester Stitzinger, <i>A nonimbedding theorem of associative algebras</i> .....  | 529 |
| J. Jerry Uhl, Jr., <i>Martingales of vector valued set functions</i> .....   | 533 |
| Gerald S. Ungar, <i>Conditions for a mapping to have the slicing structure property</i> .....  | 549 |
| John Mays Worrell Jr., <i>On continuous mappings of metacompact Čech complete spaces</i> .....   | 555 |