

# Pacific Journal of Mathematics

**ON THE INVERSION FORMULA FOR THE CHARACTERISTIC  
FUNCTION**

T. KAWATA

## ON THE INVERSION FORMULA FOR THE CHARACTERISTIC FUNCTION

T. KAWATA

**In the inversion formula**

$$F(x) - F(0) = \lim_{T \rightarrow \infty} \frac{1}{2\pi} \int_{-T}^T \frac{e^{-ixt} - 1}{-it} f(t) dt$$

**for the characteristic function  $f(t)$  of a distribution function  $F(x)$ , the limit of the symmetric integral is used. The purpose of this paper is to give a necessary and sufficient condition for the existence of the asymmetric improper integral  $\lim_{T, T' \rightarrow \infty} \int_{-T'}^T$  on the right of the above formula.**

Let  $F(x)$  be a probability distribution function and  $f(t)$  the corresponding characteristic function,

$$(1.1) \quad f(t) = \int_{-\infty}^{\infty} e^{itx} dF(x).$$

We assume in this note that  $F(x)$  is standardized so that

$$(1.2) \quad F(x) = \frac{1}{2}[F(x+0) + F(x-0)].$$

The well known inversion formula states that

$$(1.3) \quad F(x) - F(0) = \lim_{T \rightarrow \infty} \frac{1}{2\pi} \int_{-T}^T \frac{e^{-ixt} - 1}{-it} f(t) dt$$

for every  $-\infty < x < \infty$ .

It is also known that the symmetric integral of the right hand side cannot be replaced by the improper integral  $\lim_{T, T' \rightarrow \infty} \int_{-T'}^T$  ( $T, T'$  going to infinity independently).

Actually we may easily see that

$$(1.4) \quad \operatorname{Re} \left( \frac{1}{2\pi} \int_0^T \frac{e^{-ixt} - 1}{-it} f(t) dt \right) = \operatorname{Re} \left( \frac{1}{2\pi} \int_{-T}^0 \frac{e^{-ixt} - 1}{-it} f(t) dt \right),$$

and

$$(1.5) \quad \operatorname{Im} \left( \frac{1}{2\pi} \int_0^T \frac{e^{-ixt} - 1}{-it} f(t) dt \right) = -\operatorname{Im} \left( \frac{1}{2\pi} \int_{-T}^0 \frac{e^{-ixt} - 1}{-it} f(t) dt \right),$$

and hence  $\frac{1}{2\pi} \int_{-T}^T \frac{e^{-ixt} - 1}{-it} f(t) dt$  cancels out its imaginary part.

The real part (1.4) always converges to  $\frac{1}{2}[F(x) - F(0)]$ . This gives

the proof of (1.3). (See [1], pp. 263–264).

However the imaginary part (1.5) does not necessarily converge without some condition on  $F(x)$ . This is why the limit of the symmetric integral in (1.3) cannot be replaced by the general improper integral.

2. The condition for the existence of the improper integral. We shall give the necessary and sufficient condition for the existence of the limit of (1.5).

**THEOREM 1.** *In order that the limit of (1.5) when  $T \rightarrow \infty$  exists, it is necessary and sufficient that the integral*

$$(2.1) \quad \lim_{\varepsilon \rightarrow 0+} \int_{\varepsilon}^1 \frac{G(u, x) - G(u, 0)}{u} du$$

exists where,

$$(2.2) \quad G(u, x) = F(u + x) - F(-u + x)$$

and if (2.1) exists

$$(2.3) \quad \lim_{T \rightarrow \infty} \operatorname{Im} \left( \frac{1}{2\pi} \int_0^T \frac{e^{-ixt} - 1}{-it} f(t) dt \right) = \int_0^{\infty} \frac{G(u, x) - G(u, 0)}{u} du .$$

It must be noted that the integral of the right hand side of (2.3) exists in the neighborhood of the infinity. In fact  $G(u, x) - G(u, 0) = [F(u + x) - F(u)] - [F(-u + x) - F(-u)]$  and  $F(u + x) - F(u) \in L, (-\infty, \infty)$  for every fixed  $x$ .

We shall now prove the theorem.

Let

$$I(x, T) = \operatorname{Im} \left( \frac{1}{2\pi} \int_0^T \frac{e^{-ixt} - 1}{-it} f(t) dt \right) .$$

We then easily see that

$$\begin{aligned} I(x, T) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} dF(u) \int_0^T \frac{\sin xt \sin ut - (1 - \cos xt) \cos ut}{t} dt \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} dF(u) \int_0^T \frac{\cos(u-x)t - \cos ut}{t} dt \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} dF(u) \int_0^T dt \int_{u-x}^u \sin vt dv \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} dF(u) \int_{u-x}^u \frac{1 - \cos vT}{v} dv \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{1 - \cos vT}{v} dv \int_v^{v+x} dF(u) \\
 &= \frac{1}{2\pi} \int_0^{\infty} [G(v, x) - G(v, 0)] \frac{1 - \cos vT}{v} dv .
 \end{aligned}$$

As was mentioned before,  $G(v, x) - G(v, 0) \in L_1(-\infty, \infty)$ . Hence the Riemann-Lebesgue lemma shows that

$$\lim_{T \rightarrow \infty} \int_{\varepsilon}^{\infty} [G(v, x) - G(v, 0)] \frac{\cos vT}{v} dv = 0$$

for any  $\varepsilon > 0$ . Therefore we may write

$$\begin{aligned}
 (2.4) \quad I(x, T) &= \int_0^{\varepsilon} \frac{G(v, x) - G(v, 0)}{v} (1 - \cos vT) dv \\
 &\quad + \int_{\varepsilon}^{\infty} \frac{G(v, x) - G(v, 0)}{v} dv + o(1)
 \end{aligned}$$

as  $T \rightarrow \infty$ , for a fixed  $\varepsilon > 0$ .

Now we shall show the sufficiency of the condition of the theorem. Let  $\varepsilon > 0$  be arbitrary but fixed. Write

$$\begin{aligned}
 (2.5) \quad &\int_0^{\varepsilon} \frac{G(v, x) - G(v, 0)}{v} (1 - \cos vT) dv \\
 &= \int_0^{1/T} + \int_{1/T}^{\varepsilon} = K_1 + K_2 ,
 \end{aligned}$$

say. We have

$$\begin{aligned}
 (2.6) \quad |K_1| &\leq \int_0^{1/T} |G(v, x) - G(v, 0)| \frac{1 - \cos vt}{v} dv \\
 &\leq CT \int_0^{1/T} |G(v, x) - G(v, 0)| dv ,
 \end{aligned}$$

for some constant  $C$ .

$\lim_{v \rightarrow 0+} [G(v, x) - G(v, 0)]$  exists since  $F$  is nondecreasing and it must be zero, otherwise (2.1) does not exist. Hence the last expression converges to zero.

$$(2.7) \quad K_1 = o(1), \quad \text{as } T \rightarrow \infty .$$

Next write

$$(2.8) \quad \chi(v) = G(v, x) - G(v, 0) .$$

Choose  $\varepsilon$  such that  $|\chi(v)| < \delta$  for  $|v| \leq \varepsilon$  for an arbitrary chosen  $\delta$ . Since  $\chi(v)/v$  is of bounded variation in  $[1/T, \varepsilon]$ , we have, using the second mean value theorem,

$$\begin{aligned} K_2 &= \int_{1/T}^{\varepsilon} \frac{\chi(v)}{v} dv - \int_{1/T}^{\varepsilon} \frac{\chi(v)}{v} \cos vT dv \\ &= \int_{1/T}^{\varepsilon} \frac{\chi(v)}{v} dv - T\chi\left(\frac{1}{T}\right) \int_{1/T}^{\varepsilon} \cos vT dv - \frac{\chi(\varepsilon)}{\varepsilon} \int_{\varepsilon}^{\varepsilon} \cos vT dv \end{aligned}$$

for some  $1/T < \xi < \varepsilon$ . Thus

$$\left| K_2 - \int_{1/T}^{\varepsilon} \frac{\chi(v)}{v} dv \right| \leq 2\chi\left(\frac{1}{T}\right) + 2\chi(\varepsilon) \leq 4\delta .$$

Therefore from (2.4) and (2.5)

$$(2.9) \quad \left| I(x, T) - \frac{1}{2\pi} \int_{1/T}^{\infty} \frac{\chi(v)}{v} dv \right| \leq \frac{2\delta}{\pi} + o(1) .$$

This shows the sufficiency of the condition of the theorem and gives (2.3).

We shall next show the necessity. Define  $\chi(v)$  as before. We see that  $\chi(v)$  has the limit  $c$  as  $v \rightarrow +0$ . If  $c \neq 0$ , then from (2.4)

$$\begin{aligned} I(x, T) - c \int_0^{\varepsilon} \frac{1 - \cos vT}{v} dv \\ = \int_0^{\varepsilon} \frac{[\chi(v) - c](1 - \cos vT)}{v} dv + \int_{\varepsilon}^{\infty} \frac{\chi(v)}{v} dv + o(1) . \end{aligned}$$

The first integral of the right hand side is handled in the same way as in deriving (2.6) and (2.9) with  $\chi(v) - c$  in place of

$$\chi(v) = G(v, x) - G(v, 0) .$$

Actually instead of (2.6) we see that  $K_1$  with  $\phi(v) - c$  is bounded by  $CT \int_0^{1/T} |\chi(v) - c| dv$  which is  $o(1)$ . In place of (2.9) we have

$$(2.10) \quad \left| I(x, T) - \frac{c}{2\pi} \int_0^{\varepsilon} \frac{1 - \cos vT}{v} dv - \frac{1}{2\pi} \int_{1/T}^{\varepsilon} \frac{\chi(v) - c}{v} dv - \frac{1}{2\pi} \int_{\varepsilon}^{\infty} \frac{\chi(v)}{v} dv \right| \leq C_1 \delta + o(1) ,$$

where  $C_1$  is some constant.

$$(2.11) \quad \int_0^{\varepsilon} \frac{1 - \cos vT}{v} dv = 2 \int_0^{\varepsilon T} \frac{\sin^2 v/2}{v} dv \geq C_2 \log \varepsilon T ,$$

where  $C_2$  is an absolute constant. Choose  $\varepsilon$  for an arbitrary given  $\eta < C_2$  so that  $|\chi(v) - c| < \eta$  for  $0 < v < \varepsilon$ . Then

$$(2.12) \quad \left| \int_{1/T}^{\varepsilon} \frac{\chi(v) - c}{v} dv \right| \leq \eta \log \varepsilon T .$$

Hence if  $I(x, T)$  has a limit as  $T \rightarrow \infty$ , then in view of (2.11) and (2.12), (2.10) implies a contradiction. Hence we have that  $c = 0$ . Using (2.10), this yields

$$\left| I(x, T) - \frac{1}{2\pi} \int_{1/T}^{\infty} \frac{\chi(v)}{v} dv \right| \leq C_1 \delta + o(1).$$

This proves the necessity of the condition.

3. Remarks. From Theorem 1, we immediately obtain

THEOREM 2. *In order that*

$$\lim_{T, T' \rightarrow \infty} \frac{1}{2\pi} \int_{-T'}^T \frac{e^{-ixt} - 1}{-it} f(t) dt$$

*exists, it is necessary and sufficient that (2.1) exists for  $\varepsilon > 0$ . (The limit is  $F(x) - F(0)$ ).*

Similar arguments apply to the integral

$$(3.1) \quad J_1(x, T) = \int_1^T \frac{f(t)e^{-ixt}}{it} dt \quad \text{and} \quad J_2(x, T) = \int_{-T}^{-1} \frac{f(t)e^{-ixt}}{it} dt.$$

We easily see that  $J_1$  and  $J_2$  are conjugate complex. We may show that *in order for  $J_1(x, T)$  or  $J_2(x, T)$  to converge as  $T \rightarrow \infty$ , it is necessary and sufficient that*

$$(3.2) \quad \int_0^\varepsilon \frac{F(u+x) - F(-u+x)}{u} du < \infty$$

*for some  $\varepsilon > 0$ .*

(3.1) implies that

$$(3.3) \quad \lim_{T \rightarrow \infty} \frac{1}{2\pi} \int_{-T}^T f(t)e^{-ixt} dt = 0$$

which is very well known when  $F(x)$  is continuous at  $x$ . (3.1) says more than this about the improper integrability of  $f(t)$  near infinity with the additional condition (3.2) on  $F(x)$ .

The sufficiency of (3.2) for the existence of the limits of (3.1) was proved in [2] before.

#### REFERENCES

1. T. Kawata, *The characteristic function of a probability distribution*, Tohoku Math. J. **48** (1941).
2. A. Rényi, *Wahrscheinlichkeitsrechnung*, Berlin, 1962.

Received November 20, 1968. This work was supported under National Science Foundation Grants GP-1577 and GP-3725.



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. ROYDEN  
Stanford University  
Stanford, California

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

RICHARD PIERCE  
University of Washington  
Seattle, Washington 98105

BASIL GORDON  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL WEAPONS CENTER



# Pacific Journal of Mathematics

Vol. 31, No. 1

November, 1969

James Burton Ax, <i>Injective endomorphisms of varieties and schemes</i> . . . . .	1
Richard Hindman Bouldin, <i>A generalization of the Weinstein-Aronszajn formula</i> . . . . .	9
John Martin Chadam, <i>The asymptotic behavior of the Klein-Gordon equation with external potential. II</i> . . . . .	19
Rina Hadass, <i>On the zeros of the solutions of the differential equation <math>y^{(n)}(z) + p(z) = 0</math></i> . . . . .	33
John Sollion Hsia, <i>Integral equivalence of vectors over local modular lattices. II</i> . . . . .	47
Robert Hughes, <i>Boundary behavior of random valued heat polynomial expansions</i> . . . . .	61
Surender Kumar Jain, Saad H. Mohamed and Surjeet Singh, <i>Rings in which every right ideal is quasi-injective</i> . . . . .	73
T. Kawata, <i>On the inversion formula for the characteristic function</i> . . . . .	81
Erwin Kleinfeld, <i>On right alternative rings without proper right ideals</i> . . . . .	87
Robert Leroy Kruse and David Thomas Price, <i>On the subring structure of finite nilpotent rings</i> . . . . .	103
Marvin David Marcus and Stephen J. Pierce, <i>Symmetric positive definite multilinear functionals with a given automorphism</i> . . . . .	119
William Schumacher Massey, <i>Pontryagin squares in the Thom space of a bundle</i> . . . . .	133
William Schumacher Massey, <i>Proof of a conjecture of Whitney</i> . . . . .	143
John William Neuberger, <i>Existence of a spectrum for nonlinear transformations</i> . . . . .	157
Stephen E. Newman, <i>Measure algebras on idempotent semigroups</i> . . . . .	161
K. Chandrasekhara Rao, <i>Matrix transformations of some sequence spaces</i> . . . . .	171
Robert Bruce Schneider, <i>Some theorems in Fourier analysis on symmetric sets</i> . . . . .	175
Ulrich F. K. Schoenwaelder, <i>Centralizers of abelian, normal subgroups of hypercyclic groups</i> . . . . .	197
Jerrold Norman Siegel, <i>G-spaces, H-spaces and W-spaces</i> . . . . .	209
Robert Irving Soare, <i>Cohesive sets and recursively enumerable Dedekind cuts</i> . . . . .	215
Kwok-Wai Tam, <i>Isometries of certain function spaces</i> . . . . .	233
Awadhesh Kumar Tiwary, <i>Injective hulls of semi-simple modules over regular rings</i> . . . . .	247
Eldon Jon Vought, <i>Concerning continua not separated by any nonaposyndetic subcontinuum</i> . . . . .	257
Robert Breckenridge Warfield, Jr., <i>Decompositions of injective modules</i> . . . . .	263