

Pacific Journal of Mathematics

MATRIX TRANSFORMATIONS OF SOME SEQUENCE SPACES

K. CHANDRASEKHARA RAO

MATRIX TRANSFORMATIONS OF SOME SEQUENCE SPACES

K. CHANDRASEKHARA RAO

One of the important investigations in the theory of summability is that of finding necessary and sufficient conditions on an infinite matrix in order that the matrix should transform one (complex) sequence space into the same or another sequence space. In this note some such theorems are given.

Let

C_0 = the space of null sequences;

C = the space of convergent sequences;

Γ = the space of sequences $x = \{x_p\}$ such that $|x_p|^{1/p} \rightarrow 0$, as $p \rightarrow \infty$.

The space Γ can be regarded as the space of all integral functions $f(z) = \sum_{p=1}^{\infty} x_p z^p$;

Γ^* = the space of sequences $s = \{s_p\}$ such that the sequence $\{|s_p|^{1/p}\}$ is bounded. Γ^* may also be considered as the space conjugate to Γ regarded as the space of integral functions $f(z) = \sum_{p=1}^{\infty} x_p z^p$.

Each continuous linear functional $U \in \Gamma^*$ is of the form $U(f) = \sum_{p=1}^{\infty} s_p z^p$.

Let $A = (a_{np})$, $(n, p = 1, 2, \dots)$, be an infinite matrix of complex elements. The A transform of $x = \{x_p\}$, $y = \{y_n\}$ is the sequence defined by the equations

$$(1) \quad y_n = \sum_{p=1}^{\infty} a_{np} x_p, \quad (n = 1, 2, \dots).$$

Here $y = \{y_n\}$ and $x = \{x_p\}$ are complex sequences. Similarly, the A transform of $s = \{s_p\}$, $t = \{t_n\}$ is the sequence defined by the equations

$$(2) \quad t_n = \sum_{p=1}^{\infty} a_{np} s_p, \quad (n = 1, 2, \dots).$$

Here also $t = \{t_n\}$ and $s = \{s_p\}$ are both complex sequences.

The following theorems are true:

THEOREM I. *Let (1) hold. In order that $\{y_n\}$ should belong to Γ whenever $\{x_p\}$ belongs to C_0 , it is necessary and sufficient that (I, 1) the sequence $\{\theta_n\}$ is a null sequence, where*

$$(3) \quad \theta_n = \left(\sum_{p=1}^{\infty} |a_{np}| \right)^{1/n}, \quad (n = 1, 2, \dots).$$

Theorem I holds even if C_0 is replaced by C .

THEOREM II. *Let (1) hold. In order that $\{y_n\}$ should belong to Γ^* whenever $\{x_p\}$ belongs to C , it is necessary and sufficient that*

(II, 1) *the sequence $\{\theta_n\}$ is bounded, where $\theta_n, (n = 1, 2, \dots)$, are given by (3).*

THEOREM III. *Let (1) hold. In order that $\{y_n\}$ should belong to C whenever $\{x_p\}$ belongs to Γ , it is necessary and sufficient that*

(III, 1) $|a_{np}|^{1/p} \leq M$ *independently of n, p ;*

(III, 2) $\lim_{n \rightarrow \infty} a_{np} = a_p$ *exists for each fixed p .*

THEOREM IV. *Let (2) hold. In order that $\{t_n\}$ should belong to C whenever $\{s_p\}$ belongs to Γ^* , it is necessary and sufficient that*

(IV, 1) *the sequence $\{f_n(z)\}$ of integral functions*

$$(4) \quad f_n(z) = \sum_{p=1}^{\infty} a_{np} z^p, \quad (n = 1, 2, \dots),$$

is uniformly bounded on every compact set (of the complex plane);

(IV, 2) = (III, 2) $\lim_{n \rightarrow \infty} a_{np} = a_p$ *exists for each fixed p .*

THEOREM V. *Let (1) hold. In order that $\{y_n\}$ should belong to Γ^* whenever $\{x_p\}$ belongs to Γ , it is necessary and sufficient that*

(V, 1) $|a_{np}|^{1/(n+p)} \leq M$ *independently of n, p .*

THEOREM VI. *Let (2) hold. In order that $\{t_n\}$ should belong to Γ whenever $\{s_p\}$ belongs to Γ^* , it is necessary and sufficient that*

(VI, 1) $|f_n(z)|^{1/n} \rightarrow 0$, *as $n \rightarrow \infty$, uniformly on every compact set (of the complex plane), where $\{f_n(z)\}$ is the sequence of integral functions $f_n(z)$ given by (4).*

THEOREM VII. *Let (1) hold with $a_{ij} = 0$ for $i > j$. In order that $\{y_n\}$ should belong to Γ whenever $\{x_p\}$ belongs to Γ , it is necessary and sufficient that*

(VII, 1) $|a_{np}|^{1/p} \leq M$ *independently of n, p .*

The matrix transformation of Γ^* into Γ^* was studied by Heller [6].

The sufficiency, in each case is a straightforward calculation. The necessity of any of the above conditions is proved by taking special sequences, and constructing sequences to contradict the given condition, or by using Functional Analysis. Indeed, to prove the necessity of (III, 1), let $U_n(x) = y_n = \sum_{p=1}^{\infty} a_{np} x_p, (n = 1, 2, \dots)$, for each fixed $x = \{x_p\} \in \Gamma$. Then $\{U_n(x)\}$ represents a sequence of continuous linear functionals on Γ ([4], Th. 4). Here $\{|a_{np}|^{1/p}\}$ is bounded for each fixed n . Since $\{y_n\} \in C$, it follows that $\overline{\lim}_{n \rightarrow \infty} |U_n(x)| < \infty$ for each fixed $x \in \Gamma$. Define for each $x \in \Gamma, |x| =$ upper bound $(|x_p|^{1/p}, p \geq 1)$.

Then for $x, x' \in \Gamma$, $|x - x'|$ defines a metric or distance in Γ . With the metric, Γ is a complete metric space. Therefore, by Theorem 11 of ([1], p.19), there is a closed sphere S and a fixed number M such that

$$(5) \quad |U_n(x)| \leq M \text{ for } x \in S \text{ and all } n \geq 1.$$

Take the sphere S as $|x| \leq d$. Set $x_p = (d/2)^p$ and $x_j = 0$ for all $j \neq p$ so that $|x| \leq d/2$ and hence $x = \{x_p\} \in S \subset \Gamma$. Then, by (5), it at once follows that

$$|U_n(x)| = |a_{np}(d/2)^p| \leq M.$$

That is, $|a_{np}|^{1/p} \leq M^{1/p}(2/d) < 2m(M)/d$ where $m(M) = \max(1, M)$. This proves the necessity of (III, 1).

A similar proof applies to condition (VII, 1).

Finally, I thank Professor V. Ganapathy Iyer for his help and guidance. I also thank the referee for drawing my attention to the papers of Sheffer [10] and Zeller [14], and other useful comments. Conditions of Theorems V, VI and VII neither include nor are included in Sheffer's conditions. However, Γ and Γ^* are included in the spaces considered by Sheffer. Sheffer [10] and Zeller [14] also dealt with the spaces of all power series with a certain minimal radius of convergence.

BIBLIOGRAPHY

1. S. Banach, *Théorie des opérations linéaires*, Monogr. Mat., Warsaw, (1932).
2. R. G. Cooke, *Infinite matrices and sequence spaces*, Macmillan, London, 1950.
3. P. Dienes, *The Taylor series*, Oxford, 1931.
4. V. Ganapathy Iyer, *On the space of integral functions—I*, J. Indian Math. Soc. (2) **12** (1948), 13-30.
5. G. H. Hardy, *Divergent series*, Oxford, 1949.
6. I. Heller, *Contributions to the theory of divergent series*, Pacific J. Math. **2** (1952), 153-177.
7. T. Kojima, *On generalized Toeplitz's theorems on limit and their applications*, Tôhoku Math. J. **12** (1917), 291-326.
8. K. Knopp, und G. G. Lorentz, *Beiträge zur absoluten Limitierung*, Arch. Math. **2** (1949), 10-16.
9. I. Schur, *Über lineare Transformationen in der theorie der unendlichen Reihen*, J. Reine. Angew. Math. **151** (1920), 79-111.
10. I. M. Sheffer, *Systems of linear equations of analytic type*, Duke Math. J. **11** (1944), 167-180.
11. O. Szász, *Introduction to the theory of divergent series*, University of Cincinnati, Cincinnati, Ohio, 1946.
12. O. Toeplitz, *Über allgemeine lineare Mittelbildungen*, Prace Math. Fiz. **22** (1911) 113-119.
13. A. Wilansky, *Functional analysis*, Blaisdell, New York, 1964.
14. K. Zeller, *Transformationen des Durchschnitts und der Vereinigung von Folgen-*

raumen, Math. Nachr. **10** (1953), 175-177.

15. ———, *Matrix transformationen von Folgenraumen*, Uni. Roma. Ist. Naz. Alta. Mat. Rend. Mat. Appl. (5) **12** (1954), 340-346.

16. ———, *Theorie der Limitierungsverfahren*, Springer-Verlag, Berlin-Göttingen-Heidelberg, 1958.

Received January 8, 1968, and in revised form March 3, 1969.

ANNAMALAI UNIVERSITY
ANNAMALAINAGAR, INDIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

BASIL GORDON
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

James Burton Ax, <i>Injective endomorphisms of varieties and schemes</i>	1
Richard Hindman Bouldin, <i>A generalization of the Weinstein-Aronszajn formula</i>	9
John Martin Chadam, <i>The asymptotic behavior of the Klein-Gordon equation with external potential. II</i>	19
Rina Hadass, <i>On the zeros of the solutions of the differential equation $y^{(n)}(z) + p(z) = 0$</i>	33
John Sollion Hsia, <i>Integral equivalence of vectors over local modular lattices. II</i>	47
Robert Hughes, <i>Boundary behavior of random valued heat polynomial expansions</i>	61
Surender Kumar Jain, Saad H. Mohamed and Surjeet Singh, <i>Rings in which every right ideal is quasi-injective</i>	73
T. Kawata, <i>On the inversion formula for the characteristic function</i>	81
Erwin Kleinfeld, <i>On right alternative rings without proper right ideals</i>	87
Robert Leroy Kruse and David Thomas Price, <i>On the subring structure of finite nilpotent rings</i>	103
Marvin David Marcus and Stephen J. Pierce, <i>Symmetric positive definite multilinear functionals with a given automorphism</i>	119
William Schumacher Massey, <i>Pontryagin squares in the Thom space of a bundle</i>	133
William Schumacher Massey, <i>Proof of a conjecture of Whitney</i>	143
John William Neuberger, <i>Existence of a spectrum for nonlinear transformations</i>	157
Stephen E. Newman, <i>Measure algebras on idempotent semigroups</i>	161
K. Chandrasekhara Rao, <i>Matrix transformations of some sequence spaces</i>	171
Robert Bruce Schneider, <i>Some theorems in Fourier analysis on symmetric sets</i>	175
Ulrich F. K. Schoenwaelder, <i>Centralizers of abelian, normal subgroups of hypercyclic groups</i>	197
Jerrold Norman Siegel, <i>G-spaces, H-spaces and W-spaces</i>	209
Robert Irving Soare, <i>Cohesive sets and recursively enumerable Dedekind cuts</i>	215
Kwok-Wai Tam, <i>Isometries of certain function spaces</i>	233
Awadhesh Kumar Tiwary, <i>Injective hulls of semi-simple modules over regular rings</i>	247
Eldon Jon Vought, <i>Concerning continua not separated by any nonaposyndetic subcontinuum</i>	257
Robert Breckenridge Warfield, Jr., <i>Decompositions of injective modules</i>	263