

Pacific Journal of Mathematics

INJECTIVE HULLS OF SEMI-SIMPLE MODULES OVER REGULAR RINGS

AWADHESH KUMAR TIWARY

INJECTIVE HULLS OF SEMI-SIMPLE MODULES OVER REGULAR RINGS

A. K. TIWARY

The object of this paper is to provide an explicit construction of the injective hull of a semi-simple module over a commutative regular ring.

The existence of injective hulls of an arbitrary module M and their uniqueness upto isomorphism over M was shown by B. Eckmann and A. Schopf in 1953 [6]. But only in few cases these hulls have been described explicitly [1, 2].

In the special case when the ring is regular as well as Noetherian, the problem is already solved since over such a ring every module is known to be semi-simple [9] and hence is its own injective hull [11, 10]. To begin with we show that every monotypic component of the module is injective and then prove a topological lemma about T_1 -spaces. The Zariski topology of the maximal ideal space of the basic ring being T_1 , we make use of the lemma to obtain the desired construction of an injective hull of the module. We show by an example that a semi-simple module over a regular ring need not always be injective and obtain finally a necessary and sufficient condition for the injectivity of the module.

DEFINITION 1. A ring R is called (von Neumann) *regular* if for every $a \in R$, there exists an element $x \in R$ such that $axa = a$. This condition reduces to $a^2x = a$ if R is commutative. A Boolean ring is an example of a commutative regular ring. It is well known that a commutative ring R with unit is regular if and only if every simple R -module is injective [11].

Throughout this paper we shall consider R to be a commutative regular ring with unit 1. Let Ω denote the set of maximal ideals of R . For each $a \in R$ define Ω_a by $\Omega_a = \{P \in \Omega \mid a \notin P\}$. It follows that $\Omega_a \cap \Omega_b = \Omega_{ab}$. Thus Ω can be made into a topological space with $\{\Omega_a \mid a \in R\}$ as the system of basic open sets. This topology of Ω is known as the Zariski topology. Ω is clearly a T_1 -space since if P and Q are any two distinct points in Ω , there exists $a \in P - Q$ which implies that Ω_a is a neighbourhood of Q not containing P .

DEFINITION 2. Let M be a semi-simple R -module. For any simple submodule S of M , there exists exactly one $P \in \Omega$ with $S \cong R/P$. The

sum of all those simple submodules of M which are isomorphic to R/P , will be denoted by M_P and will be called the R/P -monotypic componet of M . The support of M , to be denoted by $\text{Supp } (M)$ is the set of all those maximal ideals P in Ω for which M_P is nonzero.

In our discussion M will always denote a semi-simple R -module with $\text{supp } (M) = S$. As usual for any function f , the symbol $\text{supp } (f)$ will mean the set of all those elements in domain (f) for which $f(x) \neq 0$. We shall write $E = H(M)$ to express the fact that E is an injective hull of M . Where no ambiguity can arise, we let $H(M)$ stand for an arbitrary injective hull of M . If α is any cardinal number and L any module, the sombol $\alpha \odot L$ will stand for the external sum of α copies of L .

THEOREM 1. *For any $P \in S$, the associated monotypic component M_P is an injective module.*

Proof. Let α be the length of M_P and T a set with $|T| = \alpha$. Then $M_P \cong \alpha \odot R/P = E$. Let π be the set of all functions from T into R/P . Now each factor R/P of π being injective [11], π is injective; hence there exists an $H(E) \subseteq \pi$. Without loss of generality we can take α to be an infinite cardinal. Assume E is not injective. Then $E \subset H(E) \subseteq \pi$. Take any element $f \in H(E) - E$. Since $H(E)$ is an essential extension of E , one has $Rf \cap E \neq 0$ which implies $0 \neq rf \in E$ for some $r \in R - P$. As R/P is a field and $f(t) \neq 0$ for infinitely many $t \in T$, we have $0 \neq (r + P)f(t) = rf(t)$ for infinitely many $t \in T$. But this contradicts the fact that $rf \in E$. Hence E is injective.

REMARK 1. $\prod_{P \in S} M_P$ is injective since each factor M_P is injective.

DEFINITION 3. Let X be any topological space and A any subset of X . An element $x \in A$ is called an *isolated point* of A if there exists a neighbourhood U of x such that $U \cap A = \{x\}$, i.e., if $\{x\}$ is an open set in the relative topology of A . A subset A of X is said to be *discrete* if every element x in A is an isolated point of A .

LEMMA 1. *Let $f \in \prod_{P \in S} M_P$ and $a \in R$ such that $0 \neq af \in \bigoplus_{P \in S} M_P$, then every element in $\text{supp } (af)$ is an isolated point of $\text{supp } (f)$.*

Proof. Let $\text{supp } (af) = \{P_1, P_2, \dots, P_n\}$ where $P_i \neq P_j$ if $i \neq j$. This implies that there exist elements $a_i \in P_i - P_1$ ($i = 2, 3, \dots, n$). Put $b = aa_2a_3 \dots a_n$. Then $b \notin P_1$ and $b \in P$ for each $P \in \text{supp } (f)$ with $P \neq P_1$. Hence $\Omega_b \cap \text{supp } (f) = \{P_1\}$ showing that P_1 is an isolated

point of $\text{supp}(f)$. Similar argument will prove that P_2, \dots, P_n are also isolated points of $\text{supp}(f)$.

REMARK 2. It follows from the lemma that the support of any nonzero element in an essential extension of $\bigoplus_{P \in S} M_P$ contains an isolated point.

LEMMA 2. *Let E be a proper essential extension of $\bigoplus_{P \in S} M_P$. Then for any $f \in E - \bigoplus_{P \in S} M_P$, $\text{supp}(f)$ contains infinitely many isolated points.*

Proof. Since E is an essential extension of $\bigoplus_{P \in S} M_P$ and $0 \neq f \in E$, we can find an element $a \in R$ such that $0 \neq af \in \bigoplus_{P \in S} M_P$. Let $\text{supp}(af) = \{P_1, P_2, \dots, P_n\}$. By Lemma 1, each P_i is an isolated point of $\text{supp}(f)$. Choose an element $Q \in \text{supp}(f) - \text{supp}(af)$. As $P_i \not\subseteq Q$, there exist elements $r_i \in P_i - Q$ ($i = 1, 2, \dots, n$). Then

$$r = r_1 r_2 \dots r_n \in (P_1 \cap P_2 \cap \dots \cap P_n) - Q.$$

It follows that $0 \neq rf \in E$. Since for some $s \in R$, $0 \neq srf \in \bigoplus_{P \in S} M_P$, we can apply Lemma 1 to show that the elements in $\text{supp}(srf)$ are isolated points of $\text{supp}(f)$ and they are all distinct from P_1, P_2, \dots, P_n . Now $\text{supp}(f)$ being infinite, we can find an element in

$$\text{supp}(f) - (\text{supp}(af) \cup \text{supp}(srf))$$

which will give rise to another set of finitely many elements isolated points of $\text{supp}(f)$ each being different from the ones obtained before. Proceeding thus we get infinitely many isolated points of $\text{supp}(f)$. This proves the lemma.

We now prove the following topological fact about T_1 -spaces:

LEMMA 3. *In any T_1 -space X , if A and B are nonvoid subsets such that A as well as every nonvoid subset of B has an isolated point, then there exists an isolated point in $A \cup B$.*

Proof. Let the complement of a subset C of X be denoted by C' . Since A is given to have an isolated point p , there exists an open neighbourhood U of p such that $U \cap A = \{p\}$. From

$$U \cap (A \cup (B \cap U')) = U \cap A$$

we conclude that p is also an isolated point of $A \cup (B \cap U')$. If $B \cap U$ is empty, then p is an isolated point of $A \cup B$ and so the lemma holds. We have therefore to consider only the case when $B \cap U$ is nonvoid.

By hypothesis $B \cap U$ contains an isolated point q which can be assumed to be distinct from p without any loss in generality. This assumption, together with the fact that X is T_1 implies that $\{p\}'$ is an open set containing q . Now q being an isolated point of $B \cap U$, we have $V \cap B \cap U = \{q\}$ for some neighbourhood V of q . Thus we obtain

$$U \cap V \cap \{p\}' \cap (A \cup B) = U \cap V \cap \{p\}' \cap B = \{q\} \cap \{p\}' = \{q\}.$$

Since $U \cap V \cap \{p\}'$ is a neighbourhood of q , the above relation implies that q is an isolated point of $A \cup B$.

REMARK 3. From Lemma 3 we immediately have the following

(i) Let B be a discrete subset of a T_1 -space X and A any subset of X with an isolated point, then $A \cup B$ has an isolated point.

(ii) If A and B are nonvoid subsets of a T_1 -space X with the property that each of their nonvoid subsets has an isolated point then $A \cup B$ has the same property.

LEMMA 4. Let $A = \bigcup_{i \in I} A_i$ where each A_i is without an isolated point. Then A has no isolated point.

Proof. Suppose A has an isolated point p . Then $p \in A_i$ for some $i \in I$ and $\{p\} = U \cap A$ for some neighbourhood U of p . Hence $\{p\} = U \cap A_i$ contrary to the hypothesis that A_i is without an isolated point. Thus A has no isolated point.

LEMMA 5. If A has no isolated point, then \bar{A} , the closure of A also has no isolated point.

Proof. Assume p is an isolated point in \bar{A} with $V \cap \bar{A} = \{p\}$ for some neighbourhood V of p , then $p \in \bar{A} \cap A'$ implies the existence of an element $q \in V \cap A \subseteq V \cap \bar{A}$ with q distinct from p , a contradiction. Hence A has no isolated point.

REMARK 4. We know that the semi-simple module $M = \sum_{P \in S} M_P$ (direct) hence $M \cong \bigoplus_{P \in S} M_P$. Since the injective module $\prod_{P \in S} M_P$ contains $\bigoplus_{P \in S} M_P$ as a submodule, it also contains an $H(\bigoplus_{P \in S} M_P)$. Thus to find an injective hull of M , it is sufficient to obtain one of $\bigoplus_{P \in S} M_P$ inside $\prod_{P \in S} M_P$. This is done in the following:

THEOREM 2. Let $H = \{f \in \prod_{P \in S} M_P \mid \text{Every nonvoid subset of } \text{supp}(f) \text{ has an isolated point}\}$. Then H is an injective hull of $\bigoplus_{P \in S} M_P$.

Proof. Let f, g be any two elements in H , then since

$\text{supp}(f + g) \subseteq \text{Supp}(f) \cup \text{supp}(g)$, we have $f + g \in H$ by Remark 3 (ii) following Lemma 3. Now if $a \in R, f \in H$, then $\text{supp}(af) = \Omega_a \cap \text{supp}(f)$ implies that $af \in H$. Hence H is an R -submodule of $\prod_{P \in S} M_P$ and it contains $\bigoplus_{P \in S} M_P$ since every nonvoid subset of a finite set is discrete. Now let $0 \neq f \in H$, then $\text{supp}(f)$ is nonempty and hence contains an isolated point P so that for some

$$a \in R, \text{supp}(af) = \Omega_a \cap \text{supp}(f) = \{P\}.$$

Thus $0 \neq af \in \bigoplus_{P \in S} M_P$. Hence H is an essential extension of $\bigoplus_{P \in S} M_P$.

As to the injectivity of H assume by way of contradiction that H has a proper essential extension E . Then $H \subset E \subseteq \prod_{P \in S} M_P$. Take $f \in E, f \notin H$. Then there exists a nonvoid subset of $\text{supp}(f)$ without isolated points. Denote by X , the union of all those subsets of $\text{supp}(f)$ which have no isolated points. By Lemma 4, X has no isolated point. Let $Y = \text{supp}(f) \cap X'$ where X' is the complement of X in S . Then Y is nonvoid since by Remark 2, Lemma 1, $\text{supp}(f)$ contains an isolated point which cannot belong to X . Thus $\text{supp}(f) = X \cup Y$ is a decomposition of $\text{supp}(f)$ into disjoint nonempty subsets X and Y . Moreover every nonvoid subset of Y contains an isolated point for otherwise it will have to be contained in X which is not possible. Now for any subset $A \subseteq \text{supp}(f)$, define f_A to be the function such that

$$f_A(P) = \begin{cases} f(P) & \text{if } P \in A \\ 0 & \text{if } P \in S - A \end{cases}$$

we can then write $f = f_X + f_Y$. Since $\text{supp}(f_Y) = Y$, one has $f_Y \in H$ and hence from $f_X = f - f_Y$, it follows that $f_X \in E$. The fact that f_X is a nonzero element in an essential extension E of $\bigoplus_{P \in S} M_P$, then implies that $X = \text{supp}(f_X)$ has an isolated point. We thus arrive at a contradiction. Hence H is injective. This completes the proof.

COROLLARY 1. $\prod_{P \in S} M_P$ is an injective hull of $\bigoplus_{P \in S} M_P$ if and only if every nonvoid subset of S has an isolated point. In particular if S is discrete in Ω , then $\prod_{P \in S} M_P \cong H(M)$.

Proof. If S has the property that each of its nonvoid subsets has an isolated point, then for every $f \in \prod_{P \in S} M_P$, $\text{supp}(f)$ has the same property. Hence by Theorem 2, $\prod_{P \in S} M_P = H(\bigoplus_{P \in S} M_P)$. On the other hand let $\prod_{P \in S} M_P = H(\bigoplus_{P \in S} M_P)$. Suppose that some non-empty subset A of S has no isolated point. Then A must be an infinite set. We can find a function $f \in \prod_{P \in S} M_P$ with $\text{supp}(f) = A$. Then $f \notin \bigoplus_{P \in S} M_P$ and hence $f \neq 0$. Since $\prod_{P \in S} M_P$ is an essential extension of $\bigoplus_{P \in S} M_P$, by

Remark 2, $\text{supp}(f)$ has an isolated point contrary to the assumption that A has no isolated point. Hence every nonvoid subset of S has an isolated point. The last part of the corollary follows immediately from the fact that every element in a discrete set is an isolated point.

COROLLARY 2. *If S contains only principal ideals, then*

$$\prod_{P \in S} M_P = H(\bigoplus_{P \in S} M_P).$$

Proof. Let Ra be any maximal ideal in S . If P in S is different from Ra , then $a \notin P$ since $a \in P$ would mean $Ra \subseteq P$, hence $Ra = P$, a contradiction. Regularity of R implies that $a = a^2x$ for some $x \in R$. Since $0 = a(1 - ax)$ belongs to every P in S , $1 - ax$ belongs to every element in S different from Ra . Also $1 - ax \notin Ra$ since otherwise $1 \in Ra$. It follows that $\Omega_{1-ax} \cap S = \{Ra\}$. Thus every element in S is an isolated point. By Corollary 1, we have $\prod_{P \in S} M_P = H(\bigoplus_{P \in S} M_P)$.

REMARK 5. For any module M over a regular and Noetherian ring R , $\prod_{P \in S} M_P = H(\bigoplus_{P \in S} M_P) = \bigoplus_{P \in S} M_P$ since every ideal of R is a principal ideal [9] and every R -module is injective [10, 11].

COROLLARY 3. *There exist semi-simple modules over a regular ring which are not injective.*

Proof. Let R_0 be the two-element Boolean ring $\{0, e_0\}$, I an infinite index set and R , the set of all functions $f: I \rightarrow R_0$. Then R is a complete Boolean ring and hence a commutative regular ring. For each $\alpha \in I$, define P_α by $P_\alpha = \{f \in R \mid f(\alpha) = 0\}$. It is easily seen that P_α is a maximal ideal of R [7]. Let $M = \bigoplus_{\alpha \in I} R/P_\alpha$. Then M is a semi-simple module with $\text{Supp}(M) = \{P_\alpha \mid \alpha \in I\}$. Take any $P_{\alpha_0} \in \text{Supp}(M)$ and define f by

$$f(\alpha) = \begin{cases} e_0 & \text{if } \alpha = \alpha_0 \\ 0 & \text{if } \alpha \neq \alpha_0 \end{cases}$$

then $f \in R - P_{\alpha_0}$ and $f \in P_\beta$ for all $\beta \in I$ with $\beta \neq \alpha_0$. Thus

$$\Omega_f \cap \text{Supp}(M) = \{P_{\alpha_0}\}$$

which implies that $\text{Supp}(M)$ is discrete. Hence by Corollary 1, $\prod_{\alpha \in I} (R/P_\alpha) = H(\bigoplus_{\alpha \in I} (R/P_\alpha))$. The fact that I is infinite then shows that $\bigoplus_{\alpha \in I} (R/P_\alpha)$ is not injective.

COROLLARY 4. *If $S = A \cup D_1 \cup D_2 \cup \dots \cup D_n$ where A has an*

isolated point and $D_i (i = 1, 2, \dots, n)$ are discrete sets, then $\prod_{P \in S} M_P \cong H(M)$.

Proof. It follows immediately from Lemma 3 and Corollary 1.

In Corollary 3 we have a concrete example showing that not every semi-simple R -module is injective. It is therefore worthwhile to ask under what conditions a semi-simple R -module is injective. The following theorem gives a characterisation for the injectivity of a semi-simple module.

THEOREM 3. *M is injective if and only if S has only finite discrete subsets.*

Proof. Let M be injective. Assume that $D \subseteq S$ is an infinite discrete subset. We can find $f \in \prod_{P \in S} M_P$ with $\text{supp}(f) = D$. Since D is infinite, $f \notin \bigoplus_{P \in S} M_P$. The fact that $\text{supp}(f)$ is discrete implies by Theorem 2, that $f \in H(\bigoplus_{P \in S} M_P) = \bigoplus_{P \in S} M_P$ and so we get a contradiction. Hence S contains only finite discrete subsets.

Conversely suppose that S has only finite discrete subsets. Assume that M is not injective. Then $\bigoplus_{P \in S} M_P$ has a proper essential extension E inside $\prod_{P \in S} M_P$. Hence for any $f \in E - \bigoplus_{P \in S} M_P$, $\text{supp}(f)$ contains an infinite discrete subset by Lemma 2. This contradiction then proves that M is injective.

Added in Proof.

REMARK 6. Under the assumptions of Theorem 3, S is a compact subset of Ω .

Proof. Let $S \subseteq \bigcup_{i \in I} \Omega_{a_i}$ so that $S = \bigcup_{i \in I} (S \cap \Omega_{a_i})$ where we assume without loss of generality that each $S \cap \Omega_{a_i}$ is nonvoid. For each i in I , pick one P_i from $S \cap \Omega_{a_i}$ and let A be the set of all such P_i . Then $\Omega_{a_i} \cap A = \{P_i\}$ for each i in I . This implies that A is a discrete subset of S and hence by Theorem 3, A is finite. Consequently S is compact.

As a consequence of the above remark, we obtain as a corollary of Theorem 3, the following result of J. Levine, announced in an abstract in the Notices:

COROLLARY. (Levine) *If an injective module M over a commutative regular ring R is a direct sum of simple submodules, then there are only finitely many nonisomorphic simples in the sum.*

Proof. Let $M^* = \sum_P X_P$ be the sum of nonisomorphic simple submodules in the direct sum decomposition of M . Then for each X_P ,

there exists exactly one P in S with X_P isomorphic to R/P and hence the R/P -monotypic component of M^* is X_P . Moreover, M^* being a direct summand of M , is injective and, therefore, by Remark 6, its support S^* is compact. Any nonvoid subset of S^* also has this property since it is injective. We propose to show that S^* is discrete. Take any P in S^* and let $\{P\}'$ be the complement of $\{P\}$ in S^* . Then $\{P\}'$ being open and compact, we have $\{P\}' = U_{i \in 1}^n S_{c_i}$, where $S_{c_i} = \Omega_{c_i} \cap S^*$. Now, c_i in R implies that there exists x_i in R with $c_i = c_i^2 x_i$, $i = 1, 2, \dots, n$. Put $d_i = 1 - c_i x_i$. Then from $c_i d_i = 0$, it follows that $d = d_1 d_2 \dots d_n$ belongs to every Q in S^* , different from P and does not belong to P . Hence $\{P\} = S_d$. Thus every point in S^* is an isolated point as was required. By Theorem 3 we have S^* finite.

REMARK 7. Theorem 1 is a special case of a more general Proposition of C. Faith [Proposition 3, Rings with ascending condition on annihilators, Nagoya Math. J. 27 (1966), 179–181]. Let a module M be called Σ -injective if it is injective and every direct sum of copies of M is also injective. Then Proposition 3 of Faith has the following corollaries:

COROLLARY 1. *Let R be any ring, and let M be any injective simple module. Then if M is finite dimensional over the field $K = \text{End } M_R$, then M is Σ -injective.*

COROLLARY 2. *If R is any commutative ring, and M is an injective simple module, then M is Σ -injective.*

Theorem 1 is a special case of Corollary 2 when R is a regular ring.

REMARK 8. Corollary 3 of Theorem 2 provides an example of a semisimple module over a commutative regular ring which is not injective. C. Faith has sketched an example of a simple module over a noncommutative regular ring which is not injective [Chapter 15, "Lectures on Injective Modules and Quotient Rings" Springer Verlag, New York 1967].

I should like to express my grateful thanks to the referee for suggesting the addition of Remarks 6, 7 and 8 in proof.

This paper is a part of a doctoral dissertation submitted to McMaster University in 1966. I should like to express my indebtedness and grateful thanks to Professor B. Banaschewski under whose guidance this work was done.

REFERENCES

1. B. Banaschewski, *On the injective hulls of cyclic modules over Dedekind domains*, Canad. Math. Bull. **9** (1966), 183-186.
2. ———, *On coverings of modules*, Math. Nachr. **31** (1966), 57-71.
3. N. Bourbaki, *Algebra*, Chap II, Herman, Paris, 1962.
4. H. Cartan and S. Eilenberg, *Homological algebra*, Princeton, 1956.
5. C. Chevalley, *Fundamental concepts of algebra*, Academic Press, 1956.
6. B. Eckmann and A. Schopf, *Über injektive Moduln*, Archiv der Math. **4** (1953), 75-78.
7. L. Gillman and M. Jerison, *Rings of continuous functions*, Van Nostrand Company 1960.
8. J.L. Kelley, *General topology*, Van Nostrand Company, 1955.
9. J. Lambek, *Lectures on rings and modules*, Blaisdell Publishing Company, 1966.
10. E. Matlis, *Injective modules over Noetherian rings*, Pacific. J. Math. **8** (1958), 511-528.
11. A. Rosenberg and D. Zelinski, *Finiteness of the injective hull*, Math. Z. **70** (1959), 373-380.

Received November 28, 1968.

BANARAS HINDU UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

BASIL GORDON
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 31, No. 1

November, 1969

James Burton Ax, <i>Injective endomorphisms of varieties and schemes</i>	1
Richard Hindman Bouldin, <i>A generalization of the Weinstein-Aronszajn formula</i>	9
John Martin Chadam, <i>The asymptotic behavior of the Klein-Gordon equation with external potential. II</i>	19
Rina Hadass, <i>On the zeros of the solutions of the differential equation $y^{(n)}(z) + p(z) = 0$</i>	33
John Sollion Hsia, <i>Integral equivalence of vectors over local modular lattices. II</i>	47
Robert Hughes, <i>Boundary behavior of random valued heat polynomial expansions</i>	61
Surender Kumar Jain, Saad H. Mohamed and Surjeet Singh, <i>Rings in which every right ideal is quasi-injective</i>	73
T. Kawata, <i>On the inversion formula for the characteristic function</i>	81
Erwin Kleinfeld, <i>On right alternative rings without proper right ideals</i>	87
Robert Leroy Kruse and David Thomas Price, <i>On the subring structure of finite nilpotent rings</i>	103
Marvin David Marcus and Stephen J. Pierce, <i>Symmetric positive definite multilinear functionals with a given automorphism</i>	119
William Schumacher Massey, <i>Pontryagin squares in the Thom space of a bundle</i>	133
William Schumacher Massey, <i>Proof of a conjecture of Whitney</i>	143
John William Neuberger, <i>Existence of a spectrum for nonlinear transformations</i>	157
Stephen E. Newman, <i>Measure algebras on idempotent semigroups</i>	161
K. Chandrasekhara Rao, <i>Matrix transformations of some sequence spaces</i>	171
Robert Bruce Schneider, <i>Some theorems in Fourier analysis on symmetric sets</i>	175
Ulrich F. K. Schoenwaelder, <i>Centralizers of abelian, normal subgroups of hypercyclic groups</i>	197
Jerrold Norman Siegel, <i>G-spaces, H-spaces and W-spaces</i>	209
Robert Irving Soare, <i>Cohesive sets and recursively enumerable Dedekind cuts</i>	215
Kwok-Wai Tam, <i>Isometries of certain function spaces</i>	233
Awadhesh Kumar Tiwary, <i>Injective hulls of semi-simple modules over regular rings</i>	247
Eldon Jon Vought, <i>Concerning continua not separated by any nonaposyndetic subcontinuum</i>	257
Robert Breckenridge Warfield, Jr., <i>Decompositions of injective modules</i>	263