

Pacific Journal of Mathematics

**ON COMMUTATIVE RINGS OVER WHICH THE SINGULAR
SUBMODULE IS A DIRECT SUMMAND FOR EVERY MODULE**

VASILY CATEFORIS AND FRANCIS LOUIS SANDOMIERSKI

ON COMMUTATIVE RINGS OVER WHICH THE SINGULAR SUBMODULE IS A DIRECT SUMMAND FOR EVERY MODULE

V. C. CATEFORIS AND F. L. SANDOMIERSKI

A commutative ring R with 1 over which the singular submodule is a direct summand for every module, is a semi-hereditary ring with finitely many large ideals. A commutative semi-simple (with d.c.c.) ring is characterized by the property that every semi-simple module is injective.

In this note we continue our investigation of the commutative non-singular rings over which for any module M , the singular submodule $Z(M)$ is a direct summand [1]. As in [1] we say that such a ring has SP . Throughout this paper a ring R is commutative with identity 1 and all modules are unitary. A ring is regular (in the sense of Von Neumann) if every finitely generated right ideal of R is generated by an idempotent; semi-simple means semi-simple with d.c.c.. Notation and terminology here is as in [1].

In [1] we established the following characterization of rings with SP , included here for easy reference:

THEOREM 1. *For a ring R the following are equivalent:*

- (a) R has SP
- (b) R is regular and has BSP [1].
- (c) $Z(R) = (0)$ and for every large ideal I of R , the ring R/I is semi-simple.
- (d) Every R -module M with $Z(M) = M$ is R -injective.

In particular if R has SP , then R is hereditary.

We shall need the following corollaries of this theorem:

COROLLARY 1.1. *Every homomorphic image of a ring R with SP , has SP .*

Proof. Let $S = R/I$ for some ideal $I (\neq (0), R)$ of R ; S is regular since R is and thus $Z(S) = (0)$. A large ideal A of S is of the form J/I where J is a large ideal of R containing I . Thus it follows from (c) that S/A is semi-simple since $S/A \cong R/J$. Now S has SP since it satisfies (c).

COROLLARY 1.2. *Every singular module over a ring R with SP*

is semi-simple.

Proof. Every cyclic singular module is semi-simple by (c).

For any R -module M we denote by $\text{So}_R(M)$ (or $\text{So}(M)$ if no ambiguity arises) the socle of M , that is the module sum of the simple submodules of M .

THEOREM 2. *Over a ring R with SP the socle, $\text{So}(M)$, of every nonzero R -module M is large in M .*

Proof. By definition of essential extension, it suffices to show that every nonzero cyclic module R/I has nonzero socle.

We have:

LEMMA 3. *Let R be a ring with SP and $\{e_n: n \in A\}$ a countable set of orthogonal idempotents of R such that $e_n R$ contains a proper ($\neq e_n R$) large R -submodule I_n , for each $n \in A$. Then, the set $\{e_n: n \in A\}$ is finite.*

Proof. For each $n \in A$ the module $e_n R/I_n = S_n \neq (0)$ is singular. It follows that $\bigoplus_{n \in A} S_n$ is singular and hence injective by Theorem 1 (d). Let $g: \bigoplus_{n \in A} e_n R \rightarrow \bigoplus_{n \in A} S_n$ be the R -homomorphism defined by $g|_{e_n R}: e_n R \rightarrow e_n R/I_n$, the natural R -epimorphism. Since $\bigoplus_{n \in A} e_n R \subseteq R$ and $\bigoplus_{n \in A} S_n$ is injective, there is extension $g^*: R \rightarrow \bigoplus S_n$ of g and since $\text{Im } g^*$ is clearly contained in only a finite number of the S_n , it follows that they are finitely many. Hence $\{e_n: n \in A\}$ is a finite set.

To return to the proof of Theorem 2, we note that Corollary 1.1 and Lemma 3 imply that if R is a ring with SP and $\{e_n: n \in A\}$ is an infinite set of orthogonal idempotents of R/I , then all but finitely many of the ideals $e_n(R/I)$ are semi-simple. If R/I has no infinite set of orthogonal idempotents, then R/I itself is semi-simple.

To establish the characterizations announced at the beginning of this paper we need the well known characterizations (e.g. [2]) of a commutative regular ring contained in Theorem 4 below. For each module M , $J(M)$ denotes the radical of M .

THEOREM 4. *For a commutative ring R the following are equivalent:*

- (a) R is regular.
- (b) Every simple R -module is injective.

(c) $J(M) = (0)$ for every R -module M .

(d) Every ideal of R is the intersection of the maximal ideals that contain it.

We can now complete the characterization of rings with SP , partially established in [1] as Theorem 2.9.

THEOREM 5. *For a ring R the following are equivalent:*

(a) R has SP .

(b) $R/\text{So}(R)$ is a semi-simple ring and R is regular.

(c) R is semi-hereditary and has only a finite number of large ideals.

Proof. (a) implies (b). A consequence of parts (b) and (c) of Theorem 1 and Theorem 2.

(b) implies (c). R is certainly semi-hereditary since it is regular. Furthermore since every large ideal of R contains the socle of R , $R/\text{So}(R)$ is semi-simple implies that R has only a finite number of maximal large ideals. Since every ideal containing a large ideal is itself large, the assertion that R has only a finite number of large ideals follows now from (d) of Theorem 4.

(c) implies (a). Theorem 2.9 [2].

The proof of Theorem 5 is now complete.

We close with the following characterization of commutative semi-simple rings:

THEOREM 6. *For a commutative ring R the following are equivalent:*

(a) R is semi-simple.

(b) Every semi-simple R -module is injective.

Proof. (b) implies (a). From (b) and part (b) of Theorem 4 it follows that R is regular. It is, hence, sufficient to show that R contains no infinite sets of orthogonal idempotents. Thus let $\{e_n: n \in A\}$ be any countable set of orthogonal idempotents. It follows from part (c) of Theorem 4 that each $e_n R$ contains a maximal R -submodule I_n and if we let $S_n = e_n R / I_n$ then $S = \bigoplus_{n \in A} S_n$ is a semi-simple R -module. From (b) S is injective and an argument similar to the one used to prove Lemma 3, gives now that the set $\{e_n: n \in A\}$ is finite. Hence R contains no infinite sets of orthogonal idempotents.

We do not know whether Theorem 6 remains true if R is not assumed commutative. In this direction it can be shown that if (b)

is assumed for right modules then R is finite dimensional on the right (i.e., R contains no infinite direct sum of nonzero right ideals).

REFERENCES

1. V. C. Cateforis, and F. L. Sandomierski, *The singular submodule splits off*, J. of Algebra **10** (1968), 149-165.
2. A. Rosenberg, and D. Zelinski, *Finiteness of the injective hull*, Math. Zeit. **70** (1959), 372-380.

Received February 18, 1969.

UNIVERSITY OF KENTUCKY
LEXINGTON, KENTUCKY

UNIVERSITY OF WISCONSIN
MADISON, WISCONSIN

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

BASIL GORDON
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Efraim Pacillas Armendariz, <i>Quasi-injective modules and stable torsion classes</i>	277
J. Adrian (John) Bondy, <i>On Ulam's conjecture for separable graphs</i>	281
Vasily Cateforis and Francis Louis Sandomierski, <i>On commutative rings over which the singular submodule is a direct summand for every module</i>	289
Rafael Van Severen Chacon, <i>Approximation of transformations with continuous spectrum</i>	293
Raymond Frank Dickman and Alan Zame, <i>Functionally compact spaces</i>	303
Ronald George Douglas and Walter Rudin, <i>Approximation by inner functions</i>	313
John Walter Duke, <i>A note on the similarity of matrix and its conjugate transpose</i>	321
Micheal Neal Dyer and Allan John Sieradski, <i>Coverings of mapping spaces</i>	325
Donald Campbell Dykes, <i>Weakly hypercentral subgroups of finite groups</i>	337
Nancy Dykes, <i>Mappings and realcompact spaces</i>	347
Edmund H. Feller and Richard Laham Gantos, <i>Completely injective semigroups</i>	359
Irving Leonard Glicksberg, <i>Semi-square-summable Fourier-Stieltjes transforms</i>	367
Samuel Irving Goldberg and Kentaro Yano, <i>Integrability of almost cosymplectic structures</i>	373
Seymour Haber and Charles Freeman Osgood, <i>On the sum $\sum \langle n\alpha \rangle^{-1}$ and numerical integration</i>	383
Sav Roman Harasymiv, <i>Dilations of rapidly decreasing functions</i>	395
William Leonard Harkness and R. Shantaram, <i>Convergence of a sequence of transformations of distribution functions</i>	403
Herbert Frederick Kreimer, Jr., <i>A note on the outer Galois theory of rings</i>	417
James Donald Kuelbs, <i>Abstract Wiener spaces and applications to analysis</i>	433
Roland Edwin Larson, <i>Minimal T_0-spaces and minimal T_D-spaces</i>	451
A. Meir and Ambikeshwar Sharma, <i>On Ilyeff's conjecture</i>	459
Isaac Namioka and Robert Ralph Phelps, <i>Tensor products of compact convex sets</i>	469
James L. Rovnyak, <i>On the theory of unbounded Toeplitz operators</i>	481
Benjamin L. Schwartz, <i>Infinite self-interchange graphs</i>	497
George Szeto, <i>On the Brauer splitting theorem</i>	505
Takayuki Tamura, <i>Semigroups satisfying identity $xy = f(x, y)$</i>	513
Kenneth Tolo, <i>Factorizable semigroups</i>	523
Mineko Watanabe, <i>On a boundary property of principal functions</i>	537
James Juei-Chin Yeh, <i>Singularity of Gaussian measures in function spaces with factorable covariance functions</i>	547