

Pacific Journal of Mathematics

ON THE BRAUER SPLITTING THEOREM

GEORGE SZETO

ON THE BRAUER SPLITTING THEOREM

GEORGE SZETO

This paper presents a proof for the Brauer splitting theorem in the context of a commutative ring with no idempotents except 0 and 1 and continues this investigation. The main results in this paper are the Brauer splitting theorem and the classification of all finitely generated projective indecomposable modules over a separable group algebra.

Throughout this paper we assume that the ring R is a commutative ring with no idempotents except 0 and 1, that the group G has order n invertible in R , and that all RG -modules are unitary left RG -modules. We know that the order of G , n , is invertible in R if and only if RG is separable.

1. First, let us recall the following Brauer splitting theorem: Let K be a field and G be a group of order n invertible in K , then $K(\sqrt[m]{1})$ is a splitting field for G , where m is the exponent of G and $\sqrt[m]{1}$ is a primitive m^{th} -root of 1 ([6], Th. 41-1, p. 292 and Corollary 70-24, p. 475). In [8], G. J. Janusz defined a ring R to be a splitting ring for G if the group algebra RG is the direct sum of central separable R -algebras each equivalent to R in the Brauer group of R ; that is, $RG \cong \bigoplus \sum_{i=1}^s \text{Hom}_R(P_i, P_i)$, where $\{P_i\}$ are finitely generated projective faithful R -modules, the number of different conjugate classes in G is equal to s . He then proved the Brauer splitting theorem for a Noetherian regular domain, R . This section gives a proof for the above theorem when R is any commutative ring with no idempotents except 0 and 1.

LEMMA 1. *Let R_0 be a subring of R . If R_0 is a splitting ring for G , then R is a splitting ring for G .*

Proof. Because R_0 is a splitting ring for G , $R_0G \cong \bigoplus \sum_{i=1}^s \text{Hom}_{R_0}(P_i, P_i)$ where $\{P_i\}$ are finitely generated projective faithful R_0 -modules. Then we have

$$\begin{aligned} RG &\cong R \otimes_{R_0} R_0G \cong R \otimes_{R_0} \left(\bigoplus \sum_{i=1}^s \text{Hom}_{R_0}(P_i, P_i) \right) \\ &\cong \bigoplus \sum_{i=1}^s R \otimes_{R_0} \text{Hom}_{R_0}(P_i, P_i) \cong \bigoplus \sum_{i=1}^s \text{Hom}_R(R \otimes_{R_0} P_i, R \otimes_{R_0} P_i), \end{aligned}$$

where $\{R \otimes_{R_0} P_i\}$ are finitely generated projective faithful R -modules. This follows since $\{P_i\}$ are finitely generated projective faithful R_0 -

modules ([1], Proposition 5-5). Thus R is a splitting ring for G .

THEOREM 2. *If R is a commutative ring with no idempotents except 0 and 1 and RG is a separable group algebra, then $R[\sqrt[m]{1}]$ is a splitting ring for G where $\sqrt[m]{1}$ is a primitive m^{th} -root of 1.*

Proof. Let Z be the set of integers, Q be the set of rationals. The proof divides into two cases.

Case 1. The prime ring of R is finite. Let $\text{Char}(R) = p^e$, where p is a prime integer and e is in Z .

$Z/(p^e)$ is a local ring with the maximal ideal $(p)/(p^e)$ which is also nilpotent. For $(Z/(p^e))[\theta]$ where $\theta = \sqrt[m]{1}$, we have

$$\frac{(Z/(p^e))[\theta]}{((p)/(p^e))[\theta]} \cong (Z/(p))(\bar{\theta})$$

where $\bar{\theta}$ is a primitive m^{th} -root of 1 over $Z/(p)$. Now $(Z/(p))(\bar{\theta})$ is a field; so $((p)/(p^e))[\theta]$ is a maximal ideal. On the other hand, since $(p)/(p^e)$ is nilpotent, $((p)/(p^e))[\theta]$ is also nilpotent. But then $((p)/(p^e))[\theta]$ is an unique maximal ideal and a nilpotent ideal of $(Z/(p^e))[\theta]$. Therefore, $(Z/(p^e))[\theta]$ is a complete local ring where the completion is in the sense of m -topology (see [9], p. 254). Then the Brauer group natural map

$$B((Z/(p^e))[\theta]) \longrightarrow B\left(\frac{(Z/(p^e))[\theta]}{((p)/(p^e))[\theta]}\right) \cong B((Z/(p))(\bar{\theta}))$$

is monomorphic ([1], Corollary 6-2). But $(Z/(p))(\bar{\theta})$ is a splitting field for G ; so $(Z/(p^e))[\theta]$ is a splitting ring for G . Thus $R[\theta]$ is a splitting ring for G by the lemma.

Case 2. The prime ring of R is $Z(n)$ which is the quotient ring of Z with respect to the multiplicative closed set $\{n, n^2, \dots\}$. Since $Z(n)[\theta]$ is a Dedekind domain, it is Noetherian and regular. Then the Brauer group natural map $B(Z(n)[\theta]) \rightarrow B(Q(\theta))$ is monomorphic ([1], Th. 7-2). But $Q(\theta)$ is the quotient field of $Z(n)[\theta]$ and a splitting field for G by the Brauer splitting theorem. Therefore, $Z(n)[\theta]$ is a splitting ring for G and so $R[\sqrt[m]{1}]$ is a splitting ring for G by the lemma. By combining Cases 1 and 2, the theorem is proved.

REMARK. The above theorem tells us the existence of a splitting ring, $R[\sqrt[m]{1}]$, for G , if RG is a separable group algebra. We also know that $R[\sqrt[m]{1}]$ is a finitely generated projective and separable R -algebra ([8], Corollary 2-4). But there exists a central separable R -

algebra without a finitely generated projective and separable splitting ring. The following example is due to O. Goldman: Let R be $Z[\sqrt{-2}]$, i, j, k be the usual quaternion basis. If $\alpha = (1 + i)/\sqrt{2}$ and $\beta = (1 + j)/\sqrt{2}$, then $R1 \oplus R\alpha \oplus R\beta \oplus R\alpha\beta$ is central separable over R . But R has no finitely generated projective and separable extension except direct sums of copies of R , and $R1 \oplus R\alpha \oplus R\beta \oplus R\alpha\beta$ cannot be split.

2. In this section, assume RG is a split group algebra,

$$RG \cong \bigoplus_i \text{Hom}_R(P_i, P_i), \quad i = 1, 2, \dots, s.$$

When $\{P_i\}$ are considered as RG -modules ([3], p. 5), the classification of all finitely generated projective indecomposable RG -modules can be obtained. Observe that the order of the group G , n , is invertible in R if and only if RG is separable. Therefore, any RG -module M is finitely generated and projective over RG if and only if M is finitely generated and projective over R (see the proof of Proposition 1-5 in [8]).

Let RG be a separable R -algebra and M be a finitely generated projective RG -module; for any x in M there exist X_1, X_2, \dots, X_q in M and F_1, F_2, \dots, F_q in $\text{Hom}_R(M, R)$ so that $x = \sum_{i=1}^q F_i(x)X_i$. We call $\{F_i, X_i, i = 1, 2, \dots, q\}$ a R -dual basis of M , and $T_M(x) = \sum_{i=1}^q F_i(x)X_i$ the character of M at x in RG ([4], Proposition 3-1). By a group character we mean the restriction of T_M to G . Obviously, a character T_M is completely determined by its restriction to G . In particular, let R be a splitting ring for G ; then

$$RG \cong \bigoplus_{i=1}^s \text{Hom}_R(P_i, P_i) \cong \bigoplus_{i=1}^s (RG)E_i,$$

where E_i is the i^{th} -central primitive idempotent of RG . We let

$$T^i = T_{P_i}.$$

PROPOSITION 3. *If M and N are two isomorphic finitely generated projective RG -modules, then they have the same characters.*

Proof. Let M and N be two isomorphic finitely generated projective RG -modules and let α be the isomorphism. If $\{F_i, X_i, i = 1, 2, \dots, q\}$ is a dual basis of M , then we claim that $\{F_i\alpha^{-1}, \alpha X_i, i = 1, 2, \dots, q\}$ is a dual basis of N . In fact, for any a in N , there exists b in M such that $\alpha(b) = a$; so

$$\begin{aligned} a &= \alpha\left(\sum_{i=1}^q F_i(b)X_i\right) = \sum_i F_i(b)(\alpha X_i) \\ &= \sum_i F_i\alpha^{-1}\alpha(b)(\alpha X_i) = \sum_i ((F_i\alpha^{-1})\alpha(b))(\alpha X_i) \\ &= \sum_i (F_i\alpha^{-1}(a))(\alpha X_i). \end{aligned}$$

This means that $\{F_i\alpha^{-1}, \alpha X_i, i = 1, 2, \dots, q\}$ is a dual basis of N . But the character of any finitely generated projective RG -module is independent of the dual basis chosen; so $T_N(g) = \sum_i F_i\alpha^{-1}(g\alpha X_i) = \sum_i F_i\alpha^{-1}(\alpha g X_i)$, (for α is a RG -isomorphism), and so $= \sum_i F_i(g X_i) = T_M(g)$.

The following proposition will play an important role in our discussion.

PROPOSITION 4. *If N is a finitely generated projective faithful R -module and M a finitely generated projective left $\text{Hom}_R(N, N)$ -module, then $M \cong N \otimes_R N'$ with N' a finitely generated projective R -module.*

Proof. By the Morita Theorem on p. 9 in [3].

REMARK. Proposition 4 gives a counter-example to the converse statement of Proposition 3. Because of Proposition 4, let M and N be two finitely generated projective indecomposable RG -modules over the same central component of the split group algebra RG ; that is, $\text{Hom}_R(P_i, P_i)$, then $M \cong P_i \otimes_R N'$ and $N \cong P_i \otimes_R N''$, where N' and N'' are finitely generated projective indecomposable R -modules. Suppose N' and N'' are in $P(R)$, the class group of R , then

$$T_M(g) = T_{P_i}(g)T_{N'}(1) = T_{P_i}(g) \cdot 1 = T_N(g) .$$

But $P_i \otimes_R N' \cong P_i \otimes_R N''$ only if $N' \cong N''$.

LEMMA 5. *If RG is a split group algebra; that is,*

$$RG \cong \bigoplus_{i=1}^s \text{Hom}_R(P_i, P_i) \cong \bigoplus_{i=1}^s (RG)E_i ,$$

then

$$E_i = \sum_g \frac{k_i T^i(g^{-1})}{n} g ,$$

where g is in G , $k_i = \text{rank}(P_i)$ and $T^i = T_{P_i}$.

Proof. Since

$$RG \cong \bigoplus_{i=1}^s (RG)E_i \cong \bigoplus_{i=1}^s \text{Hom}_R(P_i, P_i), E_i = \sum_g E_i(g)g ,$$

for all g in G , $E_i(g)$ in R . We then have

$$E_i h^{-1} = \sum_g E_i(g)(gh^{-1})$$

for some h in G . Taking the character afforded by RG , we have

$$T_{RG}(E_i h^{-1}) = \sum_g E_i(g) T_{RG}(gh^{-1}) .$$

But $T_{RG}(gh^{-1}) = 0$ in case $gh^{-1} \neq 1$, and $= n$ in case $gh^{-1} = 1$ or $g = h$. Hence $T_{RG}(E_i h^{-1}) = E_i(h)n, E_i(h) = T_{RG}(E_i h^{-1})/n$ (for n is invertible in R).

Next, we find $T_{RG}(E_i h^{-1})$. Because P_i is a finitely generated projective R -module, $\text{Hom}_R(P_i, P_i) \cong P_i \otimes_R \text{Hom}_R(P_i, R)$ ([3], Morita Theorem I). Noting that $\text{rank}(P_i) = \text{rank}(\text{Hom}_R(P_i, R))$, we have

$$T_{(RG)E_i}(g) = T^i(g)k_i \quad \text{for all } i = 1, 2, \dots, s .$$

Therefore,

$$T_{RG}(E_i h^{-1}) = \sum_{j=1}^s T_{(RG)E_j}(E_i h^{-1}) = \sum_j k_j T^j(E_i h^{-1}) .$$

But $T^j(E_i h^{-1}) = 0$ in case $i \neq j$, so

$$T_{RG}(E_i h^{-1}) = k_i T^i(E_i h^{-1}) = k_i T^i(h^{-1}) .$$

Hence,

$$E_i(h) = \frac{T_{RG}(E_i h^{-1})}{n} = \frac{k_i T^i(h^{-1})}{n} .$$

By substituting $E_i(h)$ in E_i , we have

$$E_i = \sum_g E_i(g)g = \sum_g \frac{k_i T^i(g^{-1})}{n} g .$$

This completes the proof.

LEMMA 6. For $i = 1, 2, \dots, s$, $\text{rank}(P_i)$ is neither 0 nor a zero divisor in R .

Proof. First, $\text{rank}(P_i)$ is not 0, otherwise E_i is 0 by Lemma 5. This is impossible.

Next, let $\text{rank}(P_i)$ be k_i , and suppose that k_i is a zero divisor in R . We then have a nonzero element, k' , in R such that $k'k = 0$. But by Lemma 5,

$$E_i = k_i \sum_g \frac{T^i(g^{-1})g}{n} ;$$

so,

$$k'E_i = k'k_i \sum_g \frac{T^i(g^{-1})g}{n} = (k'k_i) \sum_g \frac{T^i(g^{-1})g}{n} = 0 .$$

Noting that $(RG)E_i \cong \text{Hom}_R(P_i, P_i)$, we have

$$k' \text{Hom}_R(P_i, P_i) \cong k'(RG)E_i = k'E_i(RG) = 0 .$$

On the other hand, P_i is a faithful R -module; so $\text{Hom}_R(P_i, P_i)$ is a faithful R -module. Therefore, $k' \text{Hom}_R(P_i, P_i) = 0$ implies $k' = 0$. This is a contradiction. Thus we have proved that k_i is not a zero divisor in R .

THEOREM 7. *Suppose R is a splitting ring for G and all finitely generated projective indecomposable R -modules are of rank 1. Then for any two finitely generated projective indecomposable RG -modules M and N , we have $E_i M \neq 0$ and $E_i N \neq 0$ if and only if $T_M(g) = T_N(g)$ for all g in G , where E_i is the i^{th} -central primitive idempotent of RG .*

Proof. If $E_i M \neq 0$ and $E_i N \neq 0$, then $M \cong E_i M \oplus (1 - E_i)M$ and $N \cong E_i N \oplus (1 - E_i)N$. Since M and N are indecomposable, $(1 - E_i)M = 0$ and $(1 - E_i)N = 0$. We have $N = E_i N$ and $M = E_i M$ as left $\text{Hom}_R(P_i, P_i)$ -modules. Therefore, by Proposition 4, $M \cong P_i \otimes_R N'$ and $N \cong P_i \otimes_R N''$ where N' and N'' are finitely generated projective R -modules. Since M and N are indecomposable RG -modules, N' and N'' are in $P(R)$. Therefore,

$$\begin{aligned} T_M(g) &= T_{P_i \otimes_R N'}(g) = T_{P_i}(g) \cdot 1 \\ &= T_{P_i}(g) T_{N'}(1) = T_N(g) . \end{aligned}$$

Conversely, if $T_M(g) = T_N(g)$ for all g in G , then $T_M(a) = T_N(a)$ for all a in RG . Suppose $E_i M \neq 0$ and $E_i N = 0$ for some i ; then there exists a $j \neq i$ such that $E_j N \neq 0$. Thus M is a $(RG)E_i$ -module and N is a $(RG)E_j$ -module, and so we have

$$T_M(E_i) = T_{P_i}(E_i) = T_{P_i}(1) = \text{rank}(P_i) .$$

By Lemma 6, $\text{rank}(P_i) \neq 0$ in R , so $T_M(E_i) \neq 0$. Obviously, $T_N(E_i) = 0$. Thus $T_M \neq T_N$ on RG . Consequently, $T_M(g) \neq T_N(g)$ for some g in G . This is a contradiction to $T_M(g) = T_N(g)$ for all g in G , and hence the proof is completed.

COROLLARY 8. *If R is a splitting ring for G , and all finitely generated projective indecomposable R -modules are of rank 1; then there are exactly s -classes of finitely generated projective indecomposable RG -modules over different central components each uniquely determined up to an element in $P(R)$.*

Proof. Let M be a finitely generated projective indecomposable

RG -module. From the theorem, we have $M = E_i M \cong P_i \otimes_R N'$ where N' is in $P(R)$. On the other hand, $P_i, i = 1, 2, \dots, s$, is a finitely generated projective indecomposable RG -module over the i^{th} -central component. Therefore, there are exactly s -classes of finitely generated projective indecomposable RG -modules each uniquely determined up to an element in $P(R)$.

From the above result, we have computed the first Grothendieck group of $RG, K^0(RG)$, in the sense of [2], p. 31.

COROLLARY 9. *If R is a splitting ring for G , then*

$$K^0(RG) = (Z \oplus P(R)) \oplus (Z \oplus P(R)) \oplus \dots \oplus (Z \oplus P(R)).$$

A natural question to ask is whether the classification of all finitely generated projective indecomposable RG -modules can be obtained for a nonsplit group algebra. The answer is not known. But for some special rings, we have a definite answer.

For a separable group algebra RG , we have the decomposition, $RG \cong \bigoplus \sum_{i=1}^t A_i$, where A_i has no proper central idempotents and t is an integer.

THEOREM 10. *If R is local or semi-local, then there are exactly t -isomorphic classes of finitely generated projective indecomposable RG -modules.*

Proof. From the decomposition of RG, A_i is a central separable C_i -algebra for each A_i , where C_i is the center of A_i ([1], Th. 2-3). Since R is local or semi-local, C_i is semi-local by the lemma on p. 25 in [5]. Therefore any two finitely generated projective indecomposable RG -modules over the i^{th} -component A_i are in an isomorphic class of finitely generated projective indecomposable RG -modules ([7], Th. 1).

COROLLARY 11. *If R is local or semi-local, then*

$$K^0(RG) = Z \oplus Z \oplus \dots \oplus Z,$$

t -copies of Z .

This paper forms a part of the author's Doctoral Dissertation at Purdue University written under the guidance of Professor F. R. DeMeyer. The author wishes to thank Professor DeMeyer for his guidance, criticism, and encouragement throughout this study.

The author wishes to thank the referee for many helpful suggestions.

BIBLIOGRAPHY

1. M. Auslander and O. Goldman, *The Brauer group of a commutative ring*, Trans. Amer. Math. Soc. **97** (1960), 367-409.
2. H. Bass, *K-theory and stable algebra*, Publ. Math. I.H.E.S. Paris, **22** (1964), 5-60.
3. ———, *The Morita theorems*, Mimeographed Notes, University of Oregon, 1964.
4. H. Cartan and S. Eilenberg, *Homological algebra*, Princeton, 1956.
5. L. N. Childs and F. R. DeMeyer, *On automorphisms of separable algebras*, Pacific J. Math. **23** (1967), 25-34.
6. C. W. Curtis and I. Reiner, *Representation theory of finite groups and associated Algebras*, New York, Interscience, 1962.
7. F. R. DeMeyer, *Projective modules over central separable algebras*, Canad. J. Math. **21** (1969), 39-43.
8. G. J. Janusz, *Separable algebra over commutative rings*, Trans. Amer. Math. Soc. **122** (1966), 461-479.
9. O. Zariski and P. Samuel, *Commutative algebra*, Vol. 2, Princeton, D. Van Nostrand Co., 1960.

Received October 14, 1968.

BRADLEY UNIVERSITY
PEORIA, ILLINOIS

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

BASIL GORDON
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics

Vol. 31, No. 2

December, 1969

Efraim Pacillas Armendariz, <i>Quasi-injective modules and stable torsion classes</i>	277
J. Adrian (John) Bondy, <i>On Ulam's conjecture for separable graphs</i>	281
Vasily Cateforis and Francis Louis Sandomierski, <i>On commutative rings over which the singular submodule is a direct summand for every module</i>	289
Rafael Van Severen Chacon, <i>Approximation of transformations with continuous spectrum</i>	293
Raymond Frank Dickman and Alan Zame, <i>Functionally compact spaces</i>	303
Ronald George Douglas and Walter Rudin, <i>Approximation by inner functions</i>	313
John Walter Duke, <i>A note on the similarity of matrix and its conjugate transpose</i>	321
Micheal Neal Dyer and Allan John Sieradski, <i>Coverings of mapping spaces</i>	325
Donald Campbell Dykes, <i>Weakly hypercentral subgroups of finite groups</i>	337
Nancy Dykes, <i>Mappings and realcompact spaces</i>	347
Edmund H. Feller and Richard Laham Gantos, <i>Completely injective semigroups</i>	359
Irving Leonard Glicksberg, <i>Semi-square-summable Fourier-Stieltjes transforms</i>	367
Samuel Irving Goldberg and Kentaro Yano, <i>Integrability of almost cosymplectic structures</i>	373
Seymour Haber and Charles Freeman Osgood, <i>On the sum $\sum \langle n\alpha \rangle^{-1}$ and numerical integration</i>	383
Sav Roman Harasymiv, <i>Dilations of rapidly decreasing functions</i>	395
William Leonard Harkness and R. Shantaram, <i>Convergence of a sequence of transformations of distribution functions</i>	403
Herbert Frederick Kreimer, Jr., <i>A note on the outer Galois theory of rings</i>	417
James Donald Kuelbs, <i>Abstract Wiener spaces and applications to analysis</i>	433
Roland Edwin Larson, <i>Minimal T_0-spaces and minimal T_D-spaces</i>	451
A. Meir and Ambikeshwar Sharma, <i>On Ilyeff's conjecture</i>	459
Isaac Namioka and Robert Ralph Phelps, <i>Tensor products of compact convex sets</i>	469
James L. Rovnyak, <i>On the theory of unbounded Toeplitz operators</i>	481
Benjamin L. Schwartz, <i>Infinite self-interchange graphs</i>	497
George Szeto, <i>On the Brauer splitting theorem</i>	505
Takayuki Tamura, <i>Semigroups satisfying identity $xy = f(x, y)$</i>	513
Kenneth Tolo, <i>Factorizable semigroups</i>	523
Mineko Watanabe, <i>On a boundary property of principal functions</i>	537
James Juei-Chin Yeh, <i>Singularity of Gaussian measures in function spaces with factorable covariance functions</i>	547