

Pacific Journal of Mathematics

**ON MAJORANTS FOR SOLUTIONS OF ALGEBRAIC
DIFFERENTIAL EQUATIONS IN REGIONS OF THE COMPLEX
PLANE**

STEVEN BARRY BANK

ON MAJORANTS FOR SOLUTIONS OF ALGEBRAIC DIFFERENTIAL EQUATIONS IN REGIONS OF THE COMPLEX PLANE

STEVEN BANK

In this paper, we investigate the rate of growth of functions which are analytic in an arbitrary simplyconnected region of the complex plane and which are solutions of first order algebraic differential equations (i.e., equations of the form $\Omega(z, y, dy/dz) = 0$, where Ω is a polynomial in z, y and dy/dz).

In the course of constructing an example for second order equations, Vijayaraghavan in [5] showed that for *any* realvalued increasing function $\Phi(x)$ on the interval $(0, +\infty)$, it is possible to find a complex function $h(z)$, which is analytic in a simply-connected region R of the plane containing $(0, +\infty)$, and satisfies a first order algebraic differential equation, and which has the property that $|h(x)| > \Phi(x)$ at a sequence of real x tending to $+\infty$ ¹. For a given $\Phi(x)$, the function $h(z)$ constructed was of the form $P(az)$ where $P(u)$ is the Weierstrass P -function with primitive periods w and iw' (w, w' real), and where the constant a was of the form $a = w + ib$, where b depends on Φ and b/w' is irrational. Since $P(az)$ has poles at all points $(mw/a) + (niw'/a)$ where m and n are integers, clearly the region R associated with the solution $h(z) = P(az)$ depends on a and hence on $\Phi(x)$. A natural question is thus raised, namely, can such examples be constructed where, for all $\Phi(x)$, the simply-connected region R remains the same. That is, does there exist a simply-connected region R containing $(0, +\infty)$ with the property that for any real-valued increasing function $\Phi(x)$ on $(0, +\infty)$, there is a solution $h(z)$, analytic on R , of a first order algebraic differential equation, such that $|h(x)| > \Phi(x)$ at a sequence of real x tending to $+\infty$? In this paper we answer this question in the *negative* by proving the following more general result (§2 below): If R is any simply-connected region, then there exists a real-valued continuous function $\Psi(z)$ on R with the property that for any function $h(z)$, analytic on R , which satisfies a first order algebraic differential equation, there is a compact set K contained in R such that $|h(z)| < \Psi(z)$ on $R-K$. In the case where R is not the whole plane, we show that $\Psi(z)$ may be taken to be

¹ None of the solutions $h(z)$ constructed by Vijayaraghavan are real-valued on any interval $(x_0, +\infty)$. Of course this is in accord with the well-known result of Lindelöf [2; p. 213] that a *real-valued* solution on an interval $(x_0, +\infty)$ is majorized, on some interval $(x_1, +\infty)$, by $\exp(\exp x)$.

$\exp(\exp((1 - |f(z)|)^{-1}))$, where f is a univalent analytic mapping of R onto the unit disk (which exists by the Riemann Mapping Theorem). In the case where R is the whole plane, a wellknown result of Valiron [3; p. 41] states that any entire function satisfying a first order algebraic differential equation is of finite order, so $\Psi(z)$ can be taken to be $\exp(\exp|z|)$ in this case.

To prove our main result (assuming R is not the whole plane), the equation is transformed to the unit disk using f^{-1} . The resulting equation no longer has polynomial coefficients, but careful estimates on the growth of the coefficients can be made using certain standard estimates on the growth of the univalent analytic function f^{-1} on the unit disk. Then by using results of Valiron [4; p. 299] concerning certain analytic functions in the unit disk, we obtain (Lemma C below) the same growth condition on solutions in the unit disk as was obtained in [4; p. 294] in the case where the coefficients were polynomials. Our main result then follows by retransforming back to R .

2. We now state our main result:

THEOREM. *Let R be a simply-connected region of the complex plane which is not the whole plane. Let f be a univalent analytic mapping of R onto the unit disk. (Such a mapping exists by the Riemann Mapping Theorem.) Then, if $h(z)$ is any analytic function on R which is a solution of a first order algebraic differential equation, then there exist a constant $A \geq 0$ and a compact set X contained in R , such that the inequality $|h(z)| \leq \exp((1 - |f(z)|)^{-A})$ holds for all $z \in R - X$. (Thus clearly, for each such $h(z)$, there is a compact set K contained in R such that the inequality*

$$|h(z)| < \exp(\exp((1 - |f(z)|)^{-1})) \text{ holds on } R - K.)$$

REMARK. If R is the whole plane, then by a result of Valiron [3; p. 41], any analytic function on R (i.e., any entire function) which is a solution of a first order algebraic differential equation is of finite order.

3. *Proof of the theorem.* Let $h(z)$ be analytic on the simply-connected region R and be a solution of a first order algebraic differential equation $\Omega(z, y, y') = 0$. We may write $\Omega(z, y, y')$ in the form,

$$(1) \quad \Omega(z, y, y') = \sum H_{kj}(z) y^k (y')^j$$

where the functions $H_{kj}(z)$ are polynomials in z . Thus,

$$(2) \quad \sum H_{kj}(z) (h(z))^k (h'(z))^j \equiv 0 \quad \text{on } R.$$

Let g be the inverse of f . Thus g is a univalent analytic mapping of the unit disk onto R . Since $h(z)$ is analytic on R , clearly the function

$$(3) \quad \varphi(\zeta) = h(g(\zeta)) \text{ is analytic on } |\zeta| < 1.$$

Since $\varphi'(\zeta) = h'(g(\zeta))g'(\zeta)$, we have by (2) that $\varphi(\zeta)$ satisfies the relation,

$$(4) \quad \sum F_{kj}(\zeta)(\varphi(\zeta))^k(\varphi'(\zeta))^j \equiv 0 \text{ on } |\zeta| < 1, \text{ where}$$

$$(5) \quad F_{kj}(\zeta) = H_{kj}(g(\zeta))/(g'(\zeta))^j \text{ for each } (k, j).$$

For each (k, j) such that $H_{kj} \not\equiv 0$, let $d(k, j)$ be the degree of the polynomial H_{kj} . Define

$$(6) \quad q = 1 + \max \{j + 2d(k, j) : H_{kj} \not\equiv 0\}.$$

Thus clearly,

$$(7) \quad q > 0.$$

We now prove,

LEMMA A. *There exists a constant $K^* > 0$ such that on any circle $|\zeta| = r$, where $r \in [0, 1)$, we have $|F_{kj}(\zeta)| \leq K^*(1 - r)^{-q}$ for all (k, j) .*

Proof. Since $g(\zeta)$ is univalent on $|\zeta| < 1$, the function $G(\zeta) = (g(\zeta) - g(0))/g'(0)$ is also univalent on $|\zeta| < 1$ and $G(0) = 0$ while $G'(0) = 1$. Thus by [1; Th. 17. 4. 7, p. 353], $|G(\zeta)| \leq r(1 - r)^{-2}$ on any circle $|\zeta| = r < 1$. Since $r < 1$ and $(1 - r)^2 \leq 1$, we clearly obtain

$$(8) \quad |g(\zeta)| \leq L(1 - r)^{-2} \text{ on } |\zeta| = r < 1$$

where $L = |g'(0)| + |g(0)| > 0$ (since $g'(0) \neq 0$). Again, since $G(\zeta)$ is univalent on $|\zeta| < 1$ and $G(0) = 0$ while $G'(0) = 1$ we have by [1; Th. 17. 4. 6, p. 351] that $(1 - r)(1 + r)^{-3} \leq |G'(\zeta)| \leq (1 + r)(1 - r)^{-3}$ on any circle $|\zeta| = r < 1$. Setting $K_1 = 2|g'(0)|$ and $K_2 = |g'(0)|/8$, we have

$$(9) \quad K_1 > 0 \text{ and } K_2 > 0 \text{ (since } g'(0) \neq 0), \text{ and}$$

$$(10) \quad K_2(1 - r) \leq |g'(\zeta)| \leq K_1(1 - r)^{-3} \text{ on } |\zeta| = r < 1.$$

Now consider any (k, j) . If $H_{kj} \equiv 0$ clearly $F_{kj} \equiv 0$ by (5), so we may assume $H_{kj} \not\equiv 0$. Since H_{kj} is of degree $d = d(k, j)$, we may write, $H_{kj}(z) = B_0 + B_1z + \cdots + B_dz^d$ where the B_j are constants. Thus by (8), we have on any circle $|\zeta| = r < 1$,

$$(11) \quad |H_{kj}(g(\zeta))| \leq |B_0| + |B_1| L(1-r)^{-2} + \cdots + |B_d| L^d(1-r)^{-2d}.$$

Thus clearly there is a constant $\alpha(k, j) > 0$ such that,

$$(12) \quad |H_{kj}(g(\zeta))| \leq \alpha(k, j)(1-r)^{-2d} \quad \text{on} \quad |\zeta| = r < 1.$$

By (10), we have $|g'(\zeta)|^j \geq K_2^j(1-r)^j$ on $|\zeta| = r < 1$, so by (12) and (5),

$$(13) \quad |F_{kj}(\zeta)| \leq (\alpha(k, j)/K_2^j)(1-r)^{-(j+2d)} \quad \text{on} \quad |\zeta| = r < 1.$$

Setting

$$K^* = 1 + \max \{ \alpha(k, j)/K_2^j : H_{kj} \neq 0 \}$$

and noting that $q > j + 2d(k, j)$ by (6), we have by (13) that for any (k, j)

$$(14) \quad |F_{kj}(\zeta)| \leq K^*(1-r)^{-q} \quad \text{on any circle} \quad |\zeta| = r < 1,$$

proving Lemma A.

Now set

$$(15) \quad p = \max \{ k + j : H_{kj} \neq 0 \}, \quad \text{and}$$

$$(16) \quad m = \max \{ j : H_{p-j, j} \neq 0 \}.$$

We now prove,

LEMMA B. *There exist constants $K^* > 0, \sigma \geq 0$ and $r_0 \in [0, 1)$ such that on any circle $|\zeta| = r$, where $r_0 < r < 1$, we have $|F_{p-m, m}(\zeta)| \geq K^*(1-r)^\sigma$.*

Proof. If the degree d of $H_{p-m, m}$ is zero, then $H_{p-m, m}$ is a non-zero constant function, say $H_{p-m, m}(z) \equiv L_1$. Then by (5) and (10), we have $|F_{p-m, m}(\zeta)| \geq (|L_1|/K_1^m)(1-r)^{3m}$ on $|\zeta| = r < 1$ which proves the lemma in this case.

Thus we may assume $d > 0$. Since $H_{p-m, m}(z)$ is a polynomial of degree $d \geq 1$, we may write,

$$(17) \quad H_{p-m, m}(z) = B(z - a_1) \cdots (z - a_d) \quad \text{where} \quad B \neq 0,$$

and hence

$$(18) \quad H_{p-m, m}(g(\zeta)) = B(g(\zeta) - a_1) \cdots (g(\zeta) - a_d) \quad \text{for} \quad |\zeta| < 1.$$

We partition the set of roots $\{a_1, \dots, a_d\}$ of $H_{p-m, m}(z)$ into three classes and arrange them as follows: a_1, \dots, a_b lie in the complement of the closure of R ; a_{b+1}, \dots, a_c lie in R and a_{c+1}, \dots, a_d lie on the boundary of R , for some b and c with $0 \leq b \leq c \leq d$.

We first consider any root $a_k, 1 \leq k \leq b$. Since a_k lies in the complement of the closure of R , the distance ε_k from a_k to R is strictly positive. But if $|\zeta| < 1$ then $g(\zeta)$ lies in R so we have,

$$(19) \quad |g(\zeta) - a_k| \geq \varepsilon_k > 0 \quad \text{for} \quad |\zeta| < 1 \quad \text{and} \quad k \in \{1, \dots, b\}.$$

We now consider any root $a_j, b+1 \leq j \leq c$. Thus $a_j \in R$ so there exists t_j in the unit disk $|\zeta| < 1$ such that $g(t_j) = a_j$. Let $\delta = \max\{|t_j| : b+1 \leq j \leq c\}$ and let $r_0 = (1 + \delta)/2$. Thus $\delta < r_0 < 1$. Let N be the image under g of the annulus $r_0 \leq |\zeta| < 1$. Then no a_j (for $b+1 \leq j \leq c$) can lie in the closure of N , for otherwise there would be a sequence $\{\zeta_n\}$ with $r_0 \leq |\zeta_n| < 1$ such that $\{g(\zeta_n)\} \rightarrow a_j$. By the continuity of g^{-1} we would obtain $\{\zeta_n\} \rightarrow t_j$. Since $|\zeta_n| \geq r_0$ for all n , we would then have $|t_j| \geq r_0$ which contradicts $|t_j| \leq \delta$ (since $\delta < r_0$). Thus the distance ε_j from a_j to N is strictly positive for $j = b+1, \dots, c$. Thus for $j \in \{b+1, \dots, c\}$ we have

$$(20) \quad |g(\zeta) - a_j| \geq \varepsilon_j > 0 \quad \text{for} \quad r_0 \leq |\zeta| < 1.$$

Finally, we consider any root $a_s, c+1 \leq s \leq d$, and we consider the function $G_s(\zeta) = (g(\zeta) - a_s)^{-1}$ defined on $|\zeta| < 1$. Since a_s lies on the boundary of R , the equation $g(\zeta) = a_s$ has no roots in $|\zeta| < 1$ by definition of the map $g(\zeta)$. Thus $G_s(\zeta)$ is analytic on $|\zeta| < 1$ and is univalent on $|\zeta| < 1$ since $g(\zeta)$ is. Thus the function

$$\psi_s(\zeta) = (G_s(\zeta) - G_s(0))/G'_s(0)$$

is univalent and analytic on $|\zeta| < 1$ and $\psi_s(0) = 0$ while $\psi'_s(0) = 1$. Thus by [1; Th. 17.4.6, p. 351], we have $|\psi'_s(\zeta)| \leq (1+r)(1-r)^{-3}$ on any circle $|\zeta| = r < 1$. Hence $|G'_s(\zeta)| \leq 2|G'_s(0)|(1-r)^{-3}$ on $|\zeta| = r < 1$. But $G'_s(\zeta) = -g'(\zeta)(g(\zeta) - a_s)^{-2}$, so we obtain,

$$(21) \quad |g(\zeta) - a_s|^2 \geq (1-r)^3 |g'(\zeta)|/2 |G'_s(0)| \quad \text{on} \quad |\zeta| = r < 1.$$

But by (10), $|g'(\zeta)| \geq K_2(1-r)$ on $|\zeta| = r < 1$, where $K_2 > 0$. Thus clearly,

$$(22) \quad |g(\zeta) - a_s|^2 \geq (K_2/2 |G'_s(0)|)(1-r)^4 \quad \text{on} \quad |\zeta| = r < 1.$$

Setting $M_s = (K_2/2 |G'_s(0)|)^{1/2}$, we have $M_s > 0$ and

$$(23) \quad |g(\zeta) - a_s| \geq M_s(1-r)^2 \quad \text{on} \quad |\zeta| = r < 1 \quad \text{for} \quad s \in \{c+1, \dots, d\}.$$

Thus in view of (18), (19), (20) and (23), we obtain

$$(24) \quad |H_{p-m,m}(g(\zeta))| \geq K_0(1-r)^{2(d-c)} \quad \text{on} \quad |\zeta| = r,$$

when

$$r_0 \leq r < 1, \quad \text{where} \quad K_0 = |B| \varepsilon_1 \dots \varepsilon_c M_{c+1} \dots M_d > 0.$$

Now by (10), $|g'(\zeta)|^m \leq K_1^m(1-r)^{-3m}$ on $|\zeta| = r < 1$, so in view of (5) and (24), we obtain

$$(25) \quad |F_{p-m,m}(\zeta)| \geq K^\sharp(1-r)^\sigma \quad \text{on} \quad |\zeta| = r, \quad \text{when} \quad r_0 \leq r < 1,$$

where $K^\sharp = K_0/K_1^m > 0$ and $\sigma = 2(d-c) + 3m \geq 0$. This proves Lemma B.

Now $\varphi(\zeta)$ satisfies the relation (4) in $|\zeta| < 1$. We introduce the notation $M(r, \varphi) = \max_{|\zeta|=r} |\varphi(\zeta)|$ for $r \in [0, 1)$. We now prove,

LEMMA C. *There exist real numbers $A \geq 0$ and $r^* \in [0, 1)$ such that for any $r \in [r^*, 1)$, we have $M(r, \varphi) \leq \exp((1-r)^{-A})$.*

Proof. If $M(r, \varphi)$ is bounded on $[0, 1)$, then clearly the result holds. Hence we may assume $M(r, \varphi)$ is unbounded so $M(r, \varphi) \rightarrow +\infty$ as $r \rightarrow 1$.

We proceed by contradiction and assume the lemma is false. Then in the terminology of [4; p. 294], $\varphi(\zeta)$ is of order ∞ in $|\zeta| < 1$. Let $\sum_{j=0}^{\infty} c_j \zeta^j$ be the power series expansion of $\varphi(\zeta)$. For each $r \in [0, 1)$, let $N(r) = \max_{j \geq 0} |c_j| r^j$ and $n(r) = \max \{j: |c_j| r^j = N(r)\}$. Then since $\varphi(\zeta)$ is of order ∞ , by [4; p. 299], for any $b \in (0, 1)$, there is in $(0, 1)$ a sequence of values of r (called *remarkable*) tending to one, such that

$$(26) \quad \log M(r, \varphi) > \gamma'(n(r))^\delta \quad \text{and} \quad n(r) > \gamma''(1-r)^{-\delta}$$

where $\delta = (1-b)^{-1}$ and γ', γ'' are strictly positive constants independent of r , and such that at every point of $|\zeta| = r$ at which $|\varphi(\zeta)| = M(r, \varphi)$ we have

$$(27) \quad \varphi'(\zeta) = (1 + \varepsilon(\zeta))(n(r)/\zeta)\varphi(\zeta)$$

where $\varepsilon(\zeta)$ tends uniformly to zero as $r = |\zeta|$ tends to one.

Let q be as in (6), so $q > 0$ by (7). Let σ be as in Lemma B, so $\sigma \geq 0$. Thus $q + \sigma > 0$, so $b = (q + \sigma)/(q + \sigma + 1)$ belongs to $(0, 1)$. It is for this value of b that we apply the Valiron theory (26), (27). Then the corresponding $\delta = q + \sigma + 1$, so

$$(28) \quad \delta > q + \sigma > 0.$$

Let p and m be as in (15) and (16). Then clearly by (5),

$$(29) \quad p = \max \{k + j: F_{kj} \neq 0\} \quad \text{and} \quad m = \max \{j: F_{p-j,j} \neq 0\}.$$

Now let $r \in (0, 1)$ and let ζ be a point on $|\zeta| = r$ at which $|\varphi(\zeta)| = M(r, \varphi)$. Then $\varphi(\zeta) \neq 0$ and so by dividing (4) through by

$(\varphi(\zeta))^p$ (and using (29)), we can write (4) as

$$(30) \quad \sum_{j=0}^m F_{p-j,j}(\zeta)(\varphi'(\zeta)/\varphi(\zeta))^j \equiv -\sum_{k+j < p} F_{k,j}(\zeta)(\varphi'(\zeta)/\varphi(\zeta))^j (\varphi(\zeta))^{k+j-p}.$$

We will denote the left side of (30) by $\Lambda(\zeta)$, and the right side by $\Gamma(\zeta)$.

We now assert that there exist real numbers $L^* > 0$ and $r^* \in (1/2, 1)$ such that if $r \in (r^*, 1)$ is a remarkable value, then

$$(31) \quad |\Gamma(\zeta)| \leq L^* (1-r)^{-q} (M(r, \varphi))^{-1/2}$$

for any point on $|\zeta| = r$ at which $|\varphi(\zeta)| = M(r, \varphi)$.

To prove (31), we note first that for remarkable $r \rightarrow 1$, we have $n(r) \rightarrow +\infty$ by (26). Thus since $M(r, \varphi) \rightarrow +\infty$ and $\varepsilon(\zeta) \rightarrow 0$ for remarkable $r = |\zeta| \rightarrow 1$, we can choose $r^* \in (1/2, 1)$ such that for remarkable $r \in (r^*, 1)$, we have $M(r, \varphi) > 1$, $n(r) > 1$ and $|\varepsilon(\zeta)| < 1/2$ whenever $|\zeta| = r$ and $|\varphi(\zeta)| = M(r, \varphi)$. Now let $r \in (r^*, 1)$ be a remarkable value and let ζ be a point on $|\zeta| = r$ at which $|\varphi(\zeta)| = M(r, \varphi)$. We refer to the right side of (30). If $k+j < p$, then $p - (k+j) \geq 1$, so

$$(32) \quad |\varphi(\zeta)|^{k+j-p} \leq (M(r, \varphi))^{-1}.$$

Since $|\varepsilon(\zeta)| < 1/2$ and $|\zeta| = r > 1/2$, we have by (27) that

$$(33) \quad |\varphi'(\zeta)/\varphi(\zeta)| \leq 4n(r).$$

By Lemma A, there is a constant $K^* > 0$ such that $|F_{k,j}(\zeta)| \leq K^*(1-r)^{-q}$ for all (k, j) . Thus by (32), (33) and the definition of $\Gamma(\zeta)$, it follows that

$$(34) \quad |\Gamma(\zeta)| \leq K_3 (1-r)^{-q} (n(r))^\theta (M(r, \varphi))^{-1}$$

where $K_3 > 0$ and $\theta \geq 0$ are constants (independent of r). But by (26), $\gamma'(n(r))^b < \log M(r, \varphi)$, so by (34),

$$(35) \quad |\Gamma(\zeta)| \leq L^* (1-r)^{-q} (\log M(r, \varphi))^{\theta/b} (M(r, \varphi))^{-1}$$

where $L^* > 0$ is a constant (independent of r). But since $M(r, \varphi) \rightarrow +\infty$ as $r \rightarrow 1$, clearly there exists $r' \in (0, 1)$ such that if $r \in (r', 1)$, then $(\log M(r, \varphi))^{\theta/b} < (M(r, \varphi))^{1/2}$. Setting $r^* = \max\{r^*, r'\}$, we see that (31) follows from (35).

We now consider $\Lambda(\zeta) = \sum_{j=0}^m F_{p-j,j}(\zeta)(\varphi'(\zeta)/\varphi(\zeta))^j$.

Case I. $m = 0$. Then $\Lambda(\zeta) = F_{p-m,m}(\zeta)$. By Lemma B there exist constants $K^* > 0$ and $r_0 \in (r^*, 1)$, such that on any circle $|\zeta| = r$ where $r \in (r_0, 1)$, we have

$$(36) \quad |A(\zeta)| = |F_{p-m,m}(\zeta)| \geq K^*(1-r)^\sigma.$$

Since $A(\zeta) = I(\zeta)$ (by (30)), we have by (31) and (36) that for all remarkable $r \in (r_0, 1)$, $K^*(1-r)^\sigma \leq L^*(1-r)^{-q}(M(r, \varphi))^{-1/2}$ and hence

$$(37) \quad (1-r)^{2(q+\sigma)}M(r, \varphi) \leq (L^*/K^*)^2.$$

But by (26), there exist constants $\gamma' > 0, \gamma'' > 0$, independent of r , such that $\log M(r, \varphi) > \gamma'(\gamma'')^b(1-r)^{-b\delta}$. Hence with (37), we see that for all remarkable $r \in (r_0, 1)$,

$$(38) \quad (1-r)^{2(q+\sigma)} \exp(\gamma(1-r)^{-b\delta}) \leq (L^*/K^*)^2$$

where $\gamma = \gamma'(\gamma'')^b > 0$. Since $b\delta = q + \sigma > 0$ (by (28)), it is clear that (38) is impossible since the left side of (38) tends to $+\infty$ as $r \rightarrow 1$. This contradiction proves Lemma C in the case $m = 0$.

Case II. $m > 0$. Since $\varepsilon(\zeta)$ (in (27)) tends to zero as $r = |\zeta| \rightarrow 1$, we can choose $r_1 \in (0, 1)$ such that for any remarkable $r \in (r_1, 1)$, we have $|\varepsilon(\zeta)| < 1/2$ for those points ζ on $|\zeta| = r$ at which $|\varphi(\zeta)| = M(r, \varphi)$. Hence for those ζ , we have by (27),

$$|\zeta\varphi'(\zeta)/n(r)\varphi(\zeta)| \geq 1 - |\varepsilon(\zeta)| > 1/2.$$

Since $|\zeta| < 1$, we have $|\varphi'(\zeta)/\varphi(\zeta)| > n(r)/2$ and so by (26), we have

$$(39) \quad |\varphi'(\zeta)/\varphi(\zeta)| > (\gamma''/2)(1-r)^{-\delta} \quad \text{where} \quad \gamma'' > 0.$$

We now assert that there exists a constant $K_0 > 0$, such that for all sufficiently large remarkable r , we have

$$(40) \quad |A(\zeta)| \geq K_0(1-r)^{\sigma-m\delta}$$

at all points of $|\zeta| = r$ at which $|\varphi(\zeta)| = M(r, \varphi)$.

To prove (40), we consider the quotients,

$$Q_j(\zeta) = (F_{p-j,j}(\zeta)(\varphi'(\zeta)/\varphi(\zeta))^j)/(F_{p-m,m}(\zeta)(\varphi'(\zeta)/\varphi(\zeta))^m),$$

for $j = 0, 1, \dots, m-1$. By Lemma A, there is a constant $K^* > 0$ such that $|F_{p-j,j}(\zeta)| \leq K^*(1-r)^{-q}$ on any circle $|\zeta| = r < 1$, for all j , while by Lemma B, there is a constant $K^* > 0$ such that for all sufficiently large $r \in (0, 1)$, we have

$$(41) \quad |F_{p-m,m}(\zeta)| \geq K^*(1-r)^\sigma \quad \text{on} \quad |\zeta| = r.$$

Thus, with (39), we see that for $j = 0, 1, \dots, m-1$, there is a constant $L_j > 0$ such that

$$(42) \quad |Q_j(\zeta)| \leq L_j(1-r)^{(m-j)\delta-(q+\sigma)}$$

for all sufficiently large remarkable r and all points ζ on $|\zeta| = r$ at which $|\varphi(\zeta)| = M(r, \varphi)$. Since $(m - j)\delta \geq \delta > q + \sigma$ (by (28)), we see that the right side of (42) tends to zero as $r \rightarrow 1$, so for all sufficiently large remarkable r , we have

$$(43) \quad |Q_j(\zeta)| \leq 1/(m + 1) \quad \text{for } j = 0, 1, \dots, m - 1,$$

at all points on $|\zeta| = r$ at which $|\varphi(\zeta)| = M(r, \varphi)$. Now by definition of A , we have

$$A(\zeta) = F_{p-m, m}(\zeta)(\varphi'(\zeta)/\varphi(\zeta))^m \left(1 + \sum_{j=0}^{m-1} Q_j(\zeta)\right),$$

and so by (43), we have

$$(44) \quad |A(\zeta)| \geq (1/(m + 1)) |F_{p-m, m}(\zeta)(\varphi'(\zeta)/\varphi(\zeta))^m|$$

for all sufficiently large remarkable r and all points ζ on $|\zeta| = r$ at which $|\varphi(\zeta)| = M(r, \varphi)$. Now it is clear that (40) follows from (39), (41) and (44).

Since $\Gamma(\zeta) = A(\zeta)$ (by (30)), we have by (31) and (40) that for all sufficiently large remarkable r , $K_0(1 - r)^{\sigma - m\delta} \leq L^*(1 - r)^{-q}(M(r, \varphi))^{-1/2}$, and so

$$(45) \quad (1 - r)^{2(q + \sigma - m\delta)} M(r, \varphi) \leq (L^*/K_0)^2.$$

But by (26), $\log M(r, \varphi) > \gamma'(\gamma'')^b(1 - r)^{-b\delta}$ (where $\gamma', \gamma'' > 0$), and so for all sufficiently large remarkable r , we have

$$(46) \quad (1 - r)^{2(q + \sigma - m\delta)} \exp(\gamma(1 - r)^{-b\delta}) \leq (L^*/K_0)^2$$

where $\gamma = \gamma'(\gamma'')^b > 0$. Since $b\delta = q + \sigma > 0$ (by (28)), it is clear that (46) is impossible, since the left side of (46) tends to $+\infty$ as $r \rightarrow 1$. This contradiction proves Lemma C in Case II, and thus the proof of Lemma C is complete.

We now conclude the proof of the theorem. By Lemma C, there exist constants $A \geq 0$ and $r^* \in [0, 1)$, such that if $r \in [r^*, 1)$, then

$$(47) \quad |\varphi(\zeta)| \leq \exp((1 - r)^{-A}) \quad \text{on } |\zeta| = r < 1.$$

Let X be the image under g of the closed disk $|\zeta| \leq r^*$. Then X is a compact set contained in R . If $z \in R - X$, then we have $r^* < |f(z)| < 1$ (since $f = g^{-1}$), and so by (47), we have

$$(48) \quad |\varphi(f(z))| \leq \exp((1 - |f(z)|)^{-A}).$$

But $\varphi = h \circ g$ (by (3)) and so $h(z) = \varphi(f(z))$. Thus

$$|h(z)| \leq \exp((1 - |f(z)|)^{-d})$$

if $z \in R - X$, which concludes the proof of the theorem.

BIBLIOGRAPHY

1. E. Hille, *Analytic function theory*, Volume II, Ginn and Company, Boston, 1962.
2. E. Lindelöf, *Sur la croissance des intégrales des équations différentielles algébrique du premier ordre*, Bull. Soc. Math. France **27** (1899), 205-215.
3. G. Valiron, *Sur les fonctions entières vérifiant une class d'équations différentielles*, Bull. Soc. Math. France **51** (1923), 33-45.
4. ———, *Fonctions analytiques et équations différentielles*, J. Math. Pures et. Appl. **31** (1952), 293-303.
5. T. Vijayaraghavan, *Sur la croissance des fonctions définies par les équations différentielles*, C. R. Acad. Sci. Paris **194** (1932), 827-829.

Received May 13, 1968. This research was supported in part by the National Science Foundation (GP 7374).

UNIVERSITY OF ILLINOIS
URBANA, ILLINOIS

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

BASIL GORDON
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics

Vol. 31, No. 3

BadMonth, 1969

George E. Andrews, <i>On a calculus of partition functions</i>	555
Silvio Aurora, <i>A representation theorem for certain connected rings</i>	563
Lawrence Wasson Baggett, <i>A note on groups with finite dual spaces</i>	569
Steven Barry Bank, <i>On majorants for solutions of algebraic differential equations in regions of the complex plane</i>	573
Klaus R. Bichteler, <i>Locally compact topologies on a group and the corresponding continuous irreducible representations</i>	583
Mario Borelli, <i>Affine complements of divisors</i>	595
Carlos Jorge Do Rego Borges, <i>A study of absolute extensor spaces</i>	609
Bruce Langworthy Chalmers, <i>Subspace kernels and minimum problems in Hilbert spaces with kernel function</i>	619
John Dauns, <i>Representation of L-groups and F-rings</i>	629
Spencer Ernest Dickson and Kent Ralph Fuller, <i>Algebras for which every indecomposable right module is invariant in its injective envelope</i>	655
Robert Fraser and Sam Bernard Nadler, Jr., <i>Sequences of contractive maps and fixed points</i>	659
Judith Lee Gersting, <i>A rate of growth criterion for universality of regressive isols</i>	669
Robert Fred Gordon, <i>Rings in which minimal left ideals are projective</i>	679
Fred Gross, <i>Entire functions of several variables with algebraic derivatives at certain algebraic points</i>	693
W. Charles (Wilbur) Holland Jr. and Stephen H. McCleary, <i>Wreath products of ordered permutation groups</i>	703
W. J. Kim, <i>The Schwarzian derivative and multivalence</i>	717
Robert Hamor La Grange, Jr., <i>On $(m - n)$ products of Boolean algebras</i>	725
Charles D. Masiello, <i>The average of a gauge</i>	733
Stephen H. McCleary, <i>The closed prime subgroups of certain ordered permutation groups</i>	745
Richard Roy Miller, <i>Gleason parts and Choquet boundary points in convolution measure algebras</i>	755
Harold L. Peterson, Jr., <i>On dyadic subspaces</i>	773
Derek J. S. Robinson, <i>Groups which are minimal with respect to normality being intransitive</i>	777
Ralph Edwin Showalter, <i>Partial differential equations of Sobolev-Galpern type</i>	787
David Slepian, <i>The content of some extreme simplexes</i>	795
Joseph L. Taylor, <i>Noncommutative convolution measure algebras</i>	809
B. S. Yadav, <i>Contractions of functions and their Fourier series</i>	827
Lindsay Nathan Childs and Frank Rimi DeMeyer, <i>Correction to: "On automorphisms of separable algebras"</i>	833
Moses Glasner and Richard Emanuel Katz, <i>Correction to: "Function-theoretic degeneracy criteria for Riemannian manifolds"</i>	834
Satish Shirali, <i>Correction to: "On the Jordan structure of complex Banach *-algebras"</i>	834
Benjamin Rigler Halpern, <i>Addendum to: "Fixed points for iterates"</i>	834