Pacific Journal of Mathematics

THE CLOSED PRIME SUBGROUPS OF CERTAIN ORDERED PERMUTATION GROUPS

STEPHEN H. MCCLEARY

Vol. 31, No. 3

BadMonth 1969

THE CLOSED PRIME SUBGROUPS OF CERTAIN ORDERED PERMUTATION GROUPS

STEPHEN H. MCCLEARY

The group $G = A(\Omega)$ of all order-preserving permutations of a chain Ω becomes a lattice-ordered group when ordered pointwise, i.e., $f \leq g$ if and only if $\beta f \leq \beta g$ for all $\beta \in \Omega$. Lloyd showed that for each $\omega \in \Omega$, the stabilizer subgroup $G_{\omega} =$ $\{g \in G \mid \omega g = \omega\}$ is a closed prime subgroup of G. Our main result (Theorem 11) states that besides G itself, these subgroups, together with the stabilizer subgroups of Dedekind cuts of Ω , comprise all of the closed prime subgroups of G.

Actually, G need not be all of $A(\Omega)$. In § 2, we use Lloyd's result to show that for depressible or complete subgroups G of $A(\Omega)$, all stabilizer subgroups are closed. We find in § 3 that every closed convex *l*-subgroup C of G is of the form $C = \{g \in G \mid \overline{Ag} = \overline{A}\}$ for some collection \overline{A} of points and cuts of Ω . In § 4 we prove the main theorem (stated above) for depressible groups. Finally, this theorem is applied to the question of the extent to which the *l*-group G determines the chain Ω , which was first considered by Holland [4]; and to the determination of the *l*-automorphisms of G, considered by Lloyd [5].

2. Stabilizer subgroups. Let Ω be a chain. A permutation g of Ω is said to *preserve order* if $\alpha \leq \beta$ implies $\alpha g \leq \beta g$ for all $\alpha, \beta \in \Omega$. The group $A(\Omega)$ of all order-preserving permutations (o-permutations) of Ω , ordered pointwise, is a lattice-ordered group (*l*-group). $A(\Omega)$ is not assumed to be transitive. For elementary information about *l*-groups, see [1].

 $\overline{\Omega}$ will denote the completion of Ω by Dedekind cuts (without end points unless these end points belong to Ω). We shall consider Ω to be a subchain of $\overline{\Omega}$. Each $g \in A(\Omega)$ can be extended to $\overline{\Omega}$ by defining $\overline{\omega}g(\overline{\omega}\in\overline{\Omega})$ to be sup $\{\beta g \mid \beta \in \Omega, \beta \leq \overline{\omega}\}$. Thus $A(\Omega)$ can be considered to be an *l*-subgroup of $A(\overline{\Omega})$, i.e., a subgroup which is also a sublattice.

Let G be an *l*-subgroup of $A(\Omega)$, and thus also of $A(\overline{\Omega})$. G, or more properly the pair (G, Ω) , is called an ordered permutation group. Holland [3, Th. 2] showed that every abstract *l*-group L is *l*-isomorphic to such a G. A subgroup H of G is convex if $h_1 \leq g \leq h_2$, with $h_1, h_2 \in H$ and $g \in G$, implies $g \in H$. H is a prime subgroup of G if H is a convex *l*-subgroup of G and if $g_1 \wedge g_2 = 1$, with $g_1, g_2 \in G$, implies $g_1 \in H$ or $g_2 \in H$. For $\overline{\omega} \in \overline{\Omega}$, we define the stabilizer subgroup $G_{\overline{\omega}}$ to be

$$\{g\in G\,|\,ar{\omega}g\,=\,ar{\omega}\}$$
 ,

S. H. McCLEARY

a prime subgroup of G. Thus we have a stabilizer subgroup for each Dedekind cut of Ω (not just for points of Ω).

We wish to consider two classes of *l*-subgroups G of $A(\Omega)$ for which the stabilizer subgroups $G_{\bar{\omega}}$ are *closed*, i.e., if $s = \sup \{s_i \mid i \in I\}$, with $s \in G$ and $s_i \in G_{\bar{\omega}}$ for each *i* in an index set *I*, then $s \in G_{\bar{\omega}}$.

For $g \in A(\Omega)$ and $\gamma \in \Omega$ such that $\gamma g \neq \gamma$, the *interval of support* $I(g, \gamma)$ of g which contains γ is $\{\beta \in \Omega \mid \gamma g^m \leq \beta \leq \gamma g^n \text{ for some integers } m \text{ and } n\}$. An *l*-subgroup G of $A(\Omega)$ will be called *depressible* [7] if it shares with $A(\Omega)$ the following property: If Δ is an interval of support of $g \in G$, so that $\Delta g = \Delta$, then there exists $h \in G$ such that $\beta h = \beta g$ if $\beta \in \Delta$, but $\beta h = \beta$ if $\beta \notin \Delta$. Intuitively, h is obtained by *depressing* g outside Δ . As noted in [7], convex *l*-subgroups of $A(\Omega)$ are depressible.

G is a complete *l*-subgroup of $A(\Omega)$ if whenever $g \in G$ is the sup in G of a collection $\{g_i \mid i \in I\}$ of elements of G, then g is also the sup in $A(\Omega)$ of $\{g_i \mid i \in I\}$. By [2, Th. 3.10], an abstract *l*-group L is *l*isomorphic to a complete *l*-subgroup G of some $A(\Omega)$ if and only if L is completely distributive, i.e., if $\bigwedge_{i \in I} \bigvee_{k \in K} g_{ik} = \bigvee_{f \in K^I} \bigwedge_{i \in I} g_{if(i)}$ for any collection $\{g_{ik} \mid i \in I, k \in K\}$ of elements of G for which the indicated sups and infs exist.

THEOREM 1. Suppose G is a depressible or complete l-subgroup of $A(\Omega)$. Then for each $\bar{\omega} \in \bar{\Omega}$, $G_{\bar{\omega}}$ is a closed prime subgroup of G.

Proof. Lloyd [6, Th. 2] proved that if $G = A(\Omega)$, then the stabilizer subgroups G_{ω} of points $\omega \in \Omega$ are closed. As noted in [7], Lloyd's proof also works when G is a depressible *l*-subgroup of $A(\Omega)$. If (G, Ω) is depressible, the extension $(G, \overline{\Omega})$ is also depressible. But $G_{\overline{\omega}}$ is the stabilizer subgroup of a point for the group $(G, \overline{\Omega})$, and hence must be closed. On the other hand, suppose that G is complete. If s = $\sup \{s_i \mid i \in I\}$, with $s \in G$ and each $s_i \in G_{\overline{\omega}}$, then since G is complete, s is also the sup in $A(\Omega)$ of $\{s_i \mid i \in I\}$. Since the stabilizer subgroups are closed for $A(\Omega)$, $\overline{\omega}s = \overline{\omega}$, and then $s \in G_{\overline{\omega}}$. Hence again $G_{\overline{\omega}}$ is closed. (Incidentally, by [7, Th. 7], if G is transitive on Ω , then G is complete if and only if the stabilizer subgroups $G_{\overline{\omega}}$ are closed.)

A word of warning: $A(\Omega)$, extended to $\overline{\Omega}$, need not be all of $A(\overline{\Omega})$. Thus the above proof, even for $G = A(\Omega)$, does not work without the introduction of the concept of depressible groups.

3. The closed convex *l*-subgroups of *G*. In this section we determine the closed convex *l*-subgroups of *G*, where *G* is an *l*-subgroup of $A(\Omega)$ which is either depressible or complete. If *C* is a subgroup of *G*, we define FxC to be $\{\overline{\tau} \in \overline{\Omega} \mid \overline{\tau}C = \overline{\tau}\}$, the collection of points in $\overline{\Omega}$ fixed by *C*. By the convexification Conv (\overline{A}) of a subset \overline{A} of $\overline{\Omega}$,

we shall mean $\{\overline{\beta} \in \overline{\Omega} \mid \overline{\gamma} \leq \overline{\beta} \leq \overline{\delta} \text{ for some } \overline{\gamma}, \overline{\delta} \in \overline{A}\}$. If C is a subgroup of G and $\overline{\omega} \in \overline{\Omega} \setminus FxC$, we define the *orbital* of C containing $\overline{\omega}$ to be Conv $(\overline{\omega}C)$.

LEMMA 2. If an orbital Γ of C is bounded above, then $\sup \Gamma \in FxC$; and dually.

Proof. $\Gamma C = \Gamma$, so $(\sup \Gamma)C = \sup \Gamma$.

If L is any *l*-group and C is a convex *l*-subgroup of L, then the closure C^* of C means the smallest closed convex *l*-subgroup of L which contains C.

THEOREM 3. Suppose that G is a depressible or complete l-subgroup of $A(\Omega)$ and that C is a convex l-subgroup of G. Then $C^* = \{g \in G \mid \bar{\omega}g = \bar{\omega} \text{ for all } \bar{\omega} \in FxC\}.$

Proof. Let $D = \{g \in G \mid \bar{\omega}g = \bar{\omega} \text{ for all } \bar{\omega} \in FxC\}$. Then $D = \bigcap \{G_{\bar{\omega}} \mid \bar{\omega} \in FxC\}$, so D is a closed convex *l*-subgroup of G. Certainly $C^* \subseteq D$.

Now pick any $1 \leq g \in D$ and any $\beta \in \Omega$. Conv $(\beta C^*) = \text{Conv}(\beta D)$, for otherwise the lemma would imply that either $\sup(\beta C^*)$ or $\inf(\beta C^*)$ would be an element of $FxC^* \setminus FxD$. Hence there exists $f \in C^*$ such that $\beta g \leq \beta f$. Then $1 \leq (g \land f) \lor 1 \leq f \lor 1 \in C^*$, so $(g \land f) \lor 1 \in C^*$; and $\beta((g \land f) \lor 1) = \beta g$. Thus for each $\beta \in \Omega$, there exists an $s_\beta \in C^*$ such that $\beta s_\beta = \beta g$ and $s_\beta \leq g$. Hence $g = \sup\{s_\beta \mid \beta \in \Omega\}$. Since C^* is closed, $g \in C^*$. Since g was an arbitrary element of D, we have shown that $C^* = D$.

COROLLARY 4. Suppose that G is a depressible or complete l-subgroup of $A(\Omega)$. Then the closed convex l-subgroups of G can be characterized as those subsets C of G for which there exist $\overline{\mathcal{A}}_c \subseteq \overline{\Omega}$ such that $C = \{g \in G \mid \overline{\omega}g = \overline{\omega} \text{ for all } \overline{\omega} \in \overline{\mathcal{A}}_c\}$. Moreover, every closed convex l-subgroup of G is the intersection of a collection of closed prime subgroups $G_{\overline{\omega}}$ of G. Hence the maximal closed convex l-subgroups of G (if any) must be prime.

COROLLARY 5. Let L be a completely distributive abstract l-group. Then every closed convex l-subgroup of L is the intersection of a collection of closed prime subgroups of L; and the maximal closed convex l-subgroup of L (if any) are prime.

Proof. By [2, Th. 3.10], L is *l*-isomorphic to a complete subgroup

G of some $A(\Omega)$.

4. The closed prime subgroups of G. We now develop the tools needed to determine (for the depressible case) that of the closed convex *l*-subgroups of G, those which are prime are (besides G itself) precisely the stabilizer subgroups. First we establish some preliminary results which apply to any *l*-subgroup G of $A(\Omega)$. Let us define two binary relations on $\overline{\Omega}: \overline{\omega}F\overline{\tau}$ (read $\overline{\omega}$ fixes $\overline{\tau}$) if $\overline{\tau} \in FxG_{\overline{\omega}}$, and $\overline{\omega}M\overline{\tau}$ (read $\overline{\omega}$ moves $\overline{\tau}$) if $\overline{\tau} \notin FxG_{\overline{\omega}}$.

As a strong hint that the only closed prime subgroups of G are the stabilizer subgroups $G_{\bar{\omega}}$, we have

PROPOSITION 6. If $\bar{\omega}M\bar{\tau}$ and $\bar{\tau}M\bar{\omega}$, then $G_{\bar{\omega},\bar{\tau}} = G_{\bar{\omega}} \cap G_{\bar{\tau}}$ is not prime.

Proof. Since $\bar{\omega}M\bar{\tau}$, we may pick $1 < g \in G_{\bar{\omega}}$ such that $\bar{\tau}g > \bar{\tau}$; and since $\bar{\tau}M\bar{\omega}$, we may pick $1 < h \in G_{\bar{\tau}}$ such that $\bar{\omega}h > \bar{\omega}$. Then $g \wedge h \in G_{\bar{\omega},\bar{\tau}}$ while neither g nor h lies in $G_{\bar{\omega},\bar{\tau}}$, so $G_{\bar{\omega},\bar{\tau}}$ is not prime.

PROPOSITION 7. $\bar{\omega}F\bar{\tau}$ if and only if $G_{\bar{\omega}} \subseteq G_{\bar{\tau}}$. Hence $G_{\bar{\omega}} = G_{\bar{\tau}}$ if and only if $\bar{\omega}F\bar{\tau}$ and $\bar{\tau}F\bar{\omega}$.

LEMMA 8. Suppose Γ is an orbital of $G_{\overline{\tau}}, \overline{\tau} \in \overline{\Omega}$. Then for $g \in G$, Γg is an orbital of $G_{\overline{\tau}g}$.

Proof. $\Gamma = \operatorname{Conv}(\bar{\omega}G_{\bar{\tau}})$ for some $\bar{\omega} \in \bar{\Omega}$, so $\Gamma g = \operatorname{Conv}(\bar{\omega}G_{\bar{\tau}}g) = \operatorname{Conv}((\bar{\omega}g)(g^{-1}G_{\bar{\tau}}g)) = \operatorname{Conv}((\bar{\omega}g)G_{\bar{\tau}g}).$

LEMMA 9. Suppose Γ is an orbital of $G_{\overline{\tau}}, \overline{\tau} \in \overline{\Omega}$. If $\overline{\tau}g = \overline{\tau}h$, with $g, h \in G$, then $\Gamma g = \Gamma h$.

Proof. If $\overline{\tau}g = \overline{\tau}h$, then $gh^{-1} \in G_{\overline{\tau}}$, so $\Gamma gh^{-1} = \Gamma$ (since Γ is an orbital of $G_{\overline{\tau}}$), and thus $\Gamma g = \Gamma h$.

By an *o-block* of G, or more properly of $(G, \overline{\Omega})$, we mean a nonempty convex subset $\overline{\Delta}$ of $\overline{\Omega}$ such that for any $g \in G$, either $\overline{\Delta}g = \overline{\Delta}$ or $\overline{\Delta}g \cap \overline{\Delta} = \square$. (\square denotes the empty set.) An *o*-block is *trivial* if it contains only one point.

LEMMA 10. If $\bar{\omega}F\bar{\tau}$, but $\bar{\tau}M\bar{\omega}$, then the orbital $\Gamma = \text{Conv}(\bar{\omega}G_{\bar{\tau}})$ of $G_{\bar{\tau}}$ is a nontrivial o-block of G. For $g \in G$, $\Gamma g = \Gamma$ if and only if $g \in G_{\bar{\tau}}$.

Proof. If $g \in G_{\overline{\tau}}$, then $\Gamma g = \Gamma 1 = \Gamma$ by Lemma 9. We now show that if $\overline{\tau} < \overline{\tau}g$, then $\Gamma g \cap \Gamma = \square$. If not, we use Lemma 8 to pick

748

 $h \in G_{\overline{\tau}g}$ such that $(\overline{\omega}g)h \in \Gamma g \cap \Gamma$, and then we pick $k \in G_{\overline{\tau}}$ such that $(\overline{\omega}gh)k \leq \overline{\omega}$. Then $\overline{\omega}(ghk \vee 1) = \overline{\omega}$, but $\overline{\tau} < \overline{\tau}g = \overline{\tau}gh$, so $\overline{\tau} = \overline{\tau}k < \overline{\tau}ghk \leq \overline{\tau}(ghk \vee 1)$. This contradicts the assumption that $\overline{\omega}F\overline{\tau}$. Similarly, if $\overline{\tau} > \overline{\tau}g, \Gamma g \cap \Gamma = \Box$.

MAIN THEOREM 11. Suppose G is a depressible l-subgroup of $A(\Omega)$. Then the closed prime subgroups of G (besides G itself) are precisely the stabilizer subgroups $G_{\overline{w}}, \overline{\omega} \in \overline{\Omega}$.

Proof. By Theorem 1, the $G_{\bar{\omega}}$'s are closed prime subgroups of G. Now let C be any closed prime subgroup of G. By Corollary 4, $C = \{g \in G \mid \bar{\omega}g = \bar{\omega} \text{ for all } \bar{\omega} \in FxC\}$. If $FxC = \Box$, C = G. Now suppose $FxC \neq \Box$. For each $\bar{\omega} \in FxC$, $C \subseteq G_{\bar{\omega}}$. In any *l*-group, the collection of convex *l*-subgroups containing a given prime subgroup is a chain under inclusion [3, Lemma 3], so $\mathscr{S} = \{G_{\bar{\omega}} \mid \bar{\omega} \in FxC\}$ is a chain. Moreover $\bigcap \mathscr{S} = C$.

If for $G_{\overline{\omega}}, G_{\overline{\tau}} \in \mathscr{S}$, we have $G_{\overline{\omega}} \subset G_{\overline{\tau}}$ (i.e., if $\overline{\omega}F\overline{\tau}$, but $\overline{\tau}M\overline{\omega}$), then by Lemma 10, $\overline{\mathcal{A}}(\overline{\omega}, \overline{\tau}) = \operatorname{Conv}(\overline{\omega}G_{\overline{\tau}})$ is a nontrivial o-block of G. We show next that $\overline{\mathcal{A}}(\overline{\omega}, \overline{\tau})$ is independent of $\overline{\omega}$. For suppose that also $G_{\overline{\sigma}} \in \mathscr{S}$, and $G_{\overline{\sigma}} \subseteq G_{\overline{\omega}}$. Then $\overline{\sigma} \in \overline{\mathcal{A}}(\overline{\omega}, \overline{\tau})$, for otherwise we could pick $h \in G_{\overline{\tau}}$ such that $\overline{\omega}h \neq \overline{\omega}$ and depress h outside $\overline{\mathcal{A}}(\overline{\omega}, \overline{\tau})$ to obtain an element of $G_{\overline{\sigma}}$ which moves $\overline{\omega}$, contradicting the fact that $\overline{\sigma}F\overline{\omega}$. Since $\overline{\sigma} \in \overline{\mathcal{A}}(\overline{\omega}, \overline{\tau}), \overline{\mathcal{A}}(\overline{\sigma}, \overline{\tau}) = \operatorname{Conv}(\overline{\sigma}G_{\overline{\tau}}) = \operatorname{Conv}(\overline{\omega}G_{\overline{\tau}}) = \overline{\mathcal{A}}(\overline{\omega}, \overline{\tau})$. Hence we may write $\overline{\mathcal{A}}(\overline{\omega}, \overline{\tau})$ simply as $\overline{\mathcal{A}}(\overline{\tau})$; and $\{\overline{\mathcal{A}}(\overline{\tau}) \mid G_{\overline{\tau}} \in \mathscr{S}\}$ forms a tower under inclusion. Let $\overline{\delta}(\overline{\tau}) = \sup \overline{\mathcal{A}}(\overline{\tau})$ and let $\overline{\xi} = \inf\{\overline{\delta}(\overline{\tau}) \mid G_{\overline{\tau}} \in \mathscr{S}\}$. Since Cfixes each $\overline{\tau}$ such that $G_{\overline{\tau}} \in \mathscr{S}$, it fixes each $\overline{\mathcal{A}}(\overline{\tau})$ (by Lemma 10) and thus each $\overline{\delta}(\overline{\tau})$, so it fixes $\overline{\xi}$. Conversely, since $\overline{\sigma} \in \overline{\mathcal{A}}(\overline{\tau})$ whenever $G_{\overline{\sigma}} \subset G_{\overline{\tau}}$, we have $\overline{\xi} \in \overline{\mathcal{A}}(\overline{\tau})$. Since also $\overline{\mathcal{A}}(\overline{\tau})$ is an o-block of $G, G_{\overline{\xi}}$ fixes each $\overline{\mathcal{A}}(\overline{\gamma})$. Hence by Lemma 10, $G_{\overline{\xi}}$ fixes each $\overline{\tau}$. Therefore $G_{\overline{\xi}} = \{g \in G \mid \overline{\omega}g = \overline{\omega}$ for all $\overline{\omega} \in FxC\} = C$. This concludes the proof.

Theorem 11 fails when completeness is substituted for depressibility, even under the additional assumption that G is transitive. However, examples of this phenomenon are too complicated to be presented here. The author is presently trying to find additional hypotheses under which the modified theorem will hold.

By Zorn's lemma, every closed prime subgroup of G contains a minimal closed prime subgroup of G, which is characterized by

COROLLARY 12. $G_{\overline{\omega}}$ is a minimal closed prime subgroup of G if and only if for all $\overline{\tau} \in \overline{\Omega}$, $\overline{\omega}M\overline{\tau}$ implies $\overline{\tau}M\overline{\omega}$.

We define Min $(\overline{\Omega})$ to be $\{\overline{\omega} \in \overline{\Omega} \mid G_{\overline{\omega}} \text{ is a minimal closed prime sub$ $group of } G\}.$

THEOREM 13. Suppose that G is a depressible l-subgroup of $A(\Omega)$

and that for each $\alpha \in \Omega$, G_{α} has no fixed points in Ω except α . Then $\{G_{\overline{\omega}} \mid \overline{\omega} \in \operatorname{Min}(\overline{\Omega})\}$ are distinct and include all G_{α} 's, $\alpha \in \Omega$. Moreover, $\overline{\omega} \in \operatorname{Min}(\overline{\Omega})$ if and only if $\overline{\tau} M \overline{\omega}$ for every $\overline{\omega} \neq \overline{\tau} \in \overline{\Omega}$.

Proof. Suppose that $G_{\overline{\omega}} \subseteq G_{\alpha}$, with $\alpha \in \Omega$ and $\alpha \neq \overline{\omega} \in \overline{\Omega}$. If $\overline{\omega} \in \Omega$, then $\overline{\omega}F\alpha$ contradicts our hypothesis. Hence $\overline{\omega} \in \overline{\Omega} \setminus \Omega$, so we may pick $\beta \in \Omega$ lying strictly between α and $\overline{\omega}$. Then by hypothesis, we may pick $h \in G_{\beta}$ such that $\alpha h \neq \alpha$. By depressing h outside the interval of support $I(h, \alpha)$, we obtain an element of G which moves α , but fixes $\overline{\omega}$; contradicting the assumption that $G_{\overline{\omega}} \subseteq G_{\alpha}$. This shows that G_{α} is minimal, and moreover that $G_{\alpha} \neq G_{\overline{\omega}}$ for $\alpha \neq \overline{\omega} \in \overline{\Omega}$.

Now let $\bar{\omega} < \bar{\tau}$, with $\bar{\omega}, \bar{\tau} \in \text{Min}(\bar{\Omega})$, and suppose that $G_{\bar{\omega}} = G_{\bar{\tau}}$. Pick $\alpha \in \Omega$ such that $\bar{\omega} < \alpha \leq \bar{\tau}$. Since $G_{\bar{\omega}}$ is minimal, $G_{\alpha} \not\subset G_{\bar{\omega}}$, and by the previous paragraph, $G_{\alpha} \neq G_{\bar{\omega}}$. Hence we may pick $k \in G_{\alpha}$ such that $\bar{\omega}k \neq \bar{\omega}$. By depressing k outside $I(k, \bar{\omega})$, we obtain an element of G which moves $\bar{\omega}$, but fixes $\bar{\tau}$; contradicting the assumption that $G_{\bar{\omega}} = G_{\bar{\tau}}$. Hence $\{G_{\bar{\omega}} \mid \bar{\omega} \in \text{Min}(\bar{\Omega})\}$ are distinct. The last part of the theorem now follows from Corollary 12.

5. The relation between the *l*-group G and the chain Ω . Suppose now that the hypotheses of the last theorem are satisfied. We wish to determine how the chain $Min(\overline{\Omega})$ is reflected in the *l*-group structure of G. The theorem gives us a one-to-one correspondence $\overline{\omega} \leftrightarrow G_{\overline{\omega}}$ between $Min(\overline{\Omega})$ and the collection of minimal closed prime subgroups of G. Ideally, we would like to use the *l*-group structure of G to order these minimal closed prime subgroups so as to make this correspondence be an *o*-isomorphism (preserve order both ways). In general, this program will only partially succeed.

 (G, Ω) and (H, Σ) are said to be isomorphic as o-permutation groups if there is an o-isomorphism θ from Ω onto Σ and a map ψ from Gonto H such that $(\alpha g)\theta = (\alpha\theta)(g\psi)$ for all $\alpha \in \Omega, g \in G$. (It then follows that ψ is an *l*-isomorphism.) By the characteristic chains of (G, Ω) we shall mean the orbits $\bar{\omega}G$ of G (no convexification this time), for which $\bar{\omega} \in \operatorname{Min}(\bar{\Omega})$. Of course, if $\bar{\omega} \in \operatorname{Min}(\bar{\Omega})$, then $\bar{\omega}G \subseteq \operatorname{Min}(\bar{\Omega})$ since $G_{\bar{\omega}g} = g^{-1}G_{\bar{\omega}}g$. The characteristic chains of (G, Ω) partition $\operatorname{Min}(\bar{\Omega})$, and a subcollection of them partitions Ω . In particular, $(\operatorname{Min}(\bar{\Omega}))G =$ $\operatorname{Min}(\bar{\Omega})$, and G is faithful on $\operatorname{Min}(\bar{\Omega})$ since $\Omega \subseteq \operatorname{Min}(\bar{\Omega})$. The closure of a characteristic chain will refer to the order topology of $\operatorname{Min}(\bar{\Omega})$; or equivalently, since $\operatorname{Min}(\bar{\Omega})$ is dense in $\bar{\Omega}$, to the order topology of $\bar{\Omega}$. The closure of any one characteristic chain is of course a union of characteristic chains.

THEOREM 14. Suppose that (G, Ω) and (H, Σ) both satisfy the hypotheses of Theorem 13, and that ψ is an l-group isomorphism

from G onto H. Then there is an o-isomorphism θ from Min $(\overline{\Omega})$ onto Min $(\overline{\Sigma})$ such that θ and ψ provide an o-permutation group isomorphism from (G, Ω) onto (H, Σ) . (Exception: The order of $\overline{\sigma}$ and $\overline{\tau}$ need not be preserved unless there is some characteristic chain of (G, Ω) whose closure contains them both.) θ preserves characteristic chains and closures of characteristic chains. With the above exception, Σ is o-isomorphic to the union $\overline{\Gamma}$ of a collection of characteristic chains of (G, Ω) , and (H, Σ) is isomorphic as an o-permutation group to $(G, \overline{\Gamma})$.

Proof. We wish to (almost) determine the order of Min $(\overline{\Omega})$ from the *l*-group structure of G. Since $G_{\overline{\omega}} (\overline{\omega} \in \text{Min} (\overline{\Omega}))$ is a prime subgroup of G, we can totally order the set $R(G_{\overline{\omega}})$ of cosets $G_{\overline{\omega}}g (g \in G)$ by defining $G_{\overline{\omega}}g \leq G_{\overline{\omega}}k$ if and only if there exist $s \in G_{\overline{\omega}}g$ and $t \in G_{\overline{\omega}}k$ such that $s \leq t$ [3, Lemma 4]. It is easily checked that the correspondence $G_{\overline{\omega}}g \leftrightarrow \overline{\omega}g$ is an *o*-isomorphism between $R(G_{\overline{\omega}})$ and the characteristic chain $\overline{\omega}G$. Moreover, $(G_{\overline{\omega}}g)k = G_{\overline{\omega}}(gk) \leftrightarrow \overline{\omega}(gk) = (\overline{\omega}g)k, k \in G$.

Now let \overline{A} be a characteristic chain of (G, Ω) and let $(\overline{A}_1, \overline{A}_2)$ be a Dedekind cut of \overline{A} such that \overline{A}_1 has no greatest element and \overline{A}_2 no least element. Let $\overline{\delta}_1$ be the sup in $\overline{\Omega}$ of \overline{A}_1 and $\overline{\delta}_2$ the inf in $\overline{\Omega}$ of \overline{A}_2 . Let $G' = \{g \in G \mid \overline{A}_1g = \overline{A}_1 \text{ and } \overline{A}_2g = \overline{A}_2\}$. Then $G_{\overline{\delta}_1} = G' = G_{\overline{\delta}_2}$. Hence G'is a minimal closed prime subgroup of G if and only if there exists $\overline{\tau} \in \operatorname{Min}(\overline{\Omega})$ lying in the closure of \overline{A} and satisfying $\overline{A}_1 < \overline{\tau} < \overline{A}_2$. (If there exists such a $\overline{\tau}$, then $\overline{\tau} = \overline{\delta}_1$ or $\overline{\delta}_2$, say $\overline{\delta}_1$; so that $G' = G_{\overline{\delta}_1} = G_{\overline{\tau}}$ is minimal.) Moreover, if G' is minimal, then since $G_{\overline{\delta}_1} = G_{\overline{\delta}_2}$, Theorem 13 guarantees that $\overline{\delta}_1 = \overline{\delta}_2$ and thus that $\overline{\tau}$ is unique. Thus the closure of \overline{A} in Min ($\overline{\Omega}$) is determined by the *l*-group structure of G. (Of course, neither the sup nor the inf in $\overline{\Omega}$ of a nontrivial \overline{A} can lie in Min ($\overline{\Omega}$).) Similar considerations apply to (H, Σ) . For $\overline{\omega} \in \operatorname{Min}(\overline{\Omega})$, define $\overline{\omega}\theta$ to be the point in Min ($\overline{\Sigma}$) stabilized by $G_{\overline{\omega}}\psi$. The theorem follows.

The limitations of the theorem are illustrated by the following example. Let Σ_{α} be an α -set, $\alpha = 0, 1, 2$. (An α -set is a chain Σ of cardinality \aleph_{α} in which for any two subsets $\Gamma < \Delta$ of cardinality less than \aleph_{α} , there exists $\sigma \in \Sigma$ such that $\Gamma < \sigma < \Delta$.) Let Π_1 be the ordinal sum $\Sigma_0 \oplus \Sigma_1 \oplus \Sigma_2$, and Π_2 the ordinal sum $\Sigma_0 \oplus \Sigma_2 \oplus \Sigma_1$. Let $\Omega_i, i = 1, 2$, be the lexicographic product $\overline{\Pi_i \times J}$, ordered from the right, where J denotes the integers. Each copy of each Σ_{α} is an o-block of each $A(\Omega_i)$, whence it is easily seen that $A(\Omega_1)$ and $A(\Omega_2)$ are isomorphic as l-groups, despite the fact that $Min(\overline{\Omega}_1)$ and $Min(\overline{\Omega}_2)$ are not o-isomorphic.

COROLLARY 15. If the l-group G has at least one transitive representation (K, Π) of the sort described in the theorem, then the exception in the theorem can be removed.

Proof. Since K is transitive, Π is a characteristic chain of (K, Π) . The closure of Π in Min $(\overline{\Pi})$ is the entire chain Min $(\overline{\Pi})$. The theorem yields genuine *o*-isomorphisms from Min $(\overline{\Pi})$ onto Min $(\overline{\Omega})$, and also from Min $(\overline{\Pi})$ onto Min $(\overline{\Sigma})$.

For the special case in which G and H are themselves transitive and have minimal nontrivial o-blocks, this corollary is precisely Holland's main result in [4, Th. 7], applied to the depressible case. (Holland's result does not assume depressibility and is proved without the use of closed subgroups.) Further exploration of the transitive case takes on rather a different flavor and will be pursued in a later paper.

In [3, Th. 2], Holland represents an arbitrary *l*-group *L* as an *l*-subgroup *G* of an appropriate $A(\Omega)$, where the chain Ω is partitioned into convex subsets Ω_i such that for each Ω_i , $\Omega_i G = \Omega_i$ and *G* is transitive on Ω_i . If one such representation (*G*, Ω) satisfies the hypotheses of Theorem 13 (equivalently, if for each Ω_i , the restriction (*G* | Ω_i, Ω_i) satisfies the hypotheses of Theorem 13), then up to *o*-isomorphism, every such representation satisfying those hypotheses involves the same chains Min ($\overline{\Omega}_i$).

COROLLARY 16. Suppose that (G, Ω) satisfies the hypotheses of Theorem 13. Then every l-automorphism ψ of G is induced by an o-permutation θ of Min $(\overline{\Omega})$, i.e., $g\psi = \theta^{-1}g\theta$ for all $g \in G$. Moreover, θ permutes the collection of characteristic chains of (G, Ω) and preserves their closures. θ is subject to the exception of Theorem 14. This exception can be removed under the additional assumption of Corollary 15, and then the present result states that every l-automorphism of G is induced by an inner automorphism of $A(\text{Min}(\overline{\Omega}))$.

Proof. (G, Ω) and $(G\psi, \Omega)$ satisfy the hypotheses of Theorem 14. Let θ be the o-permutation of Min $(\overline{\Omega})$ provided by that theorem. Then $(\overline{\omega}\theta)(g\psi) = (\overline{\omega}g)\theta$ for all $\overline{\omega} \in \text{Min}(\overline{\Omega}), g \in G$, i.e., $\theta g\psi = g\theta$, and thus $g\psi = \theta^{-1}g\theta$.

Lloyd [5, Th. 1.10] proved Corollary 16 for the case in which G is all of $A(\Omega)$ and is transitive, except that θ was an *o*-permutation of $\overline{\Omega}$, rather than of the smaller chain Min $(\overline{\Omega})$. Lloyd's proof did not make use of closed subgroups. In many specific cases (e.g., if Ω is Dedekind complete or is the chain of rational numbers), the last statement in the corollary permits one to deduce (following Lloyd) that every *l*-automorphism of $A(\Omega)$ is inner.

References

^{1.} G. Birkhoff, Lattice theory, 3rd ed., Amer. Math. Soc. Colloq. Publns., Vol. 25, Providence, R.I., 1967.

^{2.} R. D. Byrd and J. T. Lloyd, Closed subgroups and complete distributivity in lattice-

ordered groups, Math. Zeit. 101 (1967), 123-130.

3. C. Holland, The lattice-ordered group of automorphisms of an ordered set, Michigan Math. J. 10 (1963), 399-408.

4. _____, Transitive lattice-ordered permutation groups, Math. Zeit 87 (1965), 420-433.

5. J. T. Lloyd, Lattice-ordered groups and o-permutations groups, Dissertation, Tulane University, 1964.

6. _____, Complete distributivity in certain infinite permutation groups, Michigan Math. J. 14 (1967), 393-400.

7. S. H. McCleary, *Pointwise suprema of order-preserving permutations* (to appear in Illinois J. Math.)

Received June 19, 1969.

UNIVERSITY OF GEORGIA ATHENS, GEORGIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN Stanford University Stanford, California

RICHARD PIERCE University of Washington Seattle, Washington 98105 J. DUGUNDJI Department of Mathematics University of Southern California Los Angeles, California 90007

BASIL GORDON

University of California Los Angeles, California 90024

ASSOCIATE EDITORS

YOSHIDA
τu

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA	STANFORD UNIVERSITY
CALIFORNIA INSTITUTE OF TECHNOLOGY	UNIVERSITY OF TOKYO
UNIVERSITY OF CALIFORNIA	UNIVERSITY OF UTAH
MONTANA STATE UNIVERSITY	WASHINGTON STATE UNIVERSITY
UNIVERSITY OF NEVADA	UNIVERSITY OF WASHINGTON
NEW MEXICO STATE UNIVERSITY	* * *
OREGON STATE UNIVERSITY	AMERICAN MATHEMATICAL SOCIETY
UNIVERSITY OF OREGON	CHEVRON RESEARCH CORPORATION
OSAKA UNIVERSITY	TRW SYSTEMS
UNIVERSITY OF SOUTHERN CALIFORNIA	NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539–1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The Pacific Journal of Mathematics is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics Vol. 31, No. 3 BadMonth, 1969

George E. Andrews, On a calculus of partition functions	555
Silvio Aurora, A representation theorem for certain connected rings	563
Lawrence Wasson Baggett, A note on groups with finite dual spaces	569
Steven Barry Bank, On majorants for solutions of algebraic differential equations in	
regions of the complex plane	573
Klaus R. Bichteler, Locally compact topologies on a group and the corresponding	
continuous irreducible representations	583
Mario Borelli, Affine complements of divisors	595
Carlos Jorge Do Rego Borges, A study of absolute extensor spaces	609
Bruce Langworthy Chalmers, Subspace kernels and minimum problems in Hilbert spaces with kernel function	619
John Dauns Representation of Learoups and Ferrings	629
Spencer Ernest Dickson and Kent Ralph Fuller. Algebras for which every	02)
indecomposable right module is invariant in its injective envelope	655
Robert Fraser and Sam Bernard Nadler, Ir. Sequences of contractive maps and fixed	055
noints	659
Judith Lee Gersting A rate of growth criterion for universality of regressive	057
isols	669
Robert Fred Gordon Rings in which minimal left ideals are projective	679
Fred Gross Entire functions of several variables with algebraic derivatives at	077
certain algebraic points	603
W Charles (Wilbur) Holland Ir and Stephen H. McCleary, Wradth products of	075
ordered permutation groups	703
W I Kim The Schwarzian derivative and multivalence	717
Robert Hamor I a Grange Ir. $On (m - n)$ products of Boolean algebras	725
Charles D. Masiello. The average of a gauge	723
Stephen H. McClearsy. The closed prime subgroups of cartain ordered permutation	133
arouns	745
Richard Roy Miller, Gleason parts and Choquet boundary points in convolution	745
menard Roy Miner, Orcason parts and Choquer boundary points in convolution measure aloebras	755
Harold I. Peterson Ir. On dvadic subspaces	773
Derek I S Robinson Groups which are minimal with respect to normality being	115
intransitive	777
Ralph Edwin Showalter Partial differential equations of Sobolev-Galpern type	787
David Slepian The content of some extreme simplexes	795
Joseph J. Taylor. Noncommutative convolution measure algebras	800
B. S. Vaday, Contractions of functions and their Fourier series	827
Lindsay Nathan Childs and Frank Pimi DeMayor, Correction to: "On	027
automorphisms of separable algebras"	833
Moses Classer and Pichard Emanuel Katz, Correction to: "Function theoretic	055
degeneracy criteria for Riemannian manifolds"	834
Satish Shirali, Correction to: "On the Jordan structure of complex Banach	
*algebras"	834
Benjamin Rigler Halpern, Addendum to: "Fixed points for iterates"	834