

Pacific Journal of Mathematics

**PARTIAL DIFFERENTIAL EQUATIONS OF
SOBOLEV-GALPERN TYPE**

RALPH EDWIN SHOWALTER

PARTIAL DIFFERENTIAL EQUATIONS OF SOBOLEV-GALPERN TYPE

R. E. SHOWALTER

A mixed initial and boundary value problem is considered for a partial differential equation of the form $Mu_t(x, t) + Lu(x, t) = 0$, where M and L are elliptic differential operators of orders $2m$ and $2l$, respectively, with $m \leq l$. The existence and uniqueness of a strong solution of this equation in $H_0^k(G)$ is proved by semigroup methods.

We are concerned here with a mixed initial boundary value problem for the equation

$$(1) \quad Mu_t + Lu = 0$$

in which M and L are elliptic differential operators. Equations of this type have been studied using various methods in [2, 3, 4, 6, 7, 10, 11, 13, 14, 15, 17, 18]. We will make use of the L^2 -estimates and related results on elliptic operators to obtain a generalized solution to this problem similar to that obtained for the parabolic equation

$$u_t + Lu = 0$$

as in [7].

Let G be a bounded open domain in R^n whose boundary ∂G is an $(n - 1)$ -dimensional manifold with G lying on one side of ∂G . By $H^k(G) \equiv H^k$ we mean the Hilbert space (of equivalence classes) of functions in $L^2(G)$ whose distributional derivatives through order k belong to $L^2(G)$ with the inner product and norm given, respectively, by

$$(f, g)_k = \sum \left\{ \int_G D^\alpha f \overline{D^\alpha g} dx : |\alpha| \leq k \right\}$$

and

$$\|f\|_k = \sqrt{(f, f)_k}.$$

$H_0^k \equiv H_0^k(G)$ will denote the closure in H^k of $C_0^\infty(G)$, the space of infinitely differentiable functions with compact support in G .

The operators are of the form

$$M = \sum \{ (-1)^{|\rho|} D^\rho m^{\rho\sigma}(x) D^\sigma : |\rho|, |\sigma| \leq m \}$$

and

$$L = \sum \{ (-1)^{|\rho|} D^\rho l^{\rho\sigma}(x) D^\sigma : |\rho|, |\sigma| \leq l \},$$

and they are uniformly strongly elliptic in G . We shall investigate the existence and uniqueness of solutions to (1) which coincide with the initial function u_0 in H_0^l where $t = 0$ and vanish on ∂G together with all derivatives of order less than or equal to $l - 1$.

If the order of M is as high as that of L ($2m \geq 2l$), then this problem can be handled as in [10] by forming the exponential of the bounded extension of $M^{-1}L$ on H_0^m and thus obtaining a group of operators on H_0^m and a corresponding solution for all t in \mathbf{R} . The case we shall consider is that of $m \leq l$, and this will include the parabolic equation as a special case. We obtain a semi-group of operators on H_0^m and, hence, a solution for all $t \geq 0$.

2. In this section we shall formulate the problem. Assume temporarily the following.

P'_1 : The coefficients $m^{\rho\sigma}$ in M belong to $H^{|\rho|}$, and $D^\sigma m^{\rho\sigma}$ is in $L^\infty(G)$ whenever $|\rho| \leq m$. A similar statement is true for the coefficients in L . From P'_1 it follows that the sesqui-linear forms defined on $C_0^\infty(G)$ by

$$B_M(\varphi, \psi) = \sum \{(m^{\rho\sigma} D^\sigma \varphi, D^\rho \psi)_0 : |\rho|, |\sigma| \leq m\}$$

and

$$B_L(\varphi, \psi) = \sum \{(l^{\rho\sigma} D^\sigma \varphi, D^\rho \psi)_0 : |\rho|, |\sigma| \leq l\}$$

satisfy the identities

$$(2) \quad B_M(\varphi, \psi) = (M\varphi, \psi)_0$$

and

$$(2') \quad B_L(\varphi, \psi) = (L\varphi, \psi)_0$$

for all φ, ψ in $C_0^\infty(G)$. Since P'_1 implies that

$$K_m = \sup \{\|m^{\rho\sigma}\|_\infty : |\rho|, |\sigma| \leq m\}$$

and

$$K_l = \sup \{\|l^{\rho\sigma}\|_\infty : |\rho|, |\sigma| \leq l\}$$

are finite, we see that

$$|B_M(\varphi, \psi)| \leq K_m \|\varphi\|_m \|\psi\|_m$$

and

$$|B_L(\varphi, \psi)| \leq K_l \|\varphi\|_l \|\psi\|_l$$

for all φ, ψ in $C_0^\infty(G)$. Hence these sesqui-linear forms may be extended by continuity to all of H_0^m and H_0^l , respectively.

The final properties which we shall assume are the following. For any φ, ψ in $C^\infty(G)$ we have

$$P_2: \operatorname{Re} B_M(\varphi, \varphi) \geq k_m \|\varphi\|_m^2, k_m > 0, \\ \operatorname{Re} B_L(\varphi, \varphi) \geq k_l \|\varphi\|_l^2, k_l > 0,$$

and

$$P_3: |B_M(\varphi, \psi)|^2 \leq (\operatorname{Re} B_M(\varphi, \varphi))(\operatorname{Re} B_M(\psi, \psi)).$$

These inequalities are valid for the respective extensions to H_0^m and H_0^l . The assumptions of P_2 are inequalities of the Garding type which imply that M and L are uniformly strongly elliptic. Only the first of these is essential in applications, for the usual change of dependent variable $u = ve^{2t}$ changes our equation to one with L replaced by $L + \lambda M$, and the Garding inequality is true for $B_{L+\lambda M}$ if λ is sufficiently large and if the coefficients $l^{\rho\sigma}(x), |\rho| = |\sigma| = l$ are uniformly continuous in G . See [3, 8] for sufficient conditions that P_2 be true.

The assumption P_3 is a Cauchy-Schwarz inequality for the form B_M . In view of the positivity of B_M , a necessary and sufficient condition for P_3 is that M be symmetric, that is, $m^{\rho\sigma} = \overline{m^{\sigma\rho}}$ for all ρ, σ . Such is the case for the examples

- (i) $ku_t - \Delta u = 0$ ($m = 0$) and
- (ii) $-\gamma \Delta u_t + ku_t - \Delta u = 0$,

where Δ is the Laplacian and γ and k are positive. Example (i) is a parabolic equation, and examples like (ii) appear in various problems of fluid mechanics and soil mechanics, where a solution is sought which satisfies an initial condition $u(x, 0) = u_0(x)$ on G and the Dirichlet condition $u(x, t) = 0$ on the boundary of G . See [1, 11, 12, 13].

We shall not need the full strength of P'_1 so we replace it with the following weaker assumption.

P_1 : The coefficients $m^{\rho\sigma}$ and $l^{\rho\sigma}$ belong to $L^\infty(G)$ for all ρ, σ .

We shall proceed under the assumptions P_1, P_2 and P_3 and remark that P'_1 is needed only when we wish to interpret our weak solutions by means of (2) and (2').

Under the hypotheses above there is by the theorem of Lax and Milgram [7] a closed linear operator M_0 with domain D_m dense in H_0^m and range equal to $H^0 = L^2(G)$ such that

$$(3) \quad B_M(\varphi, \psi) = (M_0\varphi, \psi)_0$$

whenever φ belongs to D_m and ψ to H_0^m . Furthermore, M_0^{-1} is a bounded operator from H^0 into H_0^m . Similarly, there is a closed linear operator L_0 with domain D_l dense in H_0^l and range equal to H^0 with

$$(3') \quad B_L(\varphi, \psi) = (L_0\varphi, \psi)_0$$

whenever φ belongs to D_l and ψ to H_0^l . Also, L_0^{-1} is bounded from H^0 into H_0^l .

Consider the bijection $A = -M_0^{-1}L_0$ from D_l onto D_m . For any φ in D_m we have

$$\begin{aligned} k_l \|A^{-1}\varphi\|_l^2 &= k_l \|L_0^{-1}M_0\varphi\|_l^2 \\ &\leq \operatorname{Re} B_L(L_0^{-1}M_0\varphi, L_0^{-1}M_0\varphi) = \operatorname{Re} \langle M_0\varphi, L_0^{-1}M_0\varphi \rangle_0 \\ &= \operatorname{Re} B_M(\varphi, L_0^{-1}M_0\varphi) \leq K_m \|\varphi\|_m \|A^{-1}\varphi\|_m \\ &\leq K_m \|\varphi\|_m \|A^{-1}\varphi\|_l \end{aligned}$$

which yields

$$(4) \quad \|A^{-1}\varphi\|_l \leq (K_m/k_l) \|\varphi\|_m$$

for all φ in D_m . But D_m is dense in H_0^m so A^{-1} has a unique extension by continuity from H_0^m onto the set $D = A^{-1}(H_0^m)$ in H_0^l , the domain of the closed extension of A . The continuity of the injection of H_0^l into H_0^m implies that A^{-1} is a bounded operator on H_0^m , and this is the space in which we formulate the Generalized Problem:

For a given initial function u_0 in D , find a differentiable map $u(t)$ of R^+ into H_0^m for which $u(t)$ belongs to H_0^l for all $t \geq 0$, $u(0) = u_0$, and

$$(5) \quad B_M(u'(t), \varphi) + B_L(u(t), \varphi) = 0$$

for all φ in $C_0^\infty(G)$ and $t \geq 0$.

Sufficient conditions for a solution of this generalized problem to be a classical solution will be discussed in [9].

3. The objective of this section is to prove the following results.

THEOREM. *There exists a unique solution of the generalized problem. If $u(t)$ is in D_l then $u'(t)$ is in D_m and*

$$(6) \quad M_0 u'(t) + L_0 u(t) = 0$$

in H^0 . The mapping of u_0 to $u(t)$ is continuous from H_0^m into itself for each $t \geq 0$ and furthermore satisfies

$$(7) \quad \|u(t)\|_m \leq \sqrt{K_m/k_m} \|u_0\|_m \exp(-k_l t/K_m).$$

We first show that the operator A is the infinitesimal generator of a semi-group of bounded operators on H_0^m ; this semi-group will provide a means of constructing a solution to the problem. From the assumptions on B_M , it follows that the function defined by

$$|\varphi|_M = \sqrt{\operatorname{Re} B_M(\varphi, \varphi)}$$

is a norm on H_0^m that is equivalent to the norm $\|\cdot\|_m$. In the following we shall use $|\cdot|_M$ as the norm on H_0^m , noting further that

$$(8) \quad k_m^{1/2} \|\varphi\|_m \leq \|\varphi\|_M \leq K_m^{1/2} \|\varphi\|_m$$

for φ in H_0^m .

To obtain the necessary estimates we let λ be a nonnegative number and consider the operator $\lambda M_0 + L_0 = N$ from the domain $D_m \cap D_l$ into H^0 . We can define a sesqui-linear form on $D_m \cap D_l$ by

$$B_N(\varphi, \psi) = ((\lambda M_0 + L_0)\varphi, \psi)_0 = \lambda B_M(\varphi, \psi) + B_L(\varphi, \psi)$$

and then note that B_N is bounded as well as positive-definite with respect to the norm of H_0^l . We extend B_N by continuity to all of H_0^l , and then by the theorem of Lax and Milgram there is a closed linear operator N_0 from a domain D_n in H_0^l onto H^0 for which

$$B_N(\varphi, \psi) = (N_0\varphi, \psi)_0$$

whenever φ is in D_n and ψ in H_0^l . Clearly N_0 is an extension of N whose domain is $D_m \cap D_l$.

For all φ in D_n we have

$$\begin{aligned} \operatorname{Re} (N_0\varphi, \varphi)_0 &= \lambda \operatorname{Re} B_M(\varphi, \varphi) + \operatorname{Re} B_L(\varphi, \varphi) \\ &\geq (\lambda + k_l/K_m) \operatorname{Re} B_M(\varphi, \varphi) \\ &= (\lambda + k_l/K_m) \|\varphi\|_M^2. \end{aligned}$$

Thus, for any ψ in D_m we see that $N_0^{-1}M_0\psi$ belongs to D_n and from above

$$\begin{aligned} (\lambda + k_l/K_m) \|N_0^{-1}M_0\psi\|_M^2 &\leq \operatorname{Re} (M_0\psi, N_0^{-1}M_0\psi)_0 \\ &= \operatorname{Re} B_M(\psi, N_0^{-1}M_0\psi) \leq \|\psi\|_M \|(N_0^{-1}M_0\psi)\|_M \end{aligned}$$

by P_3 , so we have obtained the estimate

$$\|N_0^{-1}M_0\psi\|_M \leq (\lambda + k_l/K_m)^{-1} \|\psi\|_M$$

for all ψ in D_m .

Letting φ be an element of $D_l \cap D_m$ we see

$$\begin{aligned} (N_0^{-1}M_0)(\lambda + M_0^{-1}L_0)\varphi &= N_0^{-1}(\lambda M_0\varphi + L_0\varphi) \\ &= N_0^{-1} \cdot N\varphi = \varphi, \end{aligned}$$

so $\lambda + M_0^{-1}L_0$ is injective and satisfies

$$(\lambda + M_0^{-1}L_0)^{-1} = N_0^{-1}M_0$$

on $D_m \cap D_l$. Combining this with the estimate above we see that

$$\|(\lambda + M_0^{-1}L_0)^{-1}\psi\|_M \leq (\lambda + k_l/K_m)^{-1} \|\psi\|_M$$

for all ψ in $D_l \cap D_m$. It follows by continuity that $\lambda - A$ is invertible on H_0^m and satisfies the estimate

$$\|(\lambda - A)^{-1}\psi\|_M \leq (\lambda + k_l/K_m)^{-1} \|\psi\|_M.$$

By the theorem of Hille and Yoshida [5, 16] on the characterization of the infinitesimal generators of semi-groups of class C_0 we have the following results: there exists a unique family of bounded operators $\{S(t): t \geq 0\}$ on H_0^m for which

- (i) $S(t_1 + t_2) = S(t_1)S(t_2)$,
- (ii) $S(t)x$ is strongly continuous for each x in H_0^m ,
- (iii) $S(0) = I$ and $\|S(t)\|_M \leq \exp(-k_1 t/K_m)$ for all $t \geq 0$,
- (iv) $\lim_{h \rightarrow 0} h^{-1}(S(h) - I)x_0 = Ax_0$ for each x_0 in D , and
- (v) $S(t)$ commutes with $(\lambda - A)^{-1}$ for all $\lambda \geq 0$.

The statement (v) implies in particular that D is invariant under each $S(t)$.

Having been given the initial function u_0 in D , we define

$$u(t) = S(t)u_0, t \geq 0$$

and show that $u(t)$ is a solution of the generalized problem. Clearly we see $u(t)$ belongs to H_0^m and $u(0) = u_0$. Furthermore, since $S(t)$ leaves D invariant and u_0 is in D , it follows that $u(t)$ belongs to D and thus to H_0^l . The function $u(t)$ is differentiable with

$$(9) \quad u'(t) = Au(t)$$

for all $t \geq 0$ by (i) and (iv), and hence $u'(t)$ is in H_0^m .

We shall verify that $u(t)$ satisfies the equation (5). Since D_m is dense in H_0^m there is a sequence $\{\varphi_n\}$ in D_m for which $\|\varphi_n - u'(t)\|_m \rightarrow 0$ as $n \rightarrow \infty$. Now $\{\varphi_n\}$ is a Cauchy sequence in H_0^m and it follows by (4) that $\psi_n = A^{-1}\varphi_n$ is a Cauchy sequence in the complete space H_0^l , so there is a ψ in H_0^l such that $\|\psi_n - \psi\|_l \rightarrow 0$ as $n \rightarrow \infty$. Since A^{-1} is continuous we have $\psi = u(t)$. Each ψ_n belongs to D_l , since φ_n is in D_m , and furthermore $M_0\varphi_n + L_0\psi_n = 0$. Now for each φ in $C_0^\infty(G)$ we have by the continuity of B_M and B_L

$$\begin{aligned} & B_M(u'(t), \varphi) + B_L(u(t), \varphi) \\ &= \lim_{n \rightarrow \infty} B_M(\varphi_n, \varphi) + \lim_{n \rightarrow \infty} B_L(\psi_n, \varphi) \\ &= \lim_{n \rightarrow \infty} \{B_M(\varphi_n, \varphi) + B_L(\psi_n, \varphi)\} = \lim_{n \rightarrow \infty} \{(M_0\varphi_n, \varphi)_0 + (L_0\psi_n, \varphi)_0\} \equiv 0, \end{aligned}$$

so the generalized problem does have a solution.

If $u(t)$ is in D_l then by (9) $u'(t)$ is in D_m . It follows from (5) that for every φ in $C_0^\infty(G)$

$$(M_0u'(t) + L_0u(t), \varphi)_0 = 0,$$

and this implies (6). The estimate (7) is a consequence of (iii) and (8).

To show that the generalized problem has at most one solution, we let $u(t)$ be a solution of the problem with $u_0 = 0$. By linearity it suffices to show that $u(t) \equiv 0$. The differentiability of $u(t)$ in H_0^m

implies that the real valued function

$$\alpha(t) = \operatorname{Re} B_M(u(t), u(t))$$

is differentiable and

$$\alpha'(t) = 2 \operatorname{Re} B_M(u'(t), u(t)) .$$

Since (5) is true also for all φ in H_0^1 by continuity, we have from P_2

$$\alpha'(t) = -2 \operatorname{Re} B_L(u(t), u(t)) \leq 0 .$$

But $\alpha(0) = \operatorname{Re} B_M(u(0), u(0)) = 0$, so $\alpha(t) = 0$ for all $t \geq 0$. From P_2 it follows that $u(t) = 0$ for $t \geq 0$.

REFERENCES

1. G. Barenblatt, I. Zheltov, and I. Kochiva, *Basic concepts in the theory of seepage of homogeneous liquids in fissured rocks*, J. Appl. Math. **24** (1960), 1286-1303.
2. R. Carroll, *Partial differential equations* (Harper and Row, to appear)
3. A. Friedman, *Partial differential equations of parabolic type*, Prentice-Hall, 1964.
4. C. A. Galpern, *Doklady Akad. Nauk. SSSR* **119** (1958), 640-643.
5. E. Hille and R. S. Phillips, *Functional analysis and semi-groups*, A.M.S. Colloquium Publications, vol. 31, New York, 1957.
6. A. G. Kostachenko and G. I. Eskin, *Cauchy problem for an equation of Sobolev-Galpern type*, Trudy Moscow Math. Obshch. **10** (1961), 273-284.
7. P. D. Lax and A. Milgram, *Parabolic equations*, Contributions to the Theory of Partial Differential Equations, Ann. Math. Studies, no. 33 Princeton University Press, 1954, 167-190.
8. J. L. Lions, *Lectures on elliptic partial differential equations*, Tata Institute, Bombay, 1957.
9. R. E. Showalter, *Local regularity of solutions of pseudo-parabolic partial differential equations* (to appear).
10. R. E. Showalter and T. W. Ting, *Pseudo-parabolic partial differential equations*, (to appear, SIAM Journal)
11. S. Sobolev *Izv. Akad. Nauk. SSSR Ser. Mat.* **18** (1954), 3.
12. D. W. Taylor, *Research on consolidation of clays*, Massachusetts Institute of Technology, 1942.
13. T. W. Ting, *Certain non-steady flows of second order fluids*, Arch. Rat. Mech. Anal. **14** (1963), 1-26.
14. ———, *Parabolic and pseudo-parabolic partial differential equations* (to appear, J. Math. Soc. Japan).
15. T. W. Ting and R. E. Showalter, *Asymptotic behavior of solutions of pseudo-parabolic partial differential equations* (to appear, Proc. Amer. Math. Soc.)
16. K. Yoshida, *On the differentiability and representation of one-parameter semi-groups of linear operators*, J. Math. Soc. Japan **1** (1948), 15-21.
17. T. Zelenyak, *The behavior for $t \rightarrow \infty$ of solutions of a problem of S. L. Sobolev*, Soviet Math. **2** (1961), 956-958.
18. ———, *A problem of Sobolev*, Soviet Math. **3** (1962), 1756-1759.

Received February 14, 1969.

UNIVERSITY OF TEXAS
AUSTIN, TEXAS

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

BASIL GORDON
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics

Vol. 31, No. 3

BadMonth, 1969

George E. Andrews, <i>On a calculus of partition functions</i>	555
Silvio Aurora, <i>A representation theorem for certain connected rings</i>	563
Lawrence Wasson Baggett, <i>A note on groups with finite dual spaces</i>	569
Steven Barry Bank, <i>On majorants for solutions of algebraic differential equations in regions of the complex plane</i>	573
Klaus R. Bichteler, <i>Locally compact topologies on a group and the corresponding continuous irreducible representations</i>	583
Mario Borelli, <i>Affine complements of divisors</i>	595
Carlos Jorge Do Rego Borges, <i>A study of absolute extensor spaces</i>	609
Bruce Langworthy Chalmers, <i>Subspace kernels and minimum problems in Hilbert spaces with kernel function</i>	619
John Dauns, <i>Representation of L-groups and F-rings</i>	629
Spencer Ernest Dickson and Kent Ralph Fuller, <i>Algebras for which every indecomposable right module is invariant in its injective envelope</i>	655
Robert Fraser and Sam Bernard Nadler, Jr., <i>Sequences of contractive maps and fixed points</i>	659
Judith Lee Gersting, <i>A rate of growth criterion for universality of regressive isols</i>	669
Robert Fred Gordon, <i>Rings in which minimal left ideals are projective</i>	679
Fred Gross, <i>Entire functions of several variables with algebraic derivatives at certain algebraic points</i>	693
W. Charles (Wilbur) Holland Jr. and Stephen H. McCleary, <i>Wreath products of ordered permutation groups</i>	703
W. J. Kim, <i>The Schwarzian derivative and multivalence</i>	717
Robert Hamor La Grange, Jr., <i>On $(m - n)$ products of Boolean algebras</i>	725
Charles D. Masiello, <i>The average of a gauge</i>	733
Stephen H. McCleary, <i>The closed prime subgroups of certain ordered permutation groups</i>	745
Richard Roy Miller, <i>Gleason parts and Choquet boundary points in convolution measure algebras</i>	755
Harold L. Peterson, Jr., <i>On dyadic subspaces</i>	773
Derek J. S. Robinson, <i>Groups which are minimal with respect to normality being intransitive</i>	777
Ralph Edwin Showalter, <i>Partial differential equations of Sobolev-Galpern type</i>	787
David Slepian, <i>The content of some extreme simplexes</i>	795
Joseph L. Taylor, <i>Noncommutative convolution measure algebras</i>	809
B. S. Yadav, <i>Contractions of functions and their Fourier series</i>	827
Lindsay Nathan Childs and Frank Rimi DeMeyer, <i>Correction to: "On automorphisms of separable algebras"</i>	833
Moses Glasner and Richard Emanuel Katz, <i>Correction to: "Function-theoretic degeneracy criteria for Riemannian manifolds"</i>	834
Satish Shirali, <i>Correction to: "On the Jordan structure of complex Banach *algebras"</i>	834
Benjamin Rigler Halpern, <i>Addendum to: "Fixed points for iterates"</i>	834