

# Pacific Journal of Mathematics

**CONTRACTIONS OF FUNCTIONS AND THEIR FOURIER  
SERIES**

B. S. YADAV

## CONTRACTIONS OF FUNCTIONS AND THEIR FOURIER SERIES

To Professor U. N. Singh on his 49<sup>th</sup> birthday

B. S. YADAV

The object of the present paper is to define a new type of contraction, called 'Shrivel', of a function and to prove a theorem on the absolute convergence of its Fourier series. Our theorem is similar to a theorem of M. Kinukawa, but as it is shown in the end the two results are essentially different. The original results in this direction are due to A. Beurling and R. P. Boas.

According to Beurling [1] a function  $g$  is said to be a contraction of function  $f$  if  $|g(x) - g(y)| \leq A |f(x) - f(y)|$ , for all  $x, y$ , where  $A$  is an absolute constant. We shall assume throughout that the functions  $f$  and  $g$  are each  $L -$  integrable in  $(-\pi, \pi)$  and periodic with the period  $2\pi$ . We shall further let

$$f(x) \sim \frac{1}{2}a_0 + \sum_{k=1}^{\infty} (a_k \cos kx + b_k \sin kx)$$

and

$$g(x) \sim \frac{1}{2}c_0 + \sum_{k=1}^{\infty} (c_k \cos kx + d_k \sin kx).$$

Kinukawa [3] has proved the following

**THEOREM 1.** *Let  $f$  and  $g$  be each continuous and  $g$  be a contraction of  $f$ , or more generally let  $f, g \in L_2$  and*

$$\int_{-\pi}^{\pi} |g(x+h) - g(x)|^2 dx \leq \int_{-\pi}^{\pi} |f(x+h) - f(x)|^2 dx, \text{ for all } h.$$

If

$$(1) \quad \sum_{n=1}^{\infty} n^{-3\alpha/2} \left( \sum_{k=1}^n k^2 p_k^2 \right)^{\alpha/2} < \infty$$

and

$$(2) \quad \sum_{n=1}^{\infty} n^{-\alpha/2} \left( \sum_{k=n+1}^{\infty} p_k^2 \right)^{\alpha/2} < \infty,$$

where  $p_k^\alpha = |a_k|^\alpha + |b_k|^\alpha$  and  $0 < \alpha \leq 2$ , then

$$(3) \quad \sum_{k=1}^{\infty} (|c_k|^\alpha + |d_k|^\alpha) < \infty.$$

The case  $\alpha = 1$  in this theorem is the theorem of Boas [2], which in turn is a generalization of a theorem of Beurling [1] on the absolute convergence of Fourier series.

However, it has been shown by Leindler [4] and Sunouchi [5]<sup>1</sup> that the conditions (1) and (2) are equivalent. In fact, a more general result has been proved by R. Askey in his not yet published work which we shall need later and hence state here in the form of the following

LEMMA<sup>1</sup>. *Let  $s > 0$ ,  $0 < p < 1$ ,  $a_k > 0$ . There are constants  $K = K(p, r, s) > 0$  such that*

$$\sum_{n=1}^{\infty} n^r \left[ n^{-s} \sum_{k=1}^n k^s a_k \right]^p \geq K \sum_{n=1}^{\infty} n^r \left( \sum_{k=n}^{\infty} a_k \right)^p, \quad r > -1;$$

and

$$\sum_{n=1}^{\infty} n^r \left( \sum_{k=n}^{\infty} a_k \right)^p \geq K \sum_{k=1}^{\infty} n^r \left[ n^{-s} \sum_{k=1}^n k^s a_k \right]^p, \quad r < ps - 1.$$

The object of this paper is to prove Theorem 2 which is similar to Theorem 1; but as we shall see in the end the two theorems are independent of each other. Before we can state our result precisely, we need introduce a couple of notations and a definition.

We put

$$\Delta_t^m f(x) = \sum_{k=0}^m (-1)^k C_m^k f[x + (m - 2k)t]$$

and

$$L^{(m)}(h, x, f) = \frac{1}{h} \int_0^h \Delta_t^m f(x) dt.$$

DEFINITION. We shall call a function  $g$  a 'shrivel' of order  $m$  of a function  $f$  if

$$|L^{(m)}(h, x, g)| \leq A |L^{(m)}(h, x, f)|,$$

for all  $x$  and for all  $h > 0$ .

We shall prove the following

THEOREM 2. *If  $g$  is a shrivel of order  $m$  of  $f$ , or more generally, if*

---

<sup>1</sup> The author thanks the referee for pointing out to him these references and the unpublished work of R. Askey to show the equivalence of the conditions (1) and (2) which he missed in his original text of the paper.

$$(4) \quad \int_{-\pi}^{\pi} |L^{(m)}(h, x, g)|^2 dx \leq \int_{-\pi}^{\pi} |L^{(m)}(h, x, f)|^2 dx,$$

and if (2) holds, then the conclusion (3) remains valid.

Before we proceed for the proof, we mark that for  $0 < |h| \leq \pi$ ,

$$|L^{(m)}(h, x, f)| \leq \frac{A(m)}{|h|} \int_{-\pi}^{\pi} |f(t)| dt;$$

and hence  $L^{(m)}(h, x, f)$  is a bounded function of  $x$  for fixed  $h$  and  $f$ . [ $A(m)$  denotes here, as also in the sequel, a constant depending on  $m$  but not necessarily the same everywhere.] Therefore  $L^{(m)}(h, x, f) \in L_2$ . The same is true about  $L^{(m)}(h, x, g)$  also. Thus the condition ‘ $g$  is a shrivel of order  $m$  of  $f$ ’ does imply (4). Obviously, the converse is not true.

Again, it follows from the lemma that the condition (2) is equivalent to the condition

$$(5) \quad \sum_{n=1}^{\infty} n^{-(m+1/2)\alpha} \left( \sum_{k=1}^n k^{2m} p_k^2 \right)^{\alpha/2} < \infty,$$

and this is what we shall need in the proof of our theorem.

*Proof.* We can obtain by simple calculations that, for an even  $m$ ,

$$A_t^m g(x) \sim (-1)^{m/2} 2^m \sum_{k=1}^{\infty} (c_k \cos kx + d_k \sin kx) \sin^m kt;$$

and hence

$$L^{(m)}(h, x, g) \sim (-1)^{m/2} 2^m \sum_{k=1}^{\infty} (c_k \cos kx + d_k \sin kx) \left( \frac{1}{h} \int_0^h \sin^m kt dt \right).$$

Similarly for an odd  $m$ ,

$$L^{(m)}(h, x, g) \sim (-1)^{(m-1)/2} 2^m \sum_{k=1}^{\infty} (d_k \cos kx - c_k \sin kx) \left( \frac{1}{h} \int_0^h \sin^m kt dt \right).$$

Therefore by Parseval’s theorem

$$(6) \quad \begin{aligned} \sum_{k=1}^{\infty} q_k^2 \left( \frac{1}{h} \int_0^h \sin^m kt dt \right)^2 &= A(m) \int_{-\pi}^{\pi} |L^{(m)}(h, x, g)|^2 dx \\ &\leq A(m) \int_{-\pi}^{\pi} |L^{(m)}(h, x, f)|^2 dx \\ &= A(m) \sum_{k=1}^{\infty} p_k^2 \left( \frac{1}{h} \int_{-\pi}^{\pi} \sin^m kt dt \right)^2, \end{aligned}$$

where

$$q_k^2 = c_k^2 + d_k^2 .$$

Now taking  $h = \pi/2^{n+1}$  and observing that

(i)  $\sin kt > 0$ , and is an increasing function of  $t$  for  $\pi/2^{n+2} \leq t \leq \pi/2^{n+1}$ ;  $2^{n-1} < k \leq 2^n$ ,

(ii)  $\sin^2 k\pi/2^{n+2} > \sin^2 \pi/8$  for  $2^{n-1} < k \leq 2^n$ , we get

$$\begin{aligned}
 (7) \quad \sum_{k=1}^{\infty} q_k^2 & \left( \frac{2^{n+1}}{\pi} \int_0^{\pi/2^{n+1}} \sin^m kt dt \right)^2 \\
 & \geq \sum_{k=2^{n-1}+1}^{2^n} q_k^2 \left( \frac{2^{n+1}}{\pi} \int_0^{\pi/2^{n+1}} \sin^m kt dt \right)^2 \\
 & \geq \sum_{k=2^{n-1}+1}^{2^n} q_k^2 \left( \frac{2^{n+1}}{\pi} \int_{\pi/2^{n+2}}^{\pi/2^{n+1}} \sin^m kt dt \right)^2 \\
 & \geq \sum_{k=2^{n-1}+1}^{2^n} q_k^2 \left( \frac{2^{n+1}}{\pi} \cdot \frac{\pi}{2^{n+2}} \cdot \sin^m \frac{k\pi}{2^{n+1}} \right)^2 \\
 & \geq A(m) \sum_{k=2^{n-1}+1}^{2^n} q_k^2 .
 \end{aligned}$$

Also, by Hölder’s inequality

$$(8) \quad \sum_{k=2^{n-1}+1}^{2^n} q_k^\alpha \leq 2^{(n-1)(1-\alpha/2)} \left( \sum_{k=2^{n-1}+1}^{2^n} q_k^2 \right)^{\alpha/2} .$$

Therefore we obtain from (6), (7) and (8)

$$\begin{aligned}
 \sum_{n=2}^{\infty} q_n^\alpha & = \sum_{n=1}^{\infty} \sum_{k=2^{n-1}+1}^{2^n} q_k^\alpha \\
 & \leq A(m, \alpha) \sum_{n=1}^{\infty} 2^{n(1-\alpha/2)} \left[ \sum_{k=1}^{\infty} p_k^2 \left( \frac{2^{n+1}}{\pi} \int_0^{\pi/2^{n+1}} \sin^m kt dt \right)^2 \right]^{\alpha/2} \\
 & \leq A(m, \alpha) \left[ \sum_{n=1}^{\infty} 2^{n(1-\alpha/2)} \left( \sum_{k=1}^{2^n} p_k^2 \sin^{2m} \frac{k\pi}{2^{n+1}} \right)^{\alpha/2} \right. \\
 & \quad \left. + \sum_{n=1}^{\infty} 2^{n(1-\alpha/2)} \left( \sum_{k=2^{n+1}}^{\infty} p_k^2 \right)^{\alpha/2} \right] \\
 & \leq A(m, \alpha) \left[ \sum_{n=1}^{\infty} 2^{n(1-\alpha/2-m\alpha)} \left( \sum_{k=1}^{2^n} k^{2m} p_k^2 \right)^{\alpha/2} \right. \\
 & \quad \left. + \sum_{n=1}^{\infty} 2^{n(1-\alpha/2)} \left( \sum_{k=2^{n+1}}^{\infty} p_k^2 \right)^{\alpha/2} \right] .
 \end{aligned}$$

Now it is not difficult to see that the two series within the square brackets are convergent if and only if (5) and (2) hold respectively. See Szàsz [6], Lemma 2.2. Since

$$|c_k|^\alpha \leq q_k^\alpha , \quad |d_k|^\alpha \leq q_k^\alpha ,$$

(3) follows; and hence the proof.

To prove that the result of Theorem 2 is completely independent of Kinukawa's Theorem 1, we first see that  $g$  is a shrivel of order 1 of  $f$  if for all  $x$  and all  $h > 0$ ,

$$(9) \quad \left| \int_0^h \{g(x+t) - g(x-t)\} dt \right| \leq A \left| \int_0^h \{f(x+t) - f(x-t)\} dt \right| .$$

Now consider the  $2\pi$  - periodic function defined by

$$\begin{aligned} f(x) &= 0 && \text{for } -\pi < x < 0, \\ &= \sin 2x && \text{for } 0 \leq x \leq \pi . \end{aligned}$$

Also let

$$\begin{aligned} \phi(x) &= x && \text{for } x < 0 \\ &= x - x^2 && \text{for } x \geq 0 , \end{aligned}$$

and let  $g(x) = \phi(f(x))$  for all  $x$ . The function  $\phi$  is differentiable and has a bounded derivative on the interval  $[-1, 1]$ . Hence  $\phi$  belongs to the class Lip 1 on this interval, whence,  $g$  is a contraction of  $f$  in Beurling's sense. On the other hand,

$$\int_0^\pi \{f(t) - f(-t)\} dt = \int_0^\pi \sin 2t dt = 0 ,$$

while

$$\begin{aligned} \int_0^\pi \{g(t) - g(-t)\} dt &= \int_0^\pi g(t) dt \\ &= \int_0^{\pi/2} (\sin 2t - \sin^2 2t) 2t + \int_{\pi/2}^\pi \sin 2t dt \\ &= - \int_0^{\pi/2} \sin^2 2t dt \neq 0 , \end{aligned}$$

so that (9) is not satisfied for any  $A$  when  $x = 0$  and  $h = \pi$ .

Conversely, since (9) can be satisfied when  $f, g$  are merely integrable, (9) does not imply that  $g$  is a contraction of  $f$ .

Finally we remark that Theorem 2 has its usual dual.

#### REFERENCES

1. A. Beurling, *On the spectral synthesis of bounded functions*, Acta Math. **81** (1949), 225-238.
2. R. P. Boas, *Beurling's test for absolute convergence of Fourier series*, Bull. Amer. Math. Soc. **66** (1960), 24-27.
3. M. Kinukawa, *Contractions of Fourier coefficients and Fourier integrals*, J. Analyse Math. **8** (1960/1961), 377-406.

4. L. Leindler, *Über verschiedene Konevergenzarten trigonometrischer Reihen*, Acta Sci. Math. (Szeged) **25** (1964), 233-249.
5. G. Sunouchi, *On the convolution algebra of Beurling*, Tôhoku Math. J. (2) (1967), 303-310.
6. O. Szász, *Fourier series and mean moduli of continuity*, Trans. Amer. Math. Soc. **42** (1937), 366-395.

Received October 8, 1968.

SARDAR PATEL UNIVERSITY  
VALLABH VIDYANAGAR, GUJARAT, INDIA

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. ROYDEN  
Stanford University  
Stanford, California

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

RICHARD PIERCE  
University of Washington  
Seattle, Washington 98105

BASIL GORDON  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL WEAPONS CENTER

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.



# Pacific Journal of Mathematics

Vol. 31, No. 3

BadMonth, 1969

George E. Andrews, <i>On a calculus of partition functions</i> . . . . .	555
Silvio Aurora, <i>A representation theorem for certain connected rings</i> . . . . .	563
Lawrence Wasson Baggett, <i>A note on groups with finite dual spaces</i> . . . . .	569
Steven Barry Bank, <i>On majorants for solutions of algebraic differential equations in regions of the complex plane</i> . . . . .	573
Klaus R. Bichteler, <i>Locally compact topologies on a group and the corresponding continuous irreducible representations</i> . . . . .	583
Mario Borelli, <i>Affine complements of divisors</i> . . . . .	595
Carlos Jorge Do Rego Borges, <i>A study of absolute extensor spaces</i> . . . . .	609
Bruce Langworthy Chalmers, <i>Subspace kernels and minimum problems in Hilbert spaces with kernel function</i> . . . . .	619
John Dauns, <i>Representation of L-groups and F-rings</i> . . . . .	629
Spencer Ernest Dickson and Kent Ralph Fuller, <i>Algebras for which every indecomposable right module is invariant in its injective envelope</i> . . . . .	655
Robert Fraser and Sam Bernard Nadler, Jr., <i>Sequences of contractive maps and fixed points</i> . . . . .	659
Judith Lee Gersting, <i>A rate of growth criterion for universality of regressive isols</i> . . . . .	669
Robert Fred Gordon, <i>Rings in which minimal left ideals are projective</i> . . . . .	679
Fred Gross, <i>Entire functions of several variables with algebraic derivatives at certain algebraic points</i> . . . . .	693
W. Charles (Wilbur) Holland Jr. and Stephen H. McCleary, <i>Wreath products of ordered permutation groups</i> . . . . .	703
W. J. Kim, <i>The Schwarzian derivative and multivalence</i> . . . . .	717
Robert Hamor La Grange, Jr., <i>On <math>(m - n)</math> products of Boolean algebras</i> . . . . .	725
Charles D. Masiello, <i>The average of a gauge</i> . . . . .	733
Stephen H. McCleary, <i>The closed prime subgroups of certain ordered permutation groups</i> . . . . .	745
Richard Roy Miller, <i>Gleason parts and Choquet boundary points in convolution measure algebras</i> . . . . .	755
Harold L. Peterson, Jr., <i>On dyadic subspaces</i> . . . . .	773
Derek J. S. Robinson, <i>Groups which are minimal with respect to normality being intransitive</i> . . . . .	777
Ralph Edwin Showalter, <i>Partial differential equations of Sobolev-Galpern type</i> . . . . .	787
David Slepian, <i>The content of some extreme simplexes</i> . . . . .	795
Joseph L. Taylor, <i>Noncommutative convolution measure algebras</i> . . . . .	809
B. S. Yadav, <i>Contractions of functions and their Fourier series</i> . . . . .	827
Lindsay Nathan Childs and Frank Rimi DeMeyer, <i>Correction to: "On automorphisms of separable algebras"</i> . . . . .	833
Moses Glasner and Richard Emanuel Katz, <i>Correction to: "Function-theoretic degeneracy criteria for Riemannian manifolds"</i> . . . . .	834
Satish Shirali, <i>Correction to: "On the Jordan structure of complex Banach *algebras"</i> . . . . .	834
Benjamin Rigler Halpern, <i>Addendum to: "Fixed points for iterates"</i> . . . . .	834