

# Pacific Journal of Mathematics

## **COMPACT SOBOLEV IMBEDDINGS FOR UNBOUNDED DOMAINS**

ROBERT ALEXANDER ADAMS

## COMPACT SOBOLEV IMBEDDINGS FOR UNBOUNDED DOMAINS

ROBERT A. ADAMS

**A condition on an open set  $G \subset E_n$  which is both necessary and sufficient for the compactness of the (Sobolev) imbedding  $H_0^{m+1}(G) \rightarrow H_0^m(G)$  is not yet known. C. Clark has given a necessary condition (quasiboundedness) and a much stronger sufficient condition. We show here that (unless  $n = 1$ ) quasiboundedness is not sufficient, and answer in the negative a question raised by Clark on whether the imbedding can be compact if  $\partial G$  consists of isolated points. We also substantially weaken Clark's sufficient condition so as to include a wide class of domains with null exterior. The gap between necessary and sufficient conditions is thus considerably narrowed.**

Let  $G$  be an open set in Euclidean  $n$ -space,  $E_n$ . Let  $H_0^m(G)$  for each nonnegative integer  $m$  denote the Sobolev space obtained by completing with respect to the norm

$$\|u\|_{m,G} = \left\{ \sum_{|\alpha| \leq m} \int_G |D^\alpha u(x)|^2 dx \right\}^{1/2}$$

the space  $C_0^\infty(G)$  of all infinitely differentiable complex valued functions having compact support in  $G$ . Here, as usual,  $\alpha = (\alpha_1, \dots, \alpha_n)$  is an  $n$ -tuple of nonnegative integers;  $|\alpha| = \alpha_1 + \dots + \alpha_n$ , and  $D^\alpha = D_1^{\alpha_1} \dots D_n^{\alpha_n}$  where  $D_j = \partial/\partial x_j$ ,  $j = 1, \dots, n$ .

We shall say that  $G$  has the Rellich property if for each integer  $m \geq 0$  the imbedding mapping  $H_0^{m+1}(G) \rightarrow H_0^m(G)$  is compact. It is well known that any bounded  $G$  has this property. An unbounded domain  $G$  is called quasibounded if  $\text{dist}(x, \partial G) \rightarrow 0$  whenever  $x$  tends to infinity in  $G$ . If  $G$  is unbounded and not quasibounded then it contains an infinite number of mutually disjoint, congruent balls. If  $\varphi$  is infinitely differentiable, has support in one of these balls, and has nonzero  $L^2(G)$  norm then the set of its translates with supports in the other balls provides a counterexample showing the imbedding  $H_0^1(G) \rightarrow H_0^0(G) \equiv L^2(G)$  is not compact. Thus for an unbounded domain quasiboundedness is necessary for the Rellich property.

In [2] Clark showed that the following Condition 1 is sufficient to guarantee that  $G$  has the Rellich property.

**CONDITION 1.** To each  $R \geq 0$  there correspond positive numbers  $d(R)$  and  $\delta(R)$  satisfying

- (a)  $d(R) + \delta(R) \rightarrow 0$  as  $R \rightarrow \infty$ ,

- (b)  $d(R)/\delta(R) \leq M < \infty$  for all  $R$ ,  
(c) for each  $x \in G$  with  $|x| > R$  there exists  $y$  such that  $|x - y| < d(R)$   
and  $G \cap \{z: |z - y| < \delta(R)\} = \emptyset$ .

This condition is considerably stronger than quasiboundedness. It implies, for example, that  $G$  has nonnull exterior. In [3] Clark gave an example of an unbounded domain having the Rellich property but not satisfying Condition 1. His example was the "spiny urchin," an open connected set in  $E_2$  obtained by removing from the plane all points whose polar coordinates  $(r, \theta)$  satisfy for any  $k = 1, 2, \dots$  the two restrictions  $r \geq k$  and  $\theta = 2^{-k}m\pi$ ,  $m = 1, 2, \dots, 2^{k+1}$ .

In this paper the gap between quasiboundedness as a necessary condition and Condition 1 as a sufficient condition for a domain to have the Rellich property is narrowed from both ends. On the one hand we show that if  $n \geq 2$  then no open set whose boundary consists only of isolated points with no finite accumulation point can have the Rellich property. This settles a question raised by Clark in [3]. On the other hand we show that Condition 1 can be replaced by the following weaker Condition 2, which is still sufficient to guarantee that  $G$  has the Rellich property. In the statement  $B_r(x)$  denotes the open ball of radius  $r$  about  $x$ .

CONDITION 2. There exists  $R_0 \geq 0$  such that to each  $R \geq R_0$  there correspond numbers  $d(R), \delta(R) > 0$  such that

- (a)  $d(R) + \delta(R) \rightarrow 0$  as  $R \rightarrow \infty$ ,  
(b)  $d(R)/\delta(R) < M \leq \infty$  for all  $R \geq R_0$ ,  
(c) for each  $x \in G$  such that  $|x| > R \geq R_0$  the ball  $B_{3d(R)}(x)$  is disconnected into two open components  $C_1$  and  $C_2$  by an  $n - 1$  dimensional manifold forming part of the boundary of  $G$  in such a way that each of the two open sets  $C_i \cap B_{d(R)}(x)$ ,  $i = 1, 2$ , contains a ball of radius  $\delta(R)$ .

Roughly speaking if the  $n - 1$  dimensional manifolds in the boundary of  $G$  are reasonably smooth and unbroken, and bound a quasibounded domain (containing  $G$ ) then  $G$  will satisfy Condition 2. Clark's "spiny urchin" is an example of such a domain. If  $n = 1$  any quasibounded domain satisfies Condition 2, (but not necessarily Condition 1) and so in this case quasiboundedness is necessary and sufficient for the Rellich property.

Our principal results are as follows

**THEOREM 1.** *If  $G$  is open in  $E_n$ ,  $n \geq 2$ , and the boundary of  $G$  consists only of isolated points with no finite accumulation point, then the imbedding  $H_0^1(G) \rightarrow L^2(G)$  is not compact. Thus quasibounded-*

ness is not sufficient to guarantee the Rellich property.

**THEOREM 2.** *If  $G$  is open in  $E_n$  and satisfies Condition 2 then it has the Rellich property.*

For the proof of Theorem 1 we require the following

**LEMMA 1.** *Given  $\rho, \delta > 0, x_0 \in E_n$  ( $n \geq 2$ ), there exists a function  $u \in C^\infty(E_n)$  with the following properties*

- (1)  $u(x) = 0$  in a neighbourhood of  $x_0$
- (2)  $0 \leq u(x) \leq 1$  for all  $x$
- (3)  $u(x) = 1$  outside the ball  $B_\rho(x_0)$
- (4)  $\int_{E_n} |\nabla u(x)|^2 dx \leq \delta^2$ .

*Proof.* Let  $f \in C^\infty(\mathbb{R})$  satisfy  $0 \leq f(t) \leq 1, f(t) = 1$  for  $t \geq 1$  and  $f(t) = 0$  in a neighbourhood of  $t = 0$ . Let  $m$  be a positive integer, put  $r = |x - x_0|$  and define

$$u(x) = v(r) = f([r/\rho]^{1/m}).$$

Clearly  $u \in C^\infty(E_n)$  and satisfies (1), (2) and (3). Also

$$|\nabla u(x)|^2 = \sum_{i=1}^n |D_i u(x)|^2 = |v'(r)|^2.$$

Denoting by  $\omega_n$  the surface area of the unit sphere in  $E_n$  and making the change of variables  $t = (r/\rho)^{1/m}$  we obtain

$$\begin{aligned} \int_{E_n} |\nabla u(x)|^2 dx &= \omega_n \int_0^\rho \left| \frac{d}{dr} f\left(\left[\frac{r}{\rho}\right]^{1/m}\right) \right|^2 r^{n-1} dr \\ &= \omega_n \rho^{n-2} m^{-1} \int_0^1 \left| \frac{d}{dt} f(t) \right|^2 t^{1+m(n-2)} dt \\ &\leq \omega_n \rho^{n-2} m^{-1} [2 + m(n-2)]^{-1} \sup_{0 \leq t \leq 1} |f'(t)|^2 \end{aligned}$$

which, for  $n \geq 2$ , can be made less than  $\delta^2$  for a suitably large choice of  $m$ .

**REMARK.** If  $\varphi \in C_0^\infty(E_n)$  and  $u$  is constructed as above, then  $\varphi \cdot u \in C_0^\infty(E_n - \{x_0\}) \subset H_0^1(E_n - \{x_0\})$ .

*Proof of Theorem 1.* Let  $Q$  be a fixed open ball in  $E_n$ . Let  $\varphi \in C_0^\infty(Q)$  be extended to all of  $E_n$  so that  $\varphi(x) = 0$  in  $E_n - Q$ . Suppose  $\varphi(x) \geq 0$  for all  $x$  and

$$\|\varphi\|_{0, E_n} = C > 0, \quad \|\varphi\|_{1, E_n} = K > 0.$$

There exists  $M > 0$  such that for all  $x$  in  $E_n$

$$|\varphi(x)| \leq M, \quad |D_j \varphi(x)| \leq M, \quad j = 1, \dots, n.$$

If  $Q$  contains no boundary points of  $G$  put  $\psi = \varphi$ . Otherwise  $Q$  contains only a finite number of boundary points of  $G$ , say  $x_1, \dots, x_k$ . For  $i = 1, \dots, k$  let  $B_i = B_{\rho_i}(x_i)$  where  $\rho_i$  is small enough that  $\text{vol. } B_i \leq (C/2kM)^2$ . Let  $\delta = K/Mk$  and let  $u_i$  be the function constructed as in Lemma 1 corresponding to the point  $x_i$  and the constants  $\rho_i$  and  $\delta$ . Put  $\psi = \varphi \cdot u_1 \cdots u_k$ . Clearly  $\psi \in H_0^1(Q - \{x_1, \dots, x_k\}) \subset H_0^1(G)$ . We have

$$\begin{aligned} \|\psi\|_{0,G} &\geq \|\varphi\|_{0,E_n} - \sum_{i=1}^k \|\varphi\|_{0,B_i} \\ &\geq C - \sum_{i=1}^k M(\text{vol. } B_i)^{1/2} \geq \frac{1}{2}C. \end{aligned}$$

Also

$$\begin{aligned} \|D_j \psi\|_{0,G} &\leq \|D_j \varphi\|_{0,E_n} + \sum_{i=1}^k \|\varphi u_1 \cdots D_j u_i \cdots u_k\|_{0,B_i} \\ &\leq K + kM\delta = 2K. \end{aligned}$$

Since  $\|\psi\|_{0,G} \leq \|\varphi\|_{0,G} = C$  we have

$$\|\psi\|_{1,G} \leq (C^2 + 4nK^2)^{1/2} = C_1.$$

Now let  $\{Q_i\}_{i=1}^\infty$  be a family of mutually disjoint open balls in  $E^n$  all congruent to  $Q$ . Let  $\varphi_i$  be a translate of  $\varphi$  with support in  $Q_i$  and let  $\psi_i \in H_0^1(G)$  be constructed from  $\varphi_i$  as above, so that

$$\|\psi_i\|_{0,G} \geq \frac{C}{2}, \quad \|\psi_i\|_{1,G} \leq C_1.$$

Then the sequence  $\{\psi_i\}_{i=1}^\infty$  is bounded in  $H_0^1(G)$  but contains no subsequence convergent in  $L^2(G)$  since for  $i \neq j$   $\|\psi_i - \psi_j\|_{0,G} \geq C/\sqrt{2}$ . Thus the imbedding  $H_0^1(G) \rightarrow L^2(G)$  is not compact.

The proof of Theorem 2 is based on the following generalization of Poincaré's inequality which is a variant on those forms appearing in Agmon [1] and Clark [2].

**LEMMA 2.** *Let  $G$  be open in  $E_n$  and satisfy Condition 2. Let  $G_R$  denote  $G \cap \{x: |x| > R\}$ . Then there exists a constant  $c$  depending only on  $n$  and  $M$  (the constant of Condition 2 (b)) such that for all  $R \geq R_0$  and every  $u \in H_0^1(G)$*

$$\int_{G_R} |u(x)|^2 dx \leq c(d(R))^2 \int_G |\nabla u(x)|^2 dx.$$

*Proof.* Fix  $R \geq R_0$  and let  $d = d(R)$ ,  $\delta = \delta(R)$ . If  $\alpha = (\alpha_1, \dots, \alpha_n)$  is an  $n$ -tuple of integers let  $Q_\alpha = \{x \in E_n : \alpha_k n^{-1/2} d \leq x_k \leq (\alpha_k + 1)n^{-1/2} d\}$ . Then  $E_n = \bigcup_\alpha Q_\alpha$ . Let  $\varphi \in C_0^\infty(G)$ . Fix  $x \in G_R$ . Then  $x \in Q_\alpha$  for some  $\alpha$ . Let  $B_d = B_d(x)$ ,  $B_{3d} = B_{3d}(x)$ . There exists an  $n - 1$  dimensional manifold forming part of  $\partial G$  which disconnects  $B_{3d}$  into open components  $C_1$  and  $C_2$  and there exist points  $y_i \in C_i$  ( $i = 1, 2$ ) such that  $B_\delta(y_i) \subset C_i$ . Thus  $\varphi$  can be written as  $\varphi = \varphi_1 + \varphi_2$  where  $\varphi_i \in C_0^\infty(G)$  and  $\varphi_1 \equiv 0$  in  $C_2$  while  $\varphi_2 \equiv 0$  in  $C_1$ . Since  $Q_\alpha \subset B_d$  we have

$$\int_{Q_\alpha \cap G_R} |\varphi(y)|^2 dy \leq \int_{C_1 \cap B_d} |\varphi_1(y)|^2 dy + \int_{C_2 \cap B_d} |\varphi_2(y)|^2 dy .$$

If  $(r, \sigma)$  and  $S$  denote respectively spherical coordinates in  $E_n$  centered at  $y_2$  and the surface of the unit sphere about  $y_2$  we have

$$\begin{aligned} \int_{C_1 \cap B_d} |\varphi_1(y)|^2 dy &\leq \int_S d\sigma \int_\delta^{2d} |\varphi_1(r, \sigma)|^2 r^{n-1} dr \\ &\leq 2d \int_S |\varphi_1(t, \sigma)|^2 t^{n-1} d\sigma \end{aligned}$$

where  $t = t(\sigma)$  satisfies  $\delta \leq t \leq 2d$ . Since  $\varphi_1(\delta, \sigma) = 0$  it follows that

$$\begin{aligned} |\varphi_1(t, \sigma)^2 t^{n-1}| &= \left| \int_\delta^t \frac{d}{dr} \varphi_1(r, \sigma) dr \right|^2 t^{n-1} \\ &\leq (2d)^n \int_\delta^{2d} \left| \frac{d}{dr} \varphi_1(r, \sigma) \right|^2 dr \\ &\leq (2d)^n \delta^{1-n} \int_\delta^{2d} \left| \frac{d}{dr} \varphi_1(r, \sigma) \right|^2 r^{n-1} dr . \end{aligned}$$

Thus, since  $d/\delta < M$ ,

$$\begin{aligned} \int_{C_1 \cap B_d} |\varphi_1(y)|^2 dy &\leq (2d)^{n+1} \delta^{1-n} \int_S d\sigma \int_\delta^{2d} \left| \frac{d}{dr} \varphi_1(r, \sigma) \right|^2 r^{n-1} dr \\ &\leq 2^{n+1} M^{n-1} d^2 \int_{\delta \leq |y-y_2| \leq 2d} |\nabla \varphi_1(y)|^2 dy \\ &\leq 2^{n+1} M^{n-1} d^2 \int_{B_{3d}} |\nabla \varphi_1(y)|^2 dy . \end{aligned}$$

Combining this with a similar expression for  $\varphi_2$  we obtain

$$\begin{aligned} \int_{Q_\alpha \cap G_R} |\varphi(y)|^2 dy &\leq 2^{n+1} M^{n-1} d^2 \int_{B_{3d}} |\nabla \varphi(y)|^2 dy \\ &\leq 2^{n+1} M^{n-1} d^2 \int_{Q'_\alpha} |\nabla \varphi(y)|^2 dy \end{aligned}$$

where  $Q'_\alpha$  is the union of all the sets  $Q_\alpha$  which intersect  $B_{3d}$ . There

is a number  $N$  depending only on  $n$  such that any  $N + 1$  of the sets  $Q'_\alpha$  have null intersection. Summing the above inequality over all  $\alpha$  for which  $Q_\alpha$  intersects  $G_R$  we obtain

$$\int_{G_R} |\varphi(y)|^2 dy \leq 2^{n+1} N M^{n-1} (d(R))^2 \int_G |\nabla \varphi(y)|^2 dy.$$

This inequality extends by completion to  $H_0^1(G)$ .

The remaining part of the proof of Theorem 2 is similar to Clark's proof [2, Th. 3] and is included here for completeness. First, however, let  $H^m(G, R)$  be the completion in the norm  $\|\cdot\|_{m, G \cap K_R}$  of the space  $C_0^\infty(G, R)$  of all  $C^\infty$  functions whose support is a compact subset of  $G \cap K_R$  where  $K_R = \overline{B_R(0)}$ . Since the imbedding  $H_0^{m+1}(K_R) \rightarrow H_0^m(K_R)$  is known to be compact [4, Chapter XIV] and since an element of  $H^m(G, R)$  can be extended to be zero outside its support so as to belong to  $H_0^m(K_R)$  it follows that the imbeddings  $H^{m+1}(G, R) \rightarrow H^m(G, R)$ ,  $m = 0, 1, 2, \dots$  are compact.

*Proof of Theorem 2.* It suffices, by an inductive argument, to prove only that the imbedding  $H_0^1(G) \rightarrow L^2(G)$  is compact. We make use of the following well known compactness criterion for sets in  $L^2(G)$ : if  $G \subset E_n$  and the sequence  $\{u_k\}_{k=1}^\infty$  is bounded in  $L^2(G)$  then it is compact in  $L^2(G)$  provided

(a) for every bounded  $G' \subset G$  the sequence  $\{u_k | G'\}$  is compact in  $L^2(G')$ , and

(b) for each  $\varepsilon > 0$  there exists  $R > 0$  such that for all  $k$

$$\int_{G_R} |u_k(x)|^2 dx < \varepsilon.$$

Now let  $\{u_k\}$  be a sequence bounded in  $H_0^1(G)$ , say  $\|u_k\|_{1,G} \leq K$ . By Lemma 2, for  $R \geq R_0$  we have  $\|u_k\|_{0,G_R} \leq C(d(R))^2 K \rightarrow 0$  as  $R \rightarrow \infty$  so condition (b) of the criterion is satisfied. To establish (a) let  $G'$  be a bounded subset of  $G$ , so that  $G' \subset K_R$  for some  $R$ . Since  $\{u_k | K_R\}$  is bounded in  $H^1(G, R)$  it is compact in  $H^0(G, R) = L^2(K_R \cap G)$  and so  $\{u_k | G'\}$  is compact in  $L^2(G')$ . Thus  $\{u_k\}$  is compact in  $L^2(G)$ , whence the theorem.

## REFERENCES

1. Shmuel, Agmon, *Lectures on elliptic boundary value problems*, Van Nostrand, Princeton, 1965.
2. Colin Clark, *An embedding theorem for function spaces*, Pacific J. Math. **19** (1966), 243-251.
3. ———, *Rellich's embedding theorem for a "spiny urchin"* Canad. Math. Bull. **10** (1967), 731-734.

4. N. Dunford and J. Schwartz, *Linear operators*, Part II, Interscience, New York, 1963.

Received January 8, 1968. Research partially supported by National Research Council of Canada Operating Grant A-3973.

THE UNIVERSITY OF BRITISH COLUMBIA





# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON  
Stanford University  
Stanford, California 94305

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

RICHARD PIERCE  
University of Washington  
Seattle, Washington 98105

BASIL GORDON\*  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL WEAPONS CENTER

Robert Alexander Adams, <i>Compact Sobolev imbeddings for unbounded domains</i> . . . . .	1
Bernhard Amberg, <i>Groups with maximum conditions</i> . . . . .	9
Tom M. (Mike) Apostol, <i>Möbius functions of order <math>k</math></i> . . . . .	21
Stefan Bergman, <i>On an initial value problem in the theory of two-dimensional transonic flow patterns</i> . . . . .	29
Geoffrey David Downs Creede, <i>Concerning semi-stratifiable spaces</i> . . . . .	47
Edmond Dale Dixon, <i>Matric polynomials which are higher commutators</i> . . . . .	55
R. L. Duncan, <i>Some continuity properties of the Schnirelmann density. II</i> . . . . .	65
Peter Larkin Duren and Allen Lowell Shields, <i>Coefficient multipliers of <math>H^p</math> and <math>B^p</math> spaces</i> . . . . .	69
Hector O. Fattorini, <i>On a class of differential equations for vector-valued distributions</i> . . . . .	79
Charles Hallahan, <i>Stability theorems for Lie algebras of derivations</i> . . . . .	105
Heinz Helfenstein, <i>Local isometries of flat tori</i> . . . . .	113
Gerald J. Janusz, <i>Some remarks on Clifford's theorem and the Schur index</i> . . . . .	119
Joe W. Jenkins, <i>Symmetry and nonsymmetry in the group algebras of discrete groups</i> . . . . .	131
Herbert Frederick Kreimer, Jr., <i>Outer Galois theory for separable algebras</i> . . . . .	147
D. G. Larman and P. Mani, <i>On visual hulls</i> . . . . .	157
R. Robert Laxton, <i>On groups of linear recurrences. II. Elements of finite order</i> . . . . .	173
Dong Hoon Lee, <i>The adjoint group of Lie groups</i> . . . . .	181
James B. Lucke, <i>Commutativity in locally compact rings</i> . . . . .	187
Charles Harris Scanlon, <i>Rings of functions with certain Lipschitz properties</i> . . . . .	197
Binyamin Schwarz, <i>Totally positive differential systems</i> . . . . .	203
James McLean Sloss, <i>The bending of space curves into piecewise helical curves</i> . . . . .	231
James D. Stafney, <i>Analytic interpolation of certain multiplier spaces</i> . . . . .	241
Patrick Noble Stewart, <i>Semi-simple radical classes</i> . . . . .	249
Hiroyuki Tachikawa, <i>On left QF - 3 rings</i> . . . . .	255
Glenn Francis Webb, <i>Product integral representation of time dependent nonlinear evolution equations in Banach spaces</i> . . . . .	269