

Pacific Journal of Mathematics

**RINGS OF FUNCTIONS WITH CERTAIN LIPSCHITZ
PROPERTIES**

CHARLES HARRIS SCANLON

RINGS OF FUNCTIONS WITH CERTAIN LIPSCHITZ PROPERTIES

C. H. SCANLON

Let (X, d) denote a metric space, $L_c(X)$ the ring of real valued functions on X which are Lipschitz on each compact subset of X , $L_l(X)$ the ring of real valued functions on X which are locally Lipschitz relative to the completion of X , and $L_c^*(X)$, $L_l^*(X)$ the bounded elements of $L_c(X)$, $L_l(X)$. The relations between equality of these rings and the topological properties of X are studied. It is shown that a subspace (S, d) of (X, d) is L_c -embedded (or L_c^* -embedded) in (X, d) if and only if S is closed. Further, every subspace of (X, d) is L_l - and L_l^* -embedded in (X, d) .

Su [3] investigated algebraic properties of the rings $L_c(X)$ and $L_c^*(X)$ similar to those of $C(X)$ and $C^*(X)$ by Gillman and Jerrison [2].

2. Equality of rings. Let f denote a real valued function defined on X . f is Lipschitz on $S \subset X$ if and only if there is a real number m , called a Lipschitz constant for f on S , such that if $x, y \in S$, then $|f(x) - f(y)| \leq md(x, y)$. f is locally Lipschitz on X if and only if for each $x \in X$, there is a neighborhood N of x such that f is Lipschitz on N . If $\text{comp } X$ denotes the completion of X , then f is locally Lipschitz with respect to $\text{comp } X$ if and only if for each $x \in \text{comp } X$ there is a neighborhood N of x such that f is Lipschitz on $N \cap X$.

THEOREM 2.1. $f \in L_c(X)$ if and only if f is locally Lipschitz on X .

Sufficiency. Let f be locally Lipschitz on X and S a compact subset of X . Then there exists a finite collection N_1, N_2, \dots, N_m of open sets covering S , on each of which f is Lipschitz and thus bounded. Assuming f is not Lipschitz on S implies that there exists a sequence $\{x_n\}$ from S converging to $x \in S$ and a sequence $\{y_n\}$ from S such that $|f(x_n) - f(y_n)|/d(x_n, y_n) > n$ for each positive integer n . Since f is bounded on S , it follows that $\{y_n\}$ converges to x . Since $x \in N_j$ for some $j = 1, 2, \dots, m$, f is not Lipschitz on N_j which contradicts the definition of N_j .

Necessity. Let $f \in L_c(X)$ and $x \in X$. Assuming f is not locally Lipschitz at x implies there exists sequences $\{x_n\}$ and $\{y_n\}$ such that

$d(x, x_n) < 1/n$, $d(x, y_n) < 1/n$, and $|f(x_n) - f(y_n)|/d(x_n, y_n) > n$. Then $\{p: p \in \{x_n\}, p \in \{y_n\}, \text{ or } p = x\}$ is a compact subset of X on which f is not Lipschitz.

COROLLARY 2.2. $f \in L_c^*(x)$ if and only if f is locally Lipschitz on X and bounded.

Proof. Follows immediately from the definition of $L_c^*(X)$.

COROLLARY 2.3. $L_1(X) \subset L_c(X)$ and $L_i^*(X) \subset L_c^*(X)$.

Proof. If f is locally Lipschitz relative to $\text{com } X$, then f is locally Lipschitz.

LEMMA 2.4. If K is a uniformly bounded set of Lipschitz functions defined on $S \subset X$ and there is a real number m which is a Lipschitz constant for each element of K , then $f(x) = \sup \{g(x): g \in K\}$ for each $x \in S$ is Lipschitz on S and m is a Lipschitz constant for f on S .

Proof. f exists since K is a uniformly bounded set. Assume $x \in S, y \in S$, and

$$(1) \quad f(y) - f(x) - md(x, y) = e > 0.$$

Let $g \in K$ such that

$$(2) \quad f(y) - g(y) < e,$$

then

$$(3) \quad g(y) - g(x) \leq md(x, y).$$

Combining (2) and (3) yields $f(y) - g(x) - md(x, y) < e$, which when combined with (1) gives $f(x) < g(x)$. This contradicts the definition of f .

LEMMA 2.5. Suppose each of c and $r > 0, p \in X$, and for

$$\text{each } x \in X, f(x) = \begin{cases} (c/r)\{r - d(x, p)\} & \text{for } d(x, p) \leq r, \\ 0 & \text{otherwise} \end{cases}$$

then f is Lipschitz on X and (c/r) is a Lipschitz constant for f on X .

Proof. Let $g(x) = (c/r)\{r - d(x, p)\}$ for each $x \in X$. Then for $x, y \in X$,

$$g(x) - g(y) = g(x) - g(p) + g(p) - g(y) ,$$

$$g(x) - g(y) = -(c/r)d(x, p) + (c/r)d(y, p) ,$$

and $g(x) - g(y) \leq (c/r)d(x, y)$ by the triangle property. Since $\sup \{g, 0\}$ is Lipschitz with a Lipschitz constant $\sup \{(c/r), 0\}$ by Lemma 2.4, the conclusion follows.

THEOREM 2.6. *Each of the following is equivalent to each of the others:*

- (1) $L_1(X) = L_c(X)$,
- (2) $L_1^*(X) = L_c^*(X)$, and
- (3) X is complete.

Proof. (1) = (2) obviously. The remaining order is (2) \Rightarrow (3) \Rightarrow (1).

Assume (2) and that X is not complete. Then there exists an $x \in (\text{comp } X) - X$ and a sequence $\{x_n\}$ of distinct points in X such that $\{x_n\}$ converges to x . For each odd integer n , let

$$r_n = \frac{1}{3} \inf \{y: y = d(x_n, x_m) \text{ for } m \neq n \text{ or } y = (1/n)\} ,$$

$$C(x_n, r_n) = \{t \in X: d(t, x_n) \leq r_n\} ,$$

and

$$f_n(t) = \begin{cases} (1/r_n)\{r_n - d(x_n, t)\} & \text{for } t \in C(x_n, r_n) \\ 0 & \text{otherwise} \end{cases}$$

for each $t \in X$. Let $f(t) = \sup \{f_n(t)\}$ for each $t \in X$. If S is a compact subset of X , then S can intersect at most a finite number of the elements of $\{C(x_n, r_n)\}$ and since only a finite number of elements of $\{f_n\}$ are nonzero on S , by Lemma 2.4 f is Lipschitz on S and $f \in L_c^*(X)$. For each neighborhood N in $\text{comp } X$ of x , there is a point $t \in N$ and a point $y \in N$ such that $f(t) = 1$ and $f(y) = 0$. Thus $f \notin L_1(X)$ and by contradiction, (2) \Rightarrow (3).

If (3) is true, $f \in L_1(X)$ if and only if f is locally Lipschitz. Thus by Theorem 2.1, $L_1(X) = L_c(X)$ and (3) \Rightarrow (1).

THEOREM 2.7. $L_c(X) = L_c^*(X)$ if and only if X is compact.

Proof. If X is compact, then each element of $L_c(X)$ is bounded.

Assume $L_c(X) = L_c^*(X)$ and X is not compact. Then there exists a sequence $\{x_n\}$ of distinct points in X which has no convergent subsequence. Let

$$r_n = \frac{1}{3} \inf \left\{ y: y = d(x_n, x_m) \text{ for } n \neq m \text{ or } y = \frac{1}{n} \right\} ,$$

and

$$f(x) = \begin{cases} (n/r_n)\{r_n - d(x_n, x)\} & \text{for } d(x_n, x) \leq r_n \\ 0 & \text{otherwise} \end{cases}$$

for each $x \in X$. By an argument similar to the one for Theorem 2.6, $f \in L_c(X)$. Since $f(x_n) = n$ for each n , $f \in L_c(X) - L_c^*(X)$ which contradicts the assumption.

THEOREM 2.8. $L_1(X) = L_1^*(X)$ if and only if $\text{comp } X$ is compact.

Proof. Each element of $L_1(X)$, $L_1^*(X)$ can be uniquely extended to an element of $L_1(\text{comp } X) = L_c(\text{comp } X)$, $L_1^*(\text{comp } X) = L_c^*(\text{comp } X)$. Since $L_c(\text{comp } X) = L_c^*(\text{comp } X)$ if and only if $\text{comp } X$ is compact by Theorem 2.7, the conclusion follows.

3. If A denotes one of L_1, L_1^*, L_c, L_c^* and $S \subset X$, then the statement that S is A -embedded in X means that if $f \in A(S)$, there is a $g \in A(X)$ such that $g|_S = f$ where $g|_S = \{(x, y) \in g: x \in S\}$.

THEOREM 3.1. If S is a subset of X , then each of the following is equivalent to each of the others:

- (1) S is L_c -embedded in X ,
- (2) S is L_c^* -embedded in X , and
- (3) S is closed.

Proof. Czipszer and Geher [1] proved that if S is a closed subset of X and f is a real valued locally Lipschitz function with domain S , then there is a real valued locally Lipschitz function g with domain X such that $g|_S = f$. Furthermore, they proved that if f is bounded, then there exists a bounded such g . Consequently, by Theorem 2.1, (3) \Rightarrow (1) and (3) \Rightarrow (2).

Assume (2) and S is not closed. Then there exists a sequence $\{x_n\}$ of distinct points in S and a point $x \in X - S$ such that $\{x_n\}$ converges to x . Construct f as in Theorem 2.6. Then $f \in L_c^*(S)$ which has no extension to X in $L_c(X)$. Thus (2) \Rightarrow (3). Note that this also shows (1) \Rightarrow (3).

COROLLARY 3.2. Every subset of X is L_1 -embedded and L_1^* -embedded in X .

Proof. If $S \subset X$, then every element of $L_1(S)$ has a unique extension to the closure of S in $\text{comp } X$ and by Theorems 2.6 and 3.1

an extension in $L_1(\text{comp } X)$ which when restricted to X is an element of $L_1(X)$.

REFERENCES

1. J. Czipser and L. Geher, *Extension of functions satisfying a Lipschitz condition*, Acta Math. Acad. Sci. Hungarica **6** (1955), 213-219.
2. L. Gillman and M. Jerison, *Rings of continuous functions*, D. Van Nostrand, New Jersey, 1960.
3. Li Pi Su, *Algebraic properties of certain rings of continuous functions*, Pacific J. Math. **27** (1968), 175-191.

Received April 3, 1969.

THE UNIVERSITY OF OKLAHOMA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

BASIL GORDON*
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 32, No. 1

January, 1970

Robert Alexander Adams, <i>Compact Sobolev imbeddings for unbounded domains</i>	1
Bernhard Amberg, <i>Groups with maximum conditions</i>	9
Tom M. (Mike) Apostol, <i>Möbius functions of order k</i>	21
Stefan Bergman, <i>On an initial value problem in the theory of two-dimensional transonic flow patterns</i>	29
Geoffrey David Downs Creede, <i>Concerning semi-stratifiable spaces</i>	47
Edmond Dale Dixon, <i>Matric polynomials which are higher commutators</i>	55
R. L. Duncan, <i>Some continuity properties of the Schnirelmann density. II</i>	65
Peter Larkin Duren and Allen Lowell Shields, <i>Coefficient multipliers of H^p and B^p spaces</i>	69
Hector O. Fattorini, <i>On a class of differential equations for vector-valued distributions</i>	79
Charles Hallahan, <i>Stability theorems for Lie algebras of derivations</i>	105
Heinz Helfenstein, <i>Local isometries of flat tori</i>	113
Gerald J. Janusz, <i>Some remarks on Clifford's theorem and the Schur index</i>	119
Joe W. Jenkins, <i>Symmetry and nonsymmetry in the group algebras of discrete groups</i>	131
Herbert Frederick Kreimer, Jr., <i>Outer Galois theory for separable algebras</i>	147
D. G. Larman and P. Mani, <i>On visual hulls</i>	157
R. Robert Laxton, <i>On groups of linear recurrences. II. Elements of finite order</i>	173
Dong Hoon Lee, <i>The adjoint group of Lie groups</i>	181
James B. Lucke, <i>Commutativity in locally compact rings</i>	187
Charles Harris Scanlon, <i>Rings of functions with certain Lipschitz properties</i>	197
Binyamin Schwarz, <i>Totally positive differential systems</i>	203
James McLean Sloss, <i>The bending of space curves into piecewise helical curves</i>	231
James D. Stafney, <i>Analytic interpolation of certain multiplier spaces</i>	241
Patrick Noble Stewart, <i>Semi-simple radical classes</i>	249
Hiroyuki Tachikawa, <i>On left QF - 3 rings</i>	255
Glenn Francis Webb, <i>Product integral representation of time dependent nonlinear evolution equations in Banach spaces</i>	269