

Pacific Journal of Mathematics

ANALYTIC INTERPOLATION OF CERTAIN MULTIPLIER SPACES

JAMES D. STAFNEY

ANALYTIC INTERPOLATION OF CERTAIN MULTIPLIER SPACES

JAMES D. STAFNEY

Let W_p denote the space of all functions on the circle which are the uniform limit of a sequence of trigonometric polynomials which is bounded as a sequence of multipliers for l_p , $1 \leq p \leq 2$. Let U_s be the interpolation space $[W_2, W_1]_s$ (see 1.1). Our main result, Theorem 2.4, states that for a compact subset E of the circle, $U_s|E = C(E)$ if and only if $W_1|E = C(E)$. A major step in the proof is a maximum principle for interpolation, Theorem 1.7. We also give a direct proof that $U_s \neq W_p$ (see Theorem 2.7) for corresponding s and p .

1. Some properties of analytic interpolation.

1.1. Let B^0 and B^1 be two Banach spaces continuously embedded in a topological vector space V such that $B^0 \cap B^1$ is dense in both B^0 and B^1 . For $0 < s < 1$, let \mathfrak{F} , $[B^0, B^1]_s$ and $B^0 + B^1$ denote the spaces as defined in [1, § 1]. For two Banach spaces X and Y we let $O(X, Y)$ denote the Banach space of bounded linear operators from X into Y where the norm is the usual operator norm. Let $O(X)$ denote $O(X, X)$.

1.2. Assume the notation and conditions of paragraph 1.1 and for convenience let B_s denote the space $[B^0, B^1]_s$, $0 < s < 1$. Let V' denote the Banach space

$$O(B^0 \cap B^1, B^0 + B^1).$$

Let A_j be a closed subspace of $O(B^j)$, $j = 0, 1$. By restricting the elements in A_j to $B^0 \cap B^1$ in the obvious way we may regard A_j as continuously embedded in the topological vector space V' , and it is with respect to this embedding that we understand $[A_0, A_1]_s$; in particular, $[A_0, A_1]_s$ is a subspace of V' . We will assume that $A_0 \cap A_1$ is dense in A_j with respect to the norm of A_j , $j = 0, 1$, when these spaces are embedded in V' as described. Since $B^0 \cap B^1$ is dense in B^0 and B^1 , we know from [1, § 9.3] that $B^0 \cap B^1$ is dense in B_s ; thus, since $B_s \subset B^0 + B^1$, the restriction of elements of $O(B_s)$ to $B^0 \cap B^1$ gives a continuous embedding of $O(B_s)$ in V' in the obvious manner. Note that each element of $A_0 \cap A_1$ is bounded with respect to the norm $\| \cdot \|_{B_s}$ restricted to $B^0 \cap B^1$ and is, therefore, contained in the embedded $O(B_s)$. Let A_s denote the closure of $A_0 \cap A_1$ in $O(B_s)$ where $O(B_s)$ is regarded as embedded in V' in the manner just described. Finally, we let M_s and N_s denote the norms of the spaces A_s and $[A_0, A_1]_s$, respectively.

LEMMA 1.3. *Assuming 1.2, $[A_0, A_1]_s \subset A_s$ and $M_s \leq N_s, 0 < s < 1$.*

This lemma is an immediate consequence of [1, § 11.1].

1.4. Assume the notation and conditions of 1.1. Let J be a closed subspace of $B^0 + B^1$. We will assume that

(1.4.1) $I^j = J \cap B_j$, is closed in $B^j, j = 0, 1$. Clearly the map α defined by

$$\alpha(x + I^j) = x + J \quad j = 0, 1$$

is a continuous one to one linear map from B^j/I^j into V/J . Let

$$D_s = [\alpha(B^0/I^0), \alpha(B^1/I^1)]_s .$$

LEMMA 1.5. *Assuming 1.4, if $x \in B_s, 0 < s < 1$, then $x + J \in D_s$ and*

$$(1.5.1) \quad \|x + J\|_{D_s} \leq \|x + (J \cap B_s)\|_{B_s/(J \cap B_s)} .$$

Proof. Let $x \in B_s, h \in J \cap B_s$ and $\varepsilon > 0$. Choose $f \in \mathfrak{F} = \mathfrak{F}(B^0, B^1)$ such that $f(s) = x + h$ and

$$(1.5.2) \quad \|f\|_{\mathfrak{F}} \leq \varepsilon + \|x + h\|_{B_s} .$$

Let $g(\xi) = f(\xi) + J$ for $1 \leq |\xi| \leq \varepsilon$. Then it is clear that $g \in \mathfrak{F}_1$ where

$$\mathfrak{F}_1 = \mathfrak{F}(\alpha(B^0/I^0), \alpha(B^1/I^1))$$

and that

$$(1.5.3) \quad g(s) = x + J .$$

Hence, $x + J \in D_s$. Furthermore, since it is clear that

$$(1.5.4) \quad \|g\|_{\mathfrak{F}_1} \leq \|f\|_{\mathfrak{F}} ,$$

(1.5.1) follows from (1.5.2), (1.5.3), (1.5.4) and the fact that h and ε were chosen arbitrarily.

The following lemma can be proved by the usual method of successive approximations.

LEMMA 1.6. *Suppose that D_1 is a Banach space that is continuously embedded in a Banach space D_0 such that D_1 is dense in D_0 with respect to the norm of D_0 . Suppose that there exist constants $c, c_1, c < 1$, with the property that for each $x \in D_1$ there is a corresponding element z in D_1 such that*

$$|z|_1 \leq c_1 |x|_0 \quad \text{and} \quad |x - z|_0 \leq c |x|_0.$$

Then $D_1 = D_0$.

We will now establish a "maximum principle" for analytic interpolation.

THEOREM 1.7. *If, in addition to the assumptions of paragraph 1.1, $B^0 = [B^0, B^1]_s$ for some s ($0 < s < 1$), then $B^0 = B^1$.*

Proof. From the fact that B^0 and B^1 are continuously embedded in V and the closed graph theorem we conclude that the norms $|\cdot|_0$ and $|\cdot|_s$ on B^0 and $[B^0, B^1]_s$, respectively, are equivalent. In particular, there is a constant c such that

$$(1.7.1) \quad |x|_0 \leq c |x|_s \quad \text{for all } x \text{ in } B^0.$$

From [1, 9.4. (ii)] we conclude that

$$(1.7.2) \quad |x|_s \leq |x|_0^{1-s} |x|_1^s \quad \text{for all } x \text{ in } B^0 \cap B^1.$$

We conclude from (1.7.1) and (1.7.2) that

$$|x|_0 \leq c^{1/s} |x|_1 \quad \text{for all } x \text{ in } B^0 \cap B^1.$$

Thus, B_1 is continuously embedded in B^0 . We shall now prove that (1.7.3) there is a constant c_1 with the property that for each x in B^1 there is a corresponding y in B^1 such that

$$|y|_1 \leq c_1 |x|_0 \quad \text{and} \quad |y - x|_0 \leq (1/2) |x|_0.$$

Let $x \in B^1$. In particular, $x \in [B^0, B^1]_s$ and, therefore, there exists an $f \in \mathfrak{G}(B^0, B^1)$ such that $f(s) = x$ and $|f|_{\mathfrak{G}(B^0, B^1)} \leq 2 |x|_s$. Since the norms $|\cdot|_0$ and $|\cdot|_s$ are equivalent we can choose a real number λ so that $2 |u|_s e^{\lambda s} \leq (1/2) |u|_0$ for every u in B^0 . Let $g(\xi) = f(\xi) e^{-\lambda(\xi-s)}$ where $0 \leq \text{Re } \xi \leq 1$. Then

$$(1.7.4) \quad \begin{aligned} x = g(s) &= \int_{-\infty}^{\infty} g(it) \mu_0(s, t) dt \\ &+ \int_{-\infty}^{\infty} g(1+it) \mu_1(s, t) dt \end{aligned}$$

where μ_0 and μ_1 are the Poisson kernels for the strip $0 \leq \text{Re } \xi \leq 1$ (see [1, 9.4]). Let y and z denote the first and second integrals, respectively, appearing in (1.7.4). Since $\int_{-\infty}^{\infty} |\mu_i(s, t)| dt \leq 1$ ($i = 0, 1$), $|g(it)|_0 \leq 2 |x|_s e^{\lambda s} \leq (1/2) |x|_0$ (all real t), and

$$|g(1+it)|_1 \leq 2 |x|_s e^{-\lambda(1-s)} \leq (1/2) e^{-\lambda} |x|_0$$

(all real t), it follows that $|x - z|_0 \leq (1/2)|x|_0$ and $|z|_1 \leq (1/2)e^{-\lambda}|x|_0$. This proves (1.7.3). Since B^1 is continuously embedded as a dense subspace in B^0 and (1.7.3) holds, the conclusion of Theorem 1.7 follows from Lemma 1.6.

2. The spaces W_p and U_s . Let $l_p, 1 \leq p < \infty$, denote the Banach space of complex valued functions x on the integers such that

$$\|x\|_{l_p} = (\sum |x(n)|^p)^{1/p} < \infty$$

where the sum is over all integers n . Each function α on the integers which vanishes outside some finite set determines a linear transformation T_α on l_p defined by

$$T_\alpha x(n) = \sum_{-\infty < l < \infty} x(n - k)\alpha(k).$$

Let W'_p denote the closure of the operators T_α in $O(l_p)$. Since l_1 is a dense subspace of each space $l_p, 1 \leq p < \infty$, the restriction of elements in $O(l_p), 1 \leq p \leq 2$, to the subspace l_1 gives a one-to-one continuous linear embedding of $O(l_p), 1 \leq p \leq 2$, into the space

$$R = O(l_1, l_2).$$

Throughout this section we will identify $O(l_p)$ with its image under this embedding without further comment. Let U'_s denote the space $[W'_2, W'_1]_s$ where V in 1.1 is, in this case, R .

Our immediate purpose is to define a "Fourier transform" on W'_p and to prove Lemmas 2.2 and 2.3.

If x is a complex valued function on the integers Z , let $\tau_n x(k) = x(k - n)$. Let δ_n denote the function on Z such that $\delta_n(n) = 1$ and $\delta_n(k) = 0, k \neq n$. If x and y are two complex valued function on Z let

$$x*y(m) = \sum_{n \in Z} x(m - n)y(n)$$

define the function $x*y$ provided the sum converges absolutely for each $m \in Z$. For each H in W'_p let H^\sim denote the function $H(\delta_0)$ in l_p . The following lemma states the needed properties of the map $H \rightarrow H^\sim$. Note that $\tau_n x = \delta_n * x$ for each $n \in Z$ and for each complex valued function x on Z .

LEMMA 2.1.

(2.1.1) $H \rightarrow H^\sim$ is a one-to-one linear transformation from W'_p into l_p .

(2.1.2) $Hx = H^\sim * x, H \in W'_p, x \in l_p$.

(2.1.3) $(HK)^\sim = H^\sim * K^\sim, H, K \in W'_p$.

Proof. The map $H \rightarrow H^\sim$ is clearly linear. Evidently, each H in

W'_p commutes with all operators $\tau_m, m \in Z$, since the operators of the form T_α commute with the operators $\tau_m, m \in Z$. Thus for $H \in W'_p$ and $m \in Z$, we see that

$$(2.1.4) \quad H(\delta_m) = H(\tau_m \delta_0) = \tau_m H(\delta_0) = \tau_m H^\sim = H^\sim * \delta_m .$$

From this we see that since the linear span of the elements δ_m is dense in l_p , the map $H \rightarrow H^\sim$ is one-to-one. Obviously, H^\sim is in l_p . To establish (2.1.2) we first note that since H^\sim is in $l_q(q^{-1} + p^{-1} = 1)$ the map $x \rightarrow H^\sim * x$ is a continuous linear map from l_p into c_0 , the space of complex valued functions on Z which tend to 0 at $\pm \infty$. The map $x \rightarrow Hx$ is also a continuous linear map from l_p into c_0 . These observations together with (2.1.4) and the density property of the δ_m 's noted above complete the proof of (2.1.2). To prove (2.1.3) we note that for H and K in $W'_p, K^\sim \in l_p$, so by (2.1.2) we have

$$H^\sim * K^\sim = H(K^\sim) = H(K\delta_0) = (HK)\delta_0 = (HK)^\sim .$$

This completes the proof of the lemma.

Let $L_p(1 \leq p < \infty)$ denote the Banach space of measurable functions $g(\theta)$ on the circle (reals mod 2π) whose norm $\|g\|_{L_p}$,

$$\|g\|_{L_p} = \left((1/2\pi) \int_0^{2\pi} |g(\theta)|^{1/p} d\theta \right)^{1/p} ,$$

is finite. Let L_∞ denote the space of essentially bounded measurable functions g with $\|g\|_{L_\infty}$ denoting the essential supremum of g .

Since each function $H^\sim, H \in W'_p$, is in l_p , which is contained in l_2 , there is a unique function H^\wedge in L_2 such that $\sum H^\sim(n)e^{in\theta}$ is the Fourier series of H^\wedge .

LEMMA 2.2. *For $1 \leq p \leq 2$ the map $H \rightarrow H^\wedge$ is a norm decreasing algebraic isomorphism from W'_p into L_∞ .*

Proof. The fact that $H \rightarrow H^\wedge$ is a one-to-one linear map from W'_p into L_2 is clear from (2.1.1) and the fact that each function in L_2 is uniquely determined by its Fourier coefficients. For each $f \in L_1$, let $\lambda(f)$ denote the function on Z defined by:

$$\lambda(f)(n) = (1/2\pi) \int_0^{2\pi} f(\theta)e^{-in\theta} d\theta .$$

It is clear from the Schwarz inequality that the map $(f, g) \rightarrow \lambda(f \cdot g)(n)$ is a continuous bilinear functional on $L_2 \oplus L_2$ for each integer n . On the other hand, the map

$$(f, g) \rightarrow (\lambda(f) * \lambda(g))(n)$$

is also a continuous bilinear functional on $L_2 \oplus L_2$. Since these functionals (for each n) clearly agree when f and g are trigonometric polynomials, they must agree on $L_2 \oplus L_2$. Since λ is a one-to-one map, the multiplicative property of $H \rightarrow H^\wedge$ now follows from (2.1.3). To prove that the map is norm decreasing we first note the following inequalities:

$$\|H^n\|_{W'_p} \geq \|H^n \delta_0\|_{l_p} = \|(H^n)^\sim\|_{l_p} \geq \|(H^n)^\sim\|_{l_2} = \|(H^n)^\wedge\|_{L_2} = \|(H^\wedge)^n\|_{L_2}.$$

It is well known that $(\|H^n\|_{W'_p})^{1/n}$ converges to the spectral radius of H , which is dominated by $\|H\|_{W'_p}$, and that $(\|(H^\wedge)^n\|_{L_2})^{1/n}$ converges to $\|H^\wedge\|_{L_\infty}$ as $n \rightarrow \infty$. This proves the lemma.

Let W_p and U_s denote the functions on the circle of the form H^\wedge where $H \in W'_p, U'_s$, respectively. The following lemma is an immediate consequence of Lemma 2.2.

LEMMA 2.3. *W_p consists precisely of the functions on the circle which are the uniform limits of sequences H_n^\wedge of trigonometric polynomials such that H_n is a Cauchy sequence in W'_p .*

For any subset E of the circle group $U_s|E$ denotes the functions on E obtained by restricting the functions of U_s to E and $C(E)$ denotes the continuous complex valued functions on E .

THEOREM 2.4. *Suppose that E is a compact subset of the circle group and $0 < s < 1$. Then $U_s|E = C(E)$ if and only if $W_1|E = C(E)$.*

Proof. First assume that $W_1|E = C(E)$. By Lemma 1.3, $U'_s \subset W'_p$; consequently, $U_s \subset W_p$. We conclude from Lemma 2.3 that $W_p \subset C(E)$. Thus, $U_s|E \subset C(E)$. Since $W'_2 \supset W'_1$, it is clear from the definition of interpolation that $U'_s \supset W'_1$. Thus, $U_s|E \supset C(E)$.

Consider the converse and assume that $U_s|E = C(E)$. In 1.4 we let $B^0 = W'_2, B^1 = W'_1, V = R$ and

$$J = \{a \in W'_2; \hat{a}(\theta) = 0, \theta \in E\}.$$

The assumptions on J in 1.4 are clearly satisfied since by Lemma 2.2, the maps $a \rightarrow \hat{a}$ are continuous on W'_1 and W'_2 . By Theorem 1.5, if $x \in U'_s$, then $x + J$ is in the space

$$(2.4.1) \quad [\alpha(W'_2/J), \alpha(W'_1/(J \cap W'_1))]_s.$$

However, by hypothesis, the cosets in V of the form $x + J, x \in U'_s$, are the same as the cosets $y + J, y \in W'_2$. Therefore, the space in (2.4.1) is $\alpha(W'_2/J)$. Since $W'_2 \supset W'_1$,

$$\alpha(W'_2/J) \supset \alpha(W'_1/(J \cap W'_1));$$

therefore, we conclude from 1.7 that

$$\alpha(W_2'/J) = \alpha(W_1'/(J \cap W_1')) ;$$

or, what is the same thing, that $W_1|E = C(E)$. This completes the proof.

COMMENT 2.5. It is natural to compare U_s and W_p where $[l_2, l_1]_s = l_p$, i.e., $(1-s)/2 + s = 1/p$. In [3] we showed that Theorem 2.4 is not valid for W_p . To be exact, there is a compact subset E of the circle such that $W_p|E \neq C(E) = W_{4/3}|E$, $1 \leq p < 4/3$. We had originally used this result to show that $W_p \neq U_s$; however, the referee has suggested a direct proof which we will now give.

LEMMA 2.6. *Let h_n be a sequence in U_s , $0 < s < 1$, such that $\|h_n\|_s \leq M$ (here $\|\cdot\|_s$ is the norm in U_s) and $h_n \rightarrow h$ almost everywhere. Then h agrees with some continuous function almost everywhere.*

Proof. Since $\|h_n\|_s \leq M$ there exist functions $f_n(\theta, \xi)$, analytic in ξ for $0 < B(\xi) < 1$ and continuous in $0 \leq B(\xi) \leq 1$, such that for any real number t , $\|f_n(\theta, it)\|_0 \leq 2M$, $\|f_n(\theta, 1+it)\|_1 \leq 2M$ and $f_n(\theta, s) = h_n(\theta)$. Let $g_n(\theta, \xi) = f_n(\theta, \xi)e^{+\lambda(\xi-s)}$. Then

$$\begin{aligned} h_n(\theta) = f_n(\theta, s) = g_n(\theta, s) &= \int_{-\infty}^{+\infty} g_n(\theta, it)\mu_0(s, t)dt \\ &+ \int_{-\infty}^{+\infty} g_n(\theta, 1+it)\mu_1(s, t)dt \\ &= u_n(\theta) + v_n(\theta) \end{aligned}$$

where μ_0 and μ_1 are the Poisson Kernels for the strip (see [1, 9.4]). Evidently $\|u_n\|_0 \leq 2e^{-\lambda s}M$, $\|v_n\|_1 \leq 2e^{\lambda(1-s)}M$. Since the v_n are uniformly bounded, by taking a subsequence if necessary, we may assume that v_n converges weakly to a bounded function $v(\theta)$, that is

$$\lim_{n \rightarrow \infty} \int v_n(\theta)\varphi(\theta)d\theta = \int v(\theta)\varphi(\theta)d\theta$$

for every integrable φ . Furthermore, as is readily seen, $v(\theta)$ belongs to U_1 and therefore is continuous. Since h_n is uniformly bounded and converges almost everywhere, h_n converges weakly. Since h_n and v_n converge weakly, u_n converges weakly to some function u . From the fact that $|u_n(\theta)| \leq \|u_n\|_0 \leq 2e^{-\lambda s}M$, it follows that $|u(\theta)| \leq 2e^{-\lambda s}$ almost everywhere. Since $h = u + v$ almost everywhere and λ can be taken arbitrarily large, h agrees almost everywhere with the uniform limit of continuous functions. This completes the proof of the lemma.

THEOREM 2.7. U_s is properly contained in W_p for $1 < p < 2$.

Proof. To prove the theorem it suffices to exhibit a sequence of functions in U_s whose norms in U_s tend to infinity and whose norms in W_p remain bounded. Let $h(e^{it}) = 1$ for $0 \leq t \leq \pi$ and $h(e^{it}) = 0$ for $\pi < t < 2\pi$. Then h is a multiplier for l_p (see [2]), which does not agree almost everywhere with any continuous function. Let φ_n be defined by: $\varphi_n(e^{it}) = n$ for $|t| \leq 1/2n$, $\varphi_n(e^{it}) = 0$ otherwise, $n = 1, 2, \dots$. Let $h_n = h * \varphi_n$, $n = 1, 2, \dots$. Since $\int_0^{2\pi} |h_n(e^{it})| dt = 1$, it follows that the W_p norm of h_n is the same as the W_p norm of h ; thus, h_n is bounded in W_p . Since both h and φ_n belong to $L_2(0, 2\pi)$, $h_n \in W_1 \subset U_s$. Obviously, h_n converges to h almost everywhere. Since h does not agree almost everywhere with any continuous function, it follows from Lemma 2.6 that h_n is not bounded in U_s .

BIBLIOGRAPHY

1. A. P. Calderón, *Intermediate spaces and interpolation, the complex method*, *Studia Math.* **24** (1964), 113-190.
2. I. I. Hirshmann, *On multiplier transformations*, *Duke Math. J.* **26** (1959), 221-242.
3. James D. Stafney, *Approximation of W_p -continuity sets by p -Sidon sets*, *Michigan Math. J.* **16** (1969), 161-176.

Received August 26, 1968.

UNIVERSITY OF CALIFORNIA, RIVERSIDE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

BASIL GORDON*
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Robert Alexander Adams, <i>Compact Sobolev imbeddings for unbounded domains</i>	1
Bernhard Amberg, <i>Groups with maximum conditions</i>	9
Tom M. (Mike) Apostol, <i>Möbius functions of order k</i>	21
Stefan Bergman, <i>On an initial value problem in the theory of two-dimensional transonic flow patterns</i>	29
Geoffrey David Downs Creede, <i>Concerning semi-stratifiable spaces</i>	47
Edmond Dale Dixon, <i>Matric polynomials which are higher commutators</i>	55
R. L. Duncan, <i>Some continuity properties of the Schnirelmann density. II</i>	65
Peter Larkin Duren and Allen Lowell Shields, <i>Coefficient multipliers of H^p and B^p spaces</i>	69
Hector O. Fattorini, <i>On a class of differential equations for vector-valued distributions</i>	79
Charles Hallahan, <i>Stability theorems for Lie algebras of derivations</i>	105
Heinz Helfenstein, <i>Local isometries of flat tori</i>	113
Gerald J. Janusz, <i>Some remarks on Clifford's theorem and the Schur index</i>	119
Joe W. Jenkins, <i>Symmetry and nonsymmetry in the group algebras of discrete groups</i>	131
Herbert Frederick Kreimer, Jr., <i>Outer Galois theory for separable algebras</i>	147
D. G. Larman and P. Mani, <i>On visual hulls</i>	157
R. Robert Laxton, <i>On groups of linear recurrences. II. Elements of finite order</i>	173
Dong Hoon Lee, <i>The adjoint group of Lie groups</i>	181
James B. Lucke, <i>Commutativity in locally compact rings</i>	187
Charles Harris Scanlon, <i>Rings of functions with certain Lipschitz properties</i>	197
Binyamin Schwarz, <i>Totally positive differential systems</i>	203
James McLean Sloss, <i>The bending of space curves into piecewise helical curves</i>	231
James D. Stafney, <i>Analytic interpolation of certain multiplier spaces</i>	241
Patrick Noble Stewart, <i>Semi-simple radical classes</i>	249
Hiroyuki Tachikawa, <i>On left QF - 3 rings</i>	255
Glenn Francis Webb, <i>Product integral representation of time dependent nonlinear evolution equations in Banach spaces</i>	269