

Pacific Journal of Mathematics

COMPACT INTEGRAL DOMAINS

JOE EBELING CUDE

COMPACT INTEGRAL DOMAINS

JOE E. CUDE

It is well known that if A is a compact integral domain and R is its Jacobson radical, then $A = R$ or A/R is a division ring and A has an identity. The object of this paper is to investigate some of the algebraic properties of A .

If A has an identity and finite characteristic, then there exists a maximal subfield F of A which is isomorphic to A/R . Furthermore A is topologically isomorphic to $F + R$. The existence of a subfield is a necessary and sufficient condition for A to have finite characteristic. If A does not have an identity but does have finite characteristic, then it can be openly embedded in a compact integral domain with an identity. Finally, the main result shows that if the center of A is open, then A is commutative.

1. Preliminaries. An integral domain is defined to be a ring with more than one element such that the nonzero elements form a multiplicative semigroup (which is not necessarily commutative). It is always assumed that a topological ring is Hausdorff.

Throughout this paper A will denote a compact integral domain and R will denote its Jacobson radical.

It has been shown [4, Lemma 2, p. 279] that A has a fundamental system of neighborhoods at zero consisting of open (two-sided) ideals. Kaplansky has shown that R is open [3, Th. 7, p. 161]. Also, that $A = R$ or A/R is a division ring and A has an identity [3, Th. 19, p. 168].

Because A is an integral domain, it can have no elements or ideals that are nilpotent in an algebraic sense, but nilpotency can be defined in a topological sense. We say that an element x is nilpotent if $\lim x^n = 0$, and that an ideal V is nilpotent if for every open set containing zero, there is an integer N such that for every $n \geq N$, $V^n \subset W$ where V^n is the set of all finite sums of the product of n elements of V .

It has been shown that R is nilpotent [3, Th. 14, p. 163].

The following theorem, which may be thought of as an extension of Wedderburn's theorem, immediately follows from the above results.

THEOREM 1. *Any compact semi-simple integral domain is a finite field.*

2. General properties. In this section it is assumed unless otherwise stated that A is a compact integral domain with an identity. The units and nilpotent elements of A are characterized by the following lemma.

LEMMA 1. *If A is a compact integral domain with an identity and $x \in A$, then x has an inverse if and only if $x \notin R$, and x is nilpotent if and only if $x \in R$.*

Proof. If $x \notin R$, then there exists an $x' \in A$ such that $xx' - 1 \in R$. Let y be the right quasi-inverse of $xx' - 1$, then

$$xx' - 1 + y + (xx' - 1)y = 0,$$

which implies that $x[x'(1 + y)] = 1$.

If x has an inverse, then it can not be nilpotent. Since R is itself nilpotent, and by the above all elements not in R have inverses, R consists of all the nilpotent elements of A .

THEOREM 2. *If A is a compact integral domain with an identity and has characteristic p , then A contains a maximal subfield which is isomorphic to A/R .*

Proof. Let $a + R$ be a generator of A/R and let the number of elements of A/R be p^q . Now for every positive integer k , $a^{p^{kq}}$ belongs to $a + R$. Since A is compact and R is closed, $a + R$ is closed and compact. Hence there exists a subnet $\{a^{p^{k(i)q}}\}$ of the sequence $\{a^{p^{kq}}\}$ which converges to some a^* belonging to $a + R$. Hence $a^* + R$ generates A/R . If $n = p^q - 1$, then $a^n - 1 \in R$ and $0 = \lim (a^n - 1)^{p^{k(i)q}} = (a^*)^n - 1$. The ring F generated by $\{j1 \mid 0 \leq j \leq p - 1\}$ and $\{a^*\}$ is a finite field containing p^q elements.

Let D be any subfield of A . For every $d \in D$, let $\theta(d) = d + R$. Clearly θ is a homomorphism mapping D into A/R . If $\theta(d_1) = \theta(d_2)$, $d_1 - d_2 \in R$, and if $d_1 - d_2 \neq 0$, then $1 \in R$. Since this is impossible, θ is a monomorphism and D contains at most p^q elements.

The field F constructed above is hence a maximal subfield of A and is isomorphic to A/R .

Note that in the above Theorem, D could have been a subdivision ring of A , and hence any subdivision ring of A is a finite field.

Zelinsky has shown [7, Th. E, p. 321] that if A has finite characteristic, then $A = S + R$ (group direct sum) where S is a compact subring of A . If in addition A has an identity, then since R is open and F is finite, the following theorem immediately follows.

THEOREM 3. *If A is a compact integral domain with an identity having finite characteristic, and if F is a maximal subfield of A , then A is topologically isomorphic to $F + R$.*

Not all compact integral domains have finite characteristic as is seen in the following example. Let Q be the field of rational numbers and P be any prime divisor of Q which is nonarchimedean. Let A^* be the valuation ring at P with the usual topology. Now A^* is a topological integral domain. Let A be the completion of A^* . A is an integral domain [5] which is compact [6, Lemma 1, p. 434], but it does not have finite characteristic.

It should also be noted that A does not have a subfield. As is seen in the following theorem, the existence of a subfield of A is equivalent to A having finite characteristic.

THEOREM 4. *If A is a compact integral domain with an identity, then A has finite characteristic if and only if A has a subfield.*

Proof. Let F be a subfield of A . As in Theorem 2, F is finite, and hence it has finite characteristic. Since A can have only one nonzero idempotent, the identity of F must be the identity of A , and hence A has finite characteristic.

The other implication is obvious.

In the rest of this section let us assume that A is a compact integral domain with characteristic p which does not have an identity. We say that a topological ring B can be openly embedded in a topological ring C if there exists a continuous open monomorphism which maps B into C .

THEOREM 5. *If A is a compact integral domain with finite characteristic and does not have an identity, then A can be openly embedded in a compact integral domain with an identity.*

Proof. Let $K = \{j \mid 0 \leq j \leq p-1\}$ be the discrete field of integers mod p and let $A^* = A \times K$. For every $(x, i) \in A^*$ and $(y, j) \in A^*$ define

$$(x, i) + (y, j) = (x + y, i + j), \quad \text{and}$$

$$(x, i) \cdot (y, j) = (xy + jx + iy, ij).$$

With the usual product topology A^* is a compact ring.

Let $P = \{(x, i) \in A^* \mid xa + ia = 0 \text{ for every } a \in A\}$. Now P is a closed two sided ideal in A^* . Furthermore it is easily seen that A^*/P is an integral domain with an identity $(0, 1)$. For every $a \in A$, let $\theta(a) = (a, 0) + P$. Clearly θ is a continuous open monomorphism, and

hence A can be openly embedded in A^*/P , a compact integral domain with an identity.

3. Commutativity of compact integral domain. If A has an identity, then A/R is of course a commutative ring, and it would be tempting to assert that A itself is commutative. However in general, this is not true as seen in the following example.

Let D be any finite discrete topological field containing p^q elements where $q > 1$. Let σ be any automorphism on D . For every integer $i \geq 0$, let $D(i) = D$. Also let $A = \prod_{i=0}^{\infty} D(i)$. With the usual product topology A is a compact Hausdorff space. For every $f \in A$ and $g \in A$, define $(f + g)(i) = f(i) + g(i)$ and $(fg)(i) = \sum_{k+m=i} f(k)\sigma^k[g(m)]$. With the above operations A is a compact integral domain with an identity and has characteristic p . A is not commutative if σ is not the identity automorphism. The above construction is called a Hilbert construction [1, pp. 43-44].

It is interesting to note that although commutativity is an algebraic property, it may depend upon a topological property as is seen in the main result.

THEOREM 6. *If A is a compact integral domain, then A is commutative if and only if its center is open.*

Proof. Assume that Z is the center of A and that it is open. The center of R , $Z(R)$, contains $Z \cap R$. Because Z is open and $0 \in Z$, for every $x \in R$, there exists an $n(x) \geq 1$ such that $x^{n(x)} \in Z$, and hence $x^{n(x)} \in Z(R)$. Since R itself is an integral domain, R is commutative [2, Corollary, p. 219]. Furthermore, $R \subset Z$ since for every $x \in R$ and $a \in A$, $axx = xax$ which implies $ax = xa$.

If A has no identity, then $A = R$, and A is commutative.

If A has an identity, then A/R is a finite field containing p^q elements for some prime p , and for every $x \in A$, $x^{p^q} - x \in R$. Thus for every $x \in A$, $x^{p^q} - x \in Z$ which implies that A is commutative by a result of Herstein [2, Th. 2, p. 221].

The other implication is trivial.

REFERENCES

1. E. Artin, *Geometric algebra*, Interscience Tracts in Pure and Applied Mathematics, 3, Interscience Publishers, Inc., New York, N. Y., 1957.
2. N. Jacobson, *Structure of rings*, Rev. ed., American Mathematical Colloquium Publications XXXVII, American Mathematical Society, Providence, Rhode Island, 1964.
3. I. Kaplansky, *Topological rings*, Amer. J. Math. **69** (1947), 153-183.
4. Y. Otobe, *On quasi-evaluations of compact rings*, Proc. Imp. Acad. Tokyo **20** (1944), 278-282.

5. E. Weiss, *Algebraic number theory*, International Series in Pure and Applied Mathematics, McGraw-Hill Book Company, Inc. New York, N. Y., 1963.
6. D. Zelinsky, *Rings with ideal nuclei*, Duke Math. J. **18** (1951), 431-442.
7. ———, *Raising idempotents*, Duke Math. J. **21** (1954), 315-321.

Received June 5, 1969.

WASHINGTON STATE UNIVERSITY
AND
DALLAS BAPTIST COLLEGE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

BASIL GORDON*
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

* Acting Managing Editor.

Pacific Journal of Mathematics

Vol. 32, No. 3

March, 1970

Shair Ahmad, <i>Dynamical systems of characteristic 0^+</i>	561
Charles A. Akemann and Bernard Russo, <i>Geometry of the unit sphere of a C^*-algebra and its dual</i>	575
Philip Bacon, <i>The compactness of countably compact spaces</i>	587
Richard Blaine Barrar and Henry Loeb, <i>On the continuity of the nonlinear Tschebyscheff operator</i>	593
L. Carlitz, <i>Factorization of a special polynomial over a finite field</i>	603
Joe Ebeling Cude, <i>Compact integral domains</i>	615
Frank Rimi DeMeyer, <i>On automorphisms of separable algebras. II</i>	621
James B. Derr, <i>Generalized Sylow tower groups</i>	633
Raouf Doss, <i>Some inclusions in multipliers</i>	643
Mary Rodriguez Embry, <i>The numerical range of an operator</i>	647
John Froese, <i>Domain-perturbed problems for ordinary linear differential operators</i>	651
Zdeněk Frolík, <i>Absolute Borel and Souslin sets</i>	663
Ronald Owen Fulp, <i>Tensor and torsion products of semigroups</i>	685
George Grätzer and J. Plonka, <i>On the number of polynomials of an idempotent algebra. I</i>	697
Newcomb Greenleaf and Walter Read, <i>Positive holomorphic differentials on Klein surfaces</i>	711
John Willard Heidel, <i>Uniqueness, continuation, and nonoscillation for a second order nonlinear differential equation</i>	715
Leon A. Henkin, <i>Extending Boolean operations</i>	723
R. Hirshon, <i>On hopfian groups</i>	753
Melvin Hochster, <i>Totally integrally closed rings and extremal spaces</i>	767
R. Mohanty and B. K. Ray, <i>On the convergence of a trigonometric integral</i>	781
Michael Rich, <i>On a class of nodal algebras</i>	787
Emile B. Roth, <i>Conjugate space representations of Banach spaces</i>	793
Rolf Schneider, <i>On the projections of a convex polytope</i>	799
Bertram Manuel Schreiber, <i>On the coset ring and strong Ditkin sets</i>	805
Edgar Lee Stout, <i>Some remarks on varieties in polydiscs and bounded holomorphic functions</i>	813
James Edward Ward, <i>Two-groups and Jordan algebras</i>	821