

Pacific Journal of Mathematics

**UNIQUENESS, CONTINUATION, AND NONOSCILLATION FOR
A SECOND ORDER NONLINEAR DIFFERENTIAL EQUATION**

JOHN WILLARD HEIDEL

UNIQUENESS, CONTINUATION, AND NONOSCILLATION FOR A SECOND ORDER NONLINEAR DIFFERENTIAL EQUATION

J. W. HEIDEL

This paper considers the differential equation (1) $y'' + f(t)y^\gamma = 0$ where $f(t)$ is nonnegative and continuous on $[0, \infty)$ and γ is the quotient of odd, positive integers. For this equation we discuss uniqueness of the zero solution, continuation of solutions to $[0, \infty)$, and nonoscillation of solutions. Using a relation between uniqueness and continuation on the one hand and nonoscillation on the other, we can show that the condition $f'(t) \leq 0$ in Atkinson's nonoscillation theorem (*Pacific J. Math.* 5 (1955), 643-647), and in a corresponding theorem for $0 < \gamma < 1$, cannot be removed entirely.

The equation to be considered is

$$(1) \quad y'' + f(t)y^\gamma = 0$$

where $f(t)$ is nonnegative and continuous on $[0, \infty)$ and $\gamma = p/q$ where p and q are odd positive integers. These assumptions will hold throughout this paper. We will be concerned only with real valued solutions. Coffman and Ullrich [2] have shown that if, in addition, $f(t)$ is positive and locally of bounded variation on $[0, \infty)$ and $\gamma > 1$, then all solutions can be continued to $[0, \infty)$. Our purpose here is threefold. First we give a result involving uniqueness of the zero solution of (1) which is analogous to Coffman and Ullrich's, but for the case $0 < \gamma < 1$. Secondly we extend these two results to allow $f(t)$ to have isolated zeros. Finally, we give a connection between either continuation or uniqueness on the one hand and the nonoscillation problem for (1).

We begin with some definitions and basic facts. The assumption on γ yields that solutions of (1) with real valued initial conditions are real valued and the negative of a solution of (1) is again a solution.

LEMMA 1. *If $0 < \gamma < 1$, all solutions of (1) can be continued to $[0, \infty)$. If $1 < \gamma$, then there is a unique solution satisfying any set of initial conditions.*

Continuation follows from a theorem of Wintner (Hartman [4, p. 29]). Uniqueness follows from the local Lipschitz condition. Here we will be concerned with continuation for $1 < \gamma$ and its analogue for $0 < \gamma < 1$ which is uniqueness of the zero solution (trivial solution). By definition we say that the zero solution is unique if any solution $y(t)$ of (1)

satisfying the initial conditions $y(t_0) = y'(t_0) = 0$ for some $t_0 \geq 0$ will then necessarily satisfy $y(t) = 0$ for all $t \geq 0$. A nontrivial solution of (1) which has zero initial conditions at some point will be called singular, following Kiguradze [6].

LEMMA 2. *Suppose $0 < \gamma < 1$. A nontrivial solution $y(t)$ of (1) is singular if and only if $y(t)$ has an infinite number of zeros on a finite interval.*

Proof. Suppose $y(t)$ is singular, $y^2(t_0) + y'^2(t_0) > 0$, $y^2(t_1) + y'^2(t_1) = 0$, and $t_0 < t_1$. Suppose that $y(t)$ has no zero on (t_0, t_1) , and that $y(t) > 0$ on (t_0, t_1) . Then $y'(t)$ is decreasing on (t_0, t_1) and therefore contradicts $y'(t_1) = 0$. If $y(t)$ has a last zero before t_1 , at t_2 , then the same argument holds on (t_2, t_1) . Therefore $y(t)$ does not have a last zero on (t_0, t_1) , that is, $y(t)$ has an infinity of zeros on (t_0, t_1) , clustering at t_1 . A similar argument applies if $t_1 < t_0$.

Conversely, suppose that $y(t)$ has an infinity of zeros on some finite interval (t_0, t_1) . Then they cluster at some t^* , $t_0 \leq t^* \leq t_1$; by continuity $y(t^*) = 0$, and by the mean value theorem and continuity $y'(t^*) = 0$.

2. In this section we consider uniqueness of zero solution when $0 < \gamma < 1$. The techniques are modified from Coffman and Ullrich [2].

THEOREM 1. *Suppose that $0 < \gamma < 1$ and that $f(t)$ is positive and class C' on $[0, \infty)$. Then the zero solution of (1) is unique.*

Proof. Suppose that $y(t)$ is a solution of (1) such that $(y(t_0))^2 + (y'(t_0))^2 > 0$ for some $t_0 \geq 0$. It suffices to show that $(y(t))^2 + (y'(t))^2 > 0$ for all $t \geq 0$. Let $t_1 > t_0$. Let

$$\Phi(t) = \frac{2f(t)}{\gamma + 1} (y(t))^{\gamma+1} + (y'(t))^2.$$

Then $\Phi'(t) = (2/(\gamma + 1))f'(t)(y(t))^{\gamma+1}$. Therefore

$$\Phi'(t) \geq -\frac{2}{\gamma + 1} \frac{|f'(t)|}{f(t)} \Phi(t).$$

An integration shows that

$$\Phi(t) \geq \Phi(t_0) \exp\left(-\int_{t_0}^t g(s) ds\right), \quad t_0 \leq t \leq t_1$$

where $g(t) = (2/(\gamma + 1))(|f'(t)|/f(t))$. Since

$$\int_{t_0}^{t_1} (|f'(x)|/|f(s)|) ds < \infty ,$$

it follows that $\Phi(t) > 0$ for $t_0 \leq t \leq t_1$. Therefore $(y(t))^2 + (y'(t))^2 > 0$ for $t_0 \leq t \leq t_1$. Since t_1 is arbitrary, $(y(t))^2 + (y'(t))^2 > 0$ for $t_0 \leq t$. A similar argument applies for $t_1 < t_0$.

REMARK. In their continuation theorem Coffman and Ullrich were able to replace class C' by locally of bounded variation. Their approximation argument breaks down in our case because we are dealing with uniqueness rather than continuation.

EXAMPLE 1. The example given by Coffman and Ullrich of an equation (1) with a noncontinuable solution can be modified to give an example where the zero solution is not unique, when $0 < \gamma < 1$. Since this example will be used later in this paper, we will outline its construction. The following lemma is a simple modification of a lemma of Coffman and Ullrich [2].

LEMMA 3. Suppose $0 < \gamma < 1$. For each positive integer n , there exists a continuous function $q_n(t)$ on $[0, 1]$ with $q_n(0) = q_n(1) = 0$ and such that

$$(2) \quad \frac{d^2 U}{dt^2} + (C^2 + q_n(t))U^\gamma = 0 ,$$

where C is a suitable constant, has a solution $U_n(t)$ satisfying

$$\frac{dU_n(0)}{dt} = \frac{dU_n(1)}{dt} = 0, \quad U_n(0) = 1, \quad U_n(1) = \left(\frac{n}{n+1}\right)^{4/(1-\gamma)}$$

and having at least two zeros in $(0, 1)$. In addition the $q_n(t)$ can be chosen in such a way that each is of bounded variation in $[0, 1]$ with

$$\int_0^1 |dq_n(t)| dt \leq k \left(\frac{1}{n}\right)$$

for a suitable k .

Construction of example. Define

$$\sigma_1 = 0, \quad \sigma_n = \sum_{k=1}^{n-1} \frac{1}{k^2}, \quad n > 1 .$$

Note that $\lim_{n \rightarrow \infty} \sigma_n = \pi^2/6$. Now $f(t)$ and $y(t)$ (which will be the solution of (1)) are defined on $[0, \pi^2/6)$ as follows,

$$\begin{aligned} f(t) &= C^2 + q_n(n^2(t - \sigma_n))\sigma_n \leq t \leq \sigma_{n+1} \\ y(t) &= (1/n^{4/(1-\gamma)})U_n(n^2(t - \sigma_n))\sigma_n \leq t \leq \sigma_{n+1} . \end{aligned}$$

Define $f(t) = C^2$ for $t \geq \pi^2/6$. Then, as in Coffman and Ullrich [2], it follows readily that $f(t)$ is positive and continuous on $[0, \infty)$ and that $y(t)$ is a solution of (1) on $[0, \pi^2/6)$. Continue $y(t)$ to $[0, \infty)$ by Lemma 1. Since $y(\sigma_n) = (1/n^{4^{1-\gamma}})$ it follows that $y(\pi^2/6) = y'(\pi^2/6) = 0$.

3. In this section we will show that $f(t)$ can have isolated zeros and continuation and uniqueness will still hold. First of all we state two nonoscillation theorems which will be needed.

THEOREM A ([1]). *Suppose that $1 < \gamma$ and that $f(t) > 0, f'(t) \leq 0, 0 \leq t < \infty$. If*

$$\int_0^\infty s^\gamma q(s) ds < \infty$$

then all nontrivial solutions of (1) are nonoscillatory (i.e., have a finite number of zeros on $[0, \infty)$).

THEOREM B ([5]). *Suppose that $0 < \gamma < 1$ and that $f(t) > 0, f'(t) \leq 0, 0 \leq t < \infty$. If*

$$\int_0^\infty sq(s) ds < \infty$$

then all nontrivial solutions of (1) are nonoscillatory.

THEOREM 2. *Let $\gamma > 1$. Suppose that on the interval $[0, \infty)$ $f(t)$ is continuous, locally of bounded variation, and positive except at a sequence of isolated points $\{t_i\}, i = 1, 2, 3, \dots$. If $f(t)$ is differentiable in a left neighborhood $(t_i - \varepsilon, t_i)$ of each t_i (ε depends on i) and if for each i the function*

$$A(x) = \frac{f\left(t_i - \frac{1}{x}\right)}{x^{\gamma+3}}$$

satisfies $A'(x) \leq 0$ for large x , then any solution of (1) existing at some $t_0 \geq 0$ can be continued to $[t_0, \infty)$. Likewise if $f(t)$ is differentiable in a right neighborhood $(t_i, t_i + \varepsilon)$ of each t_i (ε depends on i) and if for each i the function

$$A(x) = \frac{f\left(t_i + \frac{1}{x}\right)}{x^{\gamma+3}}$$

satisfies $A'(x) \leq 0$ for large x , then any solution of (1) existing at some $t_0 \geq 0$ can be continued to $[0, t_0]$.

Proof. Suppose that $y_i(t)$ is defined at t_0 . Then by Coffman and Ullrich's theorem $y_i(t)$ exists on $[t_0, t_i)$ where t_i is the first zero of $f(t)$ to the right of t_0 . Suppose $y_i(t)$ can't be continued to t_i . Then $y_i(t)$ has an infinite number of zeros on $[t_0, t_i)$ clustering at t_i . Let

$$x = \frac{1}{t_i - t} \quad y(t) = \frac{w(x)}{x}.$$

Then (1) is transformed into

$$(3) \quad w'' + A(x)w' = 0$$

where

$$A(x) = \frac{f\left(t_i - \frac{1}{x}\right)}{x^{\gamma+3}}.$$

$y_i(t)$ is transformed into a solution $w_1(x)$ of (3). $w_1(x)$ exists on $[1/(t_i - t_0), \infty)$ and is oscillatory (i.e., has arbitrarily large zeros). Also $A(x)$ satisfies

$$\int_0^\infty s^{\gamma+2-\varepsilon} A(s) ds < \infty$$

for any $\varepsilon > 0$. Since $A'(x) \leq 0$, Theorem A above says that $w_1(x)$ is nonoscillatory. This contradiction shows that $y_i(t)$ can be continued to t_i . Therefore, by standard existence theorem, $y_i(t)$ can be continued to a neighborhood of t_i . By Coffman and Ullrich's theorem $y_i(t)$ can be continued up to the next zero of $f(t)$. Since the zeros of $f(t)$ are isolated $y_i(t)$ can be continued to $[t_0, \infty)$.

The second part of the theorem follows in a similar manner by making the transformation

$$x = \frac{1}{t - t_i}, \quad y(t) = \frac{w(x)}{x}.$$

REMARK. The transformation in the preceding theorem has been used previously by Kiguradze [7].

REMARK. Kiguradze [7] has shown that if $f(t) < 0$ on some interval (t_1, t_2) then (1), with $\gamma > 1$, has a solution which can't be continued to the right of t_2 . Therefore a necessary condition for continuation of all solutions is $f(t) \geq 0$. Whether or not $f(t) \geq 0$ is sufficient (together with locally of bounded variation) is an open question.

THEOREM 3. Let $0 < \gamma < 1$. Suppose that $f(t)$ is class C' on $[0, \infty)$ and positive except for a sequence of isolated points $\{t_i\}$, $i =$

1, 2, 3, \dots . Suppose that for each i the function

$$A(x) = \frac{f\left(t_i - \frac{1}{x}\right)}{|x|^{\gamma+3}}$$

satisfies $A'(x) \leq 0$ for large $|x|$. Then the zero solution is unique on $[0, \infty)$.

Proof. Suppose there is a singular solution $y_1(t)$. Then $y_1(t)$ has an infinite number of zeros clustering at some point t^* . By Theorem 1, $t^* = t_i$ for some i . If the zeros of $y_1(t)$ cluster at t_i from the left (i.e., on $(t_i - \varepsilon, t_i)$) use the first transformation in the proof of Theorem 2. If the zeros of $y_1(t)$ cluster at t_i from the right use the second transformation in the proof of Theorem 2. Thus $y_1(t)$ is transformed into an oscillatory solution $w_1(x)$ of

$$w'' + A(x)w^\gamma = 0.$$

Since $A'(x) \leq 0$, this is a contradiction to Theorem B above.

REMARK. By using theorems of Gollwitzer [3], Theorems 2 and 3 can be improved somewhat in the sense that the condition $A'(x) \leq 0$ can be slightly relaxed.

4. Finally, we will answer a conjecture made in [5] about Theorem B and a similar question for Theorem A. The question is, can the condition $f(t) > 0, f'(t) \leq 0$ in Theorems A and B be replaced by the weaker condition $f(t) \geq 0$. The answer is no in both cases.

EXAMPLE 2. Coffman and Ullrich have shown that there is a continuous function $f(t) > 0$ defined on $[0, \infty)$ and a solution $\phi(t)$ of (1) such that $\phi(t)$ exists only on a finite interval $[t_0, t_1)$ and has an infinity of zeros clustering at t_1 . Using the transformation of the preceding section $\phi(t)$ is transformed into a solution $w(t)$ of

$$w'' + A(x)w^\gamma = 0$$

where

$$A(x) = \frac{f\left(t_1 - \frac{1}{x}\right)}{x^{\gamma+3}}, \quad x \geq \frac{1}{(t_1 - t_0)}$$

and $w(x)$ is oscillatory on $[1/(t_1 - t_0), \infty)$. Also

$$\int_{1/(t_1 - t_0)}^{\infty} x^\gamma A(x) dx < \infty.$$

This shows that the condition $f'(t) \leq 0$ can't be removed in Theorem A.

EXAMPLE 3. Similarly, by Example 1 above, let $f(t)$ be a positive and continuous function on $[0, \infty)$ and $\psi(t)$ a solution of (1) for $0 < \gamma < 1$ such that $\psi(t)$ is singular, $\psi(t_1) = \psi'(t_1) = 0$, and $\psi(t)$ has an infinity of zeros on some interval $[t_0, t_1)$ clustering at t_1 . Then $\psi(t)$ is transformed by the above transformation to $w(x)$ which is a solution of

$$w'' + A(x)w^\gamma = 0, \quad x \geq 1/(t_1 - t_0)$$

and $w(x)$ is oscillatory on $[1/(t_1 - t_0), \infty)$. Since

$$\int_1^\infty xA(x)dx < \infty$$

it is clear that $f'(t) \leq 0$ can't be removed from Theorem B.

REMARK. Again, results of Gollwitzer [3] show that the condition $f'(t) \leq 0$ can be weakened slightly.

REFERENCES

1. F. V. Atkinson, *On second-order non-linear oscillations*, Pacific J. Math. **5** (1955), 643-647.
2. C. V. Coffman and D. F. Ullrich, *On the continuation of solutions of a certain non-linear differential equation*, Monatsh. Math. **71** (1967), 385-392.
3. H. E. Gollwitzer, *Nonoscillation theorems for nonlinear second order equations* (to appear)
4. Philip Hartman, *Ordinary differential equations*, John Wiley and Sons, Inc., New York, 1964.
5. J. W. Heidel, *A nonoscillation theorem for a nonlinear second order differential equation*, Proc. Amer. Math. Soc. **22** (1969), 2, 485-488.
6. I. T. Kiguradze, *The problem of oscillation of solutions of nonlinear differential equations* (Russian), Differencial'nye Uravenija **1** (1965), 995-1006 (translated as Differential Equations) **1** (1965), 773-782.
7. ———, *Remarks on the oscillation of solutions of the equation $u'' + a(t) |u|^n \operatorname{sgn} u = 0$* (Russian), Časopis Psět. Mat. **92** (1967), 343-350.

Received March 18, 1969.

THE UNIVERSITY OF TENNESSEE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

BASIL GORDON*
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

* Acting Managing Editor.

Pacific Journal of Mathematics

Vol. 32, No. 3

March, 1970

Shair Ahmad, <i>Dynamical systems of characteristic 0^+</i>	561
Charles A. Akemann and Bernard Russo, <i>Geometry of the unit sphere of a C^*-algebra and its dual</i>	575
Philip Bacon, <i>The compactness of countably compact spaces</i>	587
Richard Blaine Barrar and Henry Loeb, <i>On the continuity of the nonlinear Tschebyscheff operator</i>	593
L. Carlitz, <i>Factorization of a special polynomial over a finite field</i>	603
Joe Ebeling Cude, <i>Compact integral domains</i>	615
Frank Rimi DeMeyer, <i>On automorphisms of separable algebras. II</i>	621
James B. Derr, <i>Generalized Sylow tower groups</i>	633
Raouf Doss, <i>Some inclusions in multipliers</i>	643
Mary Rodriguez Embry, <i>The numerical range of an operator</i>	647
John Froese, <i>Domain-perturbed problems for ordinary linear differential operators</i>	651
Zdeněk Frolík, <i>Absolute Borel and Souslin sets</i>	663
Ronald Owen Fulp, <i>Tensor and torsion products of semigroups</i>	685
George Grätzer and J. Plonka, <i>On the number of polynomials of an idempotent algebra. I</i>	697
Newcomb Greenleaf and Walter Read, <i>Positive holomorphic differentials on Klein surfaces</i>	711
John Willard Heidel, <i>Uniqueness, continuation, and nonoscillation for a second order nonlinear differential equation</i>	715
Leon A. Henkin, <i>Extending Boolean operations</i>	723
R. Hirshon, <i>On hopfian groups</i>	753
Melvin Hochster, <i>Totally integrally closed rings and extremal spaces</i>	767
R. Mohanty and B. K. Ray, <i>On the convergence of a trigonometric integral</i>	781
Michael Rich, <i>On a class of nodal algebras</i>	787
Emile B. Roth, <i>Conjugate space representations of Banach spaces</i>	793
Rolf Schneider, <i>On the projections of a convex polytope</i>	799
Bertram Manuel Schreiber, <i>On the coset ring and strong Ditkin sets</i>	805
Edgar Lee Stout, <i>Some remarks on varieties in polydiscs and bounded holomorphic functions</i>	813
James Edward Ward, <i>Two-groups and Jordan algebras</i>	821