

Pacific Journal of Mathematics

PROJECTIVE DISTRIBUTIVE LATTICES

RAYMOND BALBES AND ALFRED HORN

PROJECTIVE DISTRIBUTIVE LATTICES

RAYMOND BALBES AND ALFRED HORN

It will be shown that a countable distributive lattice is projective if and only if the product of any two join irreducible elements is join irreducible, and every element of the lattice is both a finite sum of join irreducible elements and a finite product of meet irreducible elements. For an arbitrary distributive lattice, necessary and sufficient conditions for projectivity are obtained by adding to these conditions a further condition on the set of join irreducible elements.

1. **Definitions.** We use sum and product notation for least upper bounds and greatest lower bounds. If A and B are meet semi-lattices, then a meet homomorphism $f: A \rightarrow B$ is a function such that $f(xy) = f(x)f(y)$. An element x of a lattice is called join irreducible if $x = y + z$ implies $x = y$ or $x = z$. x is called sub join irreducible if $x \leq y + z$ implies $x \leq y$ or $x \leq z$. In a distributive lattice these two notions coincide. We define meet irreducible and super meet irreducible in a dual manner. A lattice is called conditionally implicative if whenever $x \not\leq y$ there exists a largest z such that $xz \leq y$. The smallest and largest element of a lattice are denoted by 0 and 1 respectively. The cardinal of a set S is denoted by $|S|$. For definitions of projective distributive lattice and retract, see [1]. Note that epimorphisms are understood to be homomorphisms which are onto. If the term epimorphism is used as in the general theory of categories, there are no projective distributive lattices.

2. **Projective distributive lattices.** Consider the following properties of a lattice.

(P1) Every element is a sum of finitely many sub join irreducible elements.

(P2) Every element is a product of finitely many super meet irreducible elements.

(P3) The product of any two sub join irreducible elements is sub join irreducible.

(P4) The sum of any two super meet irreducible elements is super meet irreducible.

(P5) The lattice is conditionally implicative.

(P6) The lattice is dually conditionally implicative.

THEOREM 1. *Suppose A and B are lattices and A is a retract of B . Then we have:*

- (i) If B satisfies (P1), then A satisfies (P1).
- (ii) If B satisfies (P1) and (P3), then A satisfies (P4).
- (iii) If B satisfies (P5), then A satisfies (P5).

Proof. By hypothesis, there exist homomorphisms $f: B \rightarrow A$ and $g: A \rightarrow B$ such that $fg = I_A$. Suppose B satisfies (P1). Let x be any element of A . Then $g(x) = \sum(S)$, where S is a finite nonempty set of sub join irreducible elements of B . Let T be the set of maximal elements of the set $f(S)$. Then $x = \sum(T)$, and we claim that every element a of T is sub join irreducible. Suppose $a \leq u + v$ but $a \not\leq u$ and $a \not\leq v$. We have $a = f(b)$ for some $b \in S$. Then

$$b \leq g(x) \leq g(u) + g(v) + \sum(g(T - \{a\})).$$

Hence $b \leq g(u)$, or $b \leq g(v)$, or $b \leq g(c)$ for some $c \in T - \{a\}$. Applying f , we find $a \leq u$, or $a \leq v$, or $a \leq c$. This contradicts the maximality of a .

Suppose B satisfies (P1) and (P3). Let a_1 and a_2 be super meet irreducible in A . Suppose $a_1 + a_2 \geq a_3 a_4$ but $a_1 + a_2 \not\geq a_3$ and $a_1 + a_2 \not\geq a_4$. We have $g(a_3) = \sum(S)$ and $g(a_4) = \sum(T)$, where S and T are finite sets of sub join irreducible elements of B . Hence $a_3 = \sum(f(S))$ and $a_4 = \sum(f(T))$. There exists $x \in S$ and $y \in T$ such that $f(x) \not\leq a_1 + a_2$ and $f(y) \not\leq a_1 + a_2$. Now $xy \leq g(a_3)g(a_4) \leq g(a_1) + g(a_2)$. By (P3), either $xy \leq g(a_1)$ or $xy \leq g(a_2)$. Therefore $f(x)f(y) \leq a_1$ or $f(x)f(y) \leq a_2$. Since a_1 and a_2 are super meet irreducible, we have $f(x) \leq a_1 + a_2$ or $f(y) \leq a_1 + a_2$, which is a contradiction.

The proof of (iii) is given in [2, Th. 2.9].

THEOREM 2. For any lattice A , we have:

- (i) If (P1) and (P3), then (P4).
- (ii) If (P2) then (P5).
- (iii) If (P1), (P2) and (P3), then (P4), (P5) and (P6).
- (iv) If A is a retract of B , and B satisfies (P1), (P2) and (P3), then A satisfies all six properties (P1)-(P6).

Proof. (i) follows immediately from Theorem 1 (ii). Suppose A satisfies (P2). Let $x, y \in A$ and $x \not\leq y$. We have $y = \pi(S)$, where S is a finite set of super meet irreducible elements. Let

$$T = \{a \in S: x \not\leq a\},$$

and let $z = \pi(T)$. If $a \in S - T$, then $a \geq x \geq xz$. If $a \in T$, then $a \geq z \geq xz$. Hence $xz \leq \pi(S) = y$. Now suppose $xw \leq y$. Then for each $a \in T$, we have $a \geq xw$, hence $a \geq w$. Therefore $w \leq \pi(T) = z$. This proves (ii). (iii) follows from (i), (ii), and the dual of (ii).

Suppose B satisfies (P1), (P2), (P3) and A is a retract of B . Then by Theorem 1 (i) and its dual and by Theorem 1 (ii), A satisfies (P1), (P2) and (P4). (iv) now follows from the dual of (iii).

THEOREM 3. *Let A be a projective distributive lattice. Then A satisfies all six properties (P1)–(P6).*

Proof. A is a retract of a free distributive lattice F . It is well known that F satisfies (P1), (P2) and (P3), the sub join irreducible elements being the products of free generators. The result now follows from Theorem 2 (iv).

THEOREM 4. *Let A be a distributive lattice satisfying (P1), and let J be the set of join irreducible elements of A . Then any order preserving map $h: J \rightarrow B$, where B is a distributive lattice, can be extended uniquely to a join homomorphism $\hat{h}: A \rightarrow B$. If in addition, A satisfies (P3) and h is a meet homomorphism, then \hat{h} is a homomorphism.*

Proof. If $x = x_1 + \dots + x_n$, where $x_i \in J$ for all i , let

$$\hat{h}(x) = h(x_1) + \dots + h(x_n).$$

\hat{h} is well defined because $x_1 + \dots + x_n \leq y_1 + \dots + y_m$ implies each x_i is \leq some y_j . It is obvious that \hat{h} is a join homomorphism and is unique. If h is a meet homomorphism, then it is easy to see that \hat{h} also preserves products.

THEOREM 5. *Let J be any meet semi-lattice. There exists a unique distributive lattice \hat{J} such that \hat{J} satisfies (P1) and (P3), and J is the set of join irreducible elements of \hat{J} . \hat{J} is unique up to isomorphism over J .*

Proof. Let \hat{J} be the set of all finite unions of principal ideals of J . \hat{J} is a ring of sets and the map $f: J \rightarrow \hat{J}$ such that $f(x) = \{y \in J: y \leq x\}$ is a meet-monomorphism. It is clear that $f(J)$ is the set of all join irreducible elements of \hat{J} . The uniqueness of \hat{J} follows easily from Theorem 4.

In view of Theorems 3, 4 and 5, the study of projective distributive lattices can be reduced to the question: for which meet semi-lattices J is \hat{J} projective? An obvious condition is the following.

THEOREM 6. *Let A be a distributive lattice which satisfies (P1) (P3), and let J be the set of join irreducible elements of A . Then*

A is projective if and only if there exists a free distributive lattice F , a homomorphism $f: F \rightarrow A$ and a meet homomorphism $h: J \rightarrow F$ such that $fh(x) = x$ for all $x \in J$.

Proof. If A is projective, it is a retract of a free distributive lattice F . The necessity of the condition follows immediately. Conversely, given f and h , by Theorem 4, h can be extended to a homomorphism $\hat{h}: A \rightarrow F$. It is easy to see that $f\hat{h} = I_A$, and therefore A is projective.

Consider the following weakening of the condition of Theorem 6:

(P7) There exists a free distributive lattice F , a homomorphism $f: F \rightarrow A$ and an order preserving map $h: J \rightarrow F$ such that $fh(x) = x$ for all $x \in J$.

THEOREM 7. *Let A be a distributive lattice and let J be the set of join irreducible elements of A . Then A is projective if and only if A satisfies (P1), (P2), (P3) and (P7).*

Proof. The necessity of the conditions follows from Theorems 3 and 6. Suppose A satisfies the conditions. Let $f: F \rightarrow A$ and $h: J \rightarrow F$ be as in (P7). By Theorem 4, h can be extended to a join homomorphism $\hat{h}: A \rightarrow F$. It is clear that $f\hat{h} = I_A$. For each $x \in J$, there exists a finite set $S(x)$ of meet irreducible elements of A such that $x = \pi(S(x))$. Define $g: J \rightarrow F$ as follows: $g(x) = \pi(\hat{h}(S(x)))$. For any $x \in J$, $fg(x) = \pi(f\hat{h}(S(x))) = \pi(S(x)) = x$. If $x, y \in J$, $x \leq y$ and $z \in S(y)$, then $\pi(S(x)) \leq z$. Since z is super meet irreducible, z must be \geq some member of $S(x)$. Hence every member of $\hat{h}(S(y))$ is \geq some member of $\hat{h}(S(x))$. Thus $g(y) \geq g(x)$ and we have shown that g preserves order. The proof will be complete by Theorem 6 if we show that g preserves products. Suppose $x, y, z \in J$ and $xy = z$. Then $\pi(S(z)) = \pi(S(x) \cup S(y))$, and therefore each member of $S(z)$ is \geq some member of $S(x) \cup S(y)$. Therefore

$$g(z) = \pi(\hat{h}(S(z))) \geq \pi(\hat{h}(S(x) \cup S(y))) = g(x)g(y).$$

Since g preserves order, we have $g(z) \leq g(x)g(y)$, and the proof is complete.

THEOREM 8. *Let A be a countable distributive lattice and let J be the set of join irreducible elements of A . If A satisfies (P1) and (P3) then A satisfies (P7).*

Proof. There exists an epimorphism $f: F \rightarrow A$, where F is a free distributive lattice. For each $x \in J$, select an element $g(x)$ such

that $fg(x) = x$. Arrange the members of J in sequence x_0, x_1, \dots . Define $h: J \rightarrow F$ inductively as follows: $h(x_0) = g(x_0)$, and

$$h(x_n) = g(x_n) \amalg \{h(x_i): i < n, x_i > x_n\} + \sum \{h(x_i): i < n, x_i < x_n\}.$$

By induction, it is easy to see that $fh(x_n) = x_n$ and h preserves order on the set $\{x_0, \dots, x_n\}$. This proves (P7).

THEOREM 9. *If A is a countable distributive lattice, then A is projective if and only if A satisfies (P1), (P2) and (P3).*

Proof. This follows from Theorems 7 and 8.

COROLLARY ([1, Th. 7.1]). *If A is finite, then A is projective if and only if A satisfies (P3).*

Proof. Every finite distributive lattice satisfies (P1) and (P2).

Theorem 7 suggests the following question: for which semi-lattices J does $S = \hat{J}$ satisfy (P7)? Theorem 8 states that countability is a sufficient condition. Another sufficient condition is that J be projective in the category of semi-lattices. Condition (P7) may be replaced by a condition which refers more explicitly to J itself. First, if A is projective, every epimorphism $f: F \rightarrow A$ has a right inverse. Therefore in Theorem 7, we may replace (P7) by

(P8) There exists a free distributive lattice F whose set of free generators is T , a homomorphism $f: F \rightarrow A$ such that $f(T) = J$, and an order preserving map $h: J \rightarrow F$ such that $fh(x) = x$ for all $x \in J$.

THEOREM 10. *Let J be a meet semi-lattice. Then $A = \hat{J}$ satisfies (P8) if and only if for each $x \in J$ there exists a finite sequence $S_{x,0}, \dots, S_{x,p(x)}$ of nonempty finite subsets of J such that*

- (i) $\pi(S_{x,0}) = x$.
- (ii) $\pi(S_{x,j}) \leq x$ for all j .
- (iii) if $x \leq y$, there for every j there is a k such that $S_{x,j} \supseteq S_{y,k}$.

Proof. Assume (P8) holds. Let $x \in J$. Then

$$h(x) = \pi(T_{x,0}) + \dots + \pi(T_{x,p(x)}),$$

where each $T_{x,j}$ is a finite subset of T . Let $S_{x,j} = f(T_{x,j})$. Then $x = \pi(S_{x,0}) + \dots + \pi(S_{x,p(x)})$. Since x is join irreducible, we have (i) and (ii) after renumbering indices. If $x \leq y$, then $h(x) \leq h(y)$. From

this it follows that every $T_{x,j}$ contains some $T_{y,k}$ (see [1, Lemma 4.5]). This proves (iii).

Assume (i), (ii) and (iii). Let F be a free distributive lattice with a free generating set T with the same cardinal as J . There exists a homomorphism $f: F \rightarrow A$ which maps T onto J in a one-to-one way. For each $S_{x,j}$ let $T_{x,j}$ be the subset of T such that $f(T_{x,j}) = S_{x,j}$. Define $h: J \rightarrow F$ as follows:

$$h(x) = \pi(T_{x,0}) + \cdots + \pi(T_{x,p(x)}).$$

By (i) and (ii), $fh(x) = x$ for all $x \in J$, and by (iii) h is order preserving. This completes the proof.

If P is a partially ordered set, there exists a distributive lattice P^* containing P such that P generates P^* and every order preserving map from P to a distributive lattice B can be extended to a homomorphism from P^* to B . (See for example [2, Definition 1.10].) In Lemma 3.8 of [2] it was shown that P^* is projective if and only if for each $x \in P$ there exists a finite sequence $S_{x,0}, \dots, S_{x,p(x)}$ of non-empty finite subsets of P such that

- (i) $x \in S_{x,0}$ and every member of $S_{x,0}$ is $\geq x$.
- (ii) for each j , $S_{x,j}$ has a member $\leq x$
- (iii) if $x \leq y$, then for every j there is a k such that $S_{x,j} \supseteq S_{y,k}$.

Comparing with the conditions of Theorem 9, we find the following.

THEOREM 11. *If J is meet semi-lattice and J^* is projective, then \hat{J} satisfies (P8).*

A sufficient condition for the projectivity of J^* is given in [2, Th. 3.12].

3. Direct products.

LEMMA 1. *Let A_1 and A_2 be projective distributive lattices. If A_1 and A_2 have a 0 and 1, then $A_1 \times A_2$ is projective.*

Proof. We may assume $|A_i| > 1$ for $i = 1, 2$. Let F be a free distributive lattice with the free generating set $T_1 \cup T_2$, where T_1 and T_2 are disjoint and $|T_i| = |A_i|$, $i = 1, 2$. There exists an epimorphism $f: F \rightarrow A_1 \times A_2$ such that $f(T_1) = A_1 \times \{0\}$ and $f(T_2) = \{0\} \times A_2$. Let F_i be the sublattice of F generated by T_i . Then $f(F_1) = A_1 \times \{0\}$ and $f(F_2) = \{0\} \times A_2$. Define $f_i: F_i \rightarrow A_i$ by $f_i = \pi_i \cdot (f|F_i)$, where $\pi_i: A_1 \times A_2 \rightarrow A_i$ is the natural projection. Since A_i is projective, there exists a homomorphism $g_i: A_i \rightarrow F_i$ such that $f_i g_i = I_{A_i}$. Define $g: A_1 \times A_2 \rightarrow F$ by

$$g(x, y) = g_1(x) + g_2(y) + g_1(1)g_2(1).$$

Then $fg(x, y) = (x, 0) + (0, y) + (1, 0) \cdot (0, 1) = (x, y)$, and g is a homomorphism. Therefore $A_1 \times A_2$ is a retract of F , and the proof is complete.

LEMMA 2. *An element x of a direct product $\prod_{i \in I} A_i$ of distributive lattices is join irreducible if and only if for some $i \in I$, we have:*

- (i) $x(j) = 0$ for all $j \neq i$.
- (ii) $x(i)$ is join irreducible.

Proof. The proof is easy and will be omitted.

THEOREM 12. *Let $\langle A_i: i \in I \rangle$ be a family of distributive lattices. Suppose $|I| > 1$ and $|A_i| > 1$ for all i . Then $\prod_{i \in I} A_i$ is projective if and only if*

- (i) A_i is projective for each i
- (ii) I is finite, and
- (iii) each A_i has a 0 and 1.

Proof. The sufficiency follows from Lemma 1. Suppose A is projective. By hypothesis there exists $x \in A$ and $i_1, i_2 \in I$ such that $i_1 \neq i_2$, $x(i_1) \neq 0$ and $x(i_2) \neq 0$. Since x is a sum of join irreducible elements, it follows from Lemma 2 that A_i has a 0 for all $i \in I$. By duality, each A_i has a 1, which proves (iii). Finally if I is infinite, then the 1 element of A cannot be a finite sum of join irreducible elements by Lemma 2, since $|A_i| > 1$ for all i .

REFERENCES

1. R. Balbes, *Projective and injective distributive lattices*, Pacific J. Math. **21** (1967), 405-420.
2. R. Balbes and A. Horn, *Order sums of distributive lattices*, Pacific J. Math. **21** (1967), 421-435.

Received October 1, 1969. This research was supported in part by National Science Foundation grants GP-11893 and GP-9044.

UNIVERSITY OF MISSOURI, ST. LOUIS
UNIVERSITY OF CALIFORNIA, LOS ANGELES

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

BASIL GORDON*
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLE

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

* Acting Managing Editor.

Pacific Journal of Mathematics

Vol. 33, No. 2

April, 1970

Raymond Balbes and Alfred Horn, <i>Projective distributive lattices</i>	273
John Findley Berglund, <i>On extending almost periodic functions</i>	281
Günter Krause, <i>Admissible modules and a characterization of reduced left artinian rings</i>	291
Edward Milton Landesman and Alan Cecil Lazer, <i>Linear eigenvalues and a nonlinear boundary value problem</i>	311
Anthony To-Ming Lau, <i>Extremely amenable algebras</i>	329
Aldo Joram Lazar, <i>Sections and subsets of simplexes</i>	337
Vincent Mancuso, <i>Mesocompactness and related properties</i>	345
Edwin Leroy Marsden, Jr., <i>The commutator and solvability in a generalized orthomodular lattice</i>	357
Shozo Matsuura, <i>Bergman kernel functions and the three types of canonical domains</i>	363
S. Mukhoti, <i>Theorems on Cesàro summability of series</i>	385
Ngô Van Quê, <i>Classes de Chern et théorème de Gauss-Bonnet</i>	393
Ralph Tyrrell Rockafellar, <i>Generalized Hamiltonian equations for convex problems of Lagrange</i>	411
Ken-iti Sato, <i>On dispersive operators in Banach lattices</i>	429
Charles Andrew Swanson, <i>Comparison theorems for elliptic differential systems</i>	445
John Griggs Thompson, <i>Nonsolvable finite groups all of whose local subgroups are solvable. II</i>	451
David J. Winter, <i>Cartan subalgebras of a Lie algebra and its ideals</i>	537