

# Pacific Journal of Mathematics

**ON A CLASS OF NÖRLUND MEANS AND FOURIER SERIES**

H. P. DIKSHIT

## ON A CLASS OF NÖRLUND MEANS AND FOURIER SERIES

H. P. DIKSHIT

By considering a class of Nörlund means that covers as a subclass the corresponding  $(C)$  means, we obtain in the present paper, several results concerning absolute Nörlund summability and deduce from these the corresponding  $|C|$  results as special cases. What is indeed remarkable, is that a special case of our Theorem 2 improves an earlier result due to Bosanquet and Hyslop in dropping one of the two independent conditions used by them. Further, the proofs of some of our results are shorter and even more direct than the proofs given for the corresponding special cases by using equivalent Riesz means instead of  $(C)$  means.

1. Definitions and notations. Let  $\sum_n v_n$  be a given infinite series with the sequence of partial sum  $\{s_n\}$ . We shall consider sequence to sequence transformations of the type

$$t_n = \sum_{k=0}^{\infty} d_{nk} s_k; \quad d_{nk} = 0 \quad \text{for } k > n;$$

in which the elements of the matrix  $D = (d_{nk})$  are real or complex constants.  $t_n$  is called the  $n$ -th  $D$ -mean of  $\{s_n\}$ .

Let  $\{p_n\}$  be a sequence of constants, real or complex and let  $P_n = p_0 + p_1 + \dots + p_n \neq 0$ ,  $P_{-1} = p_{-1} = 0$ . Then the matrix  $D$  defines a Nörlund matrix  $(N, p)$ , if

$$d_{nk} = p_{n-k}/P_n, \quad n \geq k \geq 0.$$

In the special case in which

$$(1.1) \quad p_n = \binom{n + \alpha - 1}{\alpha - 1} = \frac{\Gamma(n + \alpha)}{\Gamma(n + 1)\Gamma(\alpha)}, \quad \alpha \neq -1, -2, \dots,$$

the  $(N, p)$  mean reduces to the familiar  $(C, \alpha)$  mean.

The  $(N, p)(C, 1)$  matrix is defined as the product of a  $(N, p)$  matrix with the  $(C, 1)$  matrix. Thus the  $(N, p)(C, 1)$  mean of  $\{s_n\}$  is

$$t_n = \frac{1}{P_n} \sum_{r=0}^n p_{n-r} \frac{1}{r+1} \sum_{k=0}^r s_k.$$

Similarly, one defines the  $(C, 1)(N, p)$  mean [5].

Let  $f(t)$  be integrable  $(L)$  in  $(-\pi, \pi)$  and periodic with period  $2\pi$ . We assume as we may without any loss of generality that

$$f(t) \sim \sum_{n=1}^{\infty} (a_n \cos nt + b_n \sin nt) = \sum_n A_n(t).$$

Then the conjugate series is

$$\sum_{n=1}^{\infty} (b_n \cos nt - a_n \sin nt) = \sum_n B_n(t).$$

We shall also consider the series

$$\sum_n \frac{1}{n} \left\{ \sum_{k=1}^n A_k(t) - s \right\} = \sum_n A_n^*(t),$$

where  $s$  is an appropriate number, independent of  $n$ .

Throughout the present paper we write  $L^\alpha(t)$  for the series  $\sum_n n^\alpha A_n(t)$ ,  $\tilde{L}^\alpha(t)$  for  $\sum_n n^\alpha B_n(t)$ , ( $\alpha \geq 0$ ),  $L^*(t)$  for  $\sum_n A_n^*(t)$  and  $\mathcal{A}$  for the class  $\{L^\alpha(t), \tilde{L}^\alpha(t), L^*(t)\}$ .

Let  $E_f$  be a point set in the interval  $(-\pi, \pi)$  for each function  $f(t)$  and such that at every point  $x \in E_f$ ,  $f(x)$  has a finite definite value and satisfies a prescribed condition of regularity.

**DEFINITION 1.** A method of summation  $D = \langle\langle d_{nk} \rangle\rangle$  is said to be  $|A(x), E_f|$ -effective, if for each  $x \in E_f$

$$\sum_{n=1}^{\infty} |t_n(A(x)) - t_{n-1}(A(x))| < \infty,$$

symbolically,  $\{t_n(A(x))\} \in BV$ ; where  $t_n(A(x))$  denotes the  $n$ th  $D$ -mean of  $A(x) \in \mathcal{A}$ .

We write

$$\phi(t) = \frac{1}{2} \{f(x+t) + f(x-t)\}; \quad \phi^*(t) = \phi(t) - s;$$

$$\Phi_\alpha(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-u)^{\alpha-1} \phi(u) du, \quad \alpha > 0; \quad \Phi_0(t) = \phi(t);$$

$$\phi_\alpha(t) = \Gamma(\alpha+1) t^{-\alpha} \Phi_\alpha(t), \quad \alpha \geq 0;$$

$$\psi(t) = \frac{1}{2} \{f(x+t) - f(x-t)\},$$

$\Psi_\alpha(t)$  and  $\psi_\alpha(t)$  have similar meanings.

$[x]$  denotes the greatest integer not greater than  $x$ .

By ' $F(t) \in BV(\alpha, b)$ ', we mean that  $F(t)$  is a function of bounded variation in  $(\alpha, b)$  and by ' $\{\lambda_n\} \in B$ ' that  $\{\lambda_n\}$  is a bounded sequence.

$K$  denotes a positive constant, not necessarily the same at each occurrence.

DEFINITION 2. For some  $\alpha \geq 0$ , the point  $x$  is said to be

- (i)  $|F_\alpha|$  - regular, if  $\phi_\alpha(t) \in BV(0, \pi)$ ,
- (ii)  $|\tilde{F}_\alpha|$  - regular, if  $\Psi_\alpha(+0) = 0$  and  $\int_0^\pi t^{-\alpha} |d\Psi_\alpha(t)| \leq K$ ,
- (iii)  $|F^\alpha|$  - regular, if  $\int_0^\pi t^{-\alpha} |d\phi(t)| \leq K$ ,
- (iv)  $|\tilde{F}^\alpha|$  - regular, if  $\psi(+0) = 0$  and  $\int_0^\pi t^{-\alpha} |d\psi(t)| \leq K$ ,
- (v)  $|F^*|$  - regular, if  $\int_0^\pi t^{-1} |d\phi^*(t)| \leq K$ ,
- (vi)  $|\tilde{F}^*|$  - regular, if  $\int_0^\pi t^{-1} |\psi(t)| dt \leq K$ .

Denoting the set of  $|X|$  - regular points with respect to  $f(t)$  in  $(-\pi, \pi)$  by  $E|X, f|$ , we know the following ([14], § 13.24)

$$(1.2) \quad E|\tilde{F}^*, f| \not\subset E|\tilde{F}_0, f| \quad \text{and} \quad E|\tilde{F}_0, f| \not\subset E|\tilde{F}^*, f|^1.$$

DEFINITION 3. A method of summation, which is  $|A(x), E_f|$  - effective is said to be

- (i)  $|F_\alpha|$  - effective, if  $A(x) = L^0(x)$  and  $E_f = E|F_\alpha, f|$ ;
- (ii)  $|\tilde{F}_\alpha|$  - effective, if  $A(x) = \tilde{L}^0(x)$  and  $E_f = E|\tilde{F}_\alpha, f|$ ;
- (iii)  $|\tilde{F}^\alpha|$  - effective, if  $A(x) = L^\alpha(x)$  and  $E_f = E|F^\alpha, f|$ ;
- (iv)  $|\tilde{F}^\alpha|$  - effective, if  $A(x) = \tilde{L}^\alpha(x)$  and  $E_f = E|\tilde{F}^\alpha, f|$ ;
- (v)  $|F^*|$  - effective, if  $A(x) = L^*(x)$  and  $E_f = E|F^*, f|$ ;
- (vi) absolute  $\alpha$ -effective or  $|\alpha|$ -effective, if it is effective in the sense of (i)-(iv) simultaneously.

The following notations will be used throughout. If for  $n = 0, 1, 2, \dots$

$$p_n > 0, p_{n+1}/p_n \leq p_{n+2}/p_{n+1} \leq 1,$$

then we shall write  $\{p_n\} \in M$ . If  $\{p_n\} \in M$  and for some  $\alpha$ ,

$$(1.3) \quad P_k \sum_{n=k}^\infty \frac{1}{n^{1-\alpha} P_n} \leq Kk^\alpha, \quad k = 1, 2, \dots$$

then we write  $\{p_n\} \in M_\alpha$ .

For a given series  $v = \sum_n v_n$ ,

$$\sigma_n(v) = \sum_{k=0}^n p_{n-k} k v_k.$$

We also write

---

<sup>1</sup> I.e.,  $x \in E|\tilde{F}^*, f|$  does not imply that  $x \in E|\tilde{F}_0, f|$  and conversely.

$$h(n, t) = \frac{2}{\pi} \sum_{k=0}^n p_{n-k} \exp(ikt),$$

$$H(n, u) = \frac{1}{\Gamma(1-\alpha)} \int_u^\pi (t-u)^{-\alpha} \frac{d}{dt} h(n, t) dt,$$

$g(n, t)$  and  $\tilde{g}(n, t)$  for the imaginary and real parts respectively of  $h(n, t)$ .  $J(n, u)$  (or  $\tilde{J}(n, u)$ ) is  $H(n, u)$  with  $h(n, t)$  replaced by  $g(n, t)$  (or  $\tilde{g}(n, t)$ ). Further

$$V(n, u) = \frac{1}{\Gamma(1+\alpha)} \int_0^u v^\alpha \frac{d}{dv} J(n, v) dv.$$

**2. Introduction.** Concerning  $|\alpha|$ -effectiveness of the (C)-method we have the following result which is known to be the best possible in the sense that it breaks down if  $\delta = 0$ .

**THEOREM A.** *If  $0 < \alpha < 1$ , then  $(C, \alpha + \delta)$  for any  $\delta > 0$  is  $|\alpha|$ -effective.*

Starting with the proof of  $|F_\alpha|$ -effectiveness of  $(C, \alpha + \delta)$  given by Bosanquet [1] in 1936, Theorem A has been completed in stages by different authors. Thus  $|\tilde{F}_\alpha|$ -effective part of Theorem A is due to Bosanquet and Hyslop ([2], Th. 2) and  $|F^\alpha|$  and  $|\tilde{F}^\alpha|$ -effective parts are due to Mohanty ([11], Th. 1 and Th. 2). It is somewhat peculiar to observe that Theorem A may be extended to cover the case  $\alpha = 0$ , as far as  $|F_\alpha|$  or  $|F^\alpha|$ -effective parts are concerned but for  $|\tilde{F}_0|$  or equivalently  $|\tilde{F}^0|$ -effectiveness of the (C) method, Bosanquet and Hyslop ([2], Th. K, with  $\alpha = 0$ ) require an additional condition that  $x$  is  $|\tilde{F}^*|$ -regular also, in the following.

**THEOREM B.** *If  $x$  is  $|\tilde{F}^*|$ -regular, then the  $(C, \delta)$  method is  $|\tilde{F}_0|$ -effective for each  $\delta > 0$ .*

The condition that  $x$  is  $|\tilde{F}^*|$ -regular is independent of the condition that  $x$  is  $|\tilde{F}_0|$ -regular in view of (1.2).

In Theorem 1 of the present paper we prove  $|\alpha|$ -effectiveness ( $0 < \alpha < 1$ ) of a  $(N, p)$  method which covers the corresponding (C) method as a special subclass and deduce the various results of Theorem A as particular cases of our Theorem 1. What is indeed remarkable is that in Theorem 2, we have succeeded in extending Theorem 1 to the case  $\alpha = 0$  by proving  $|\tilde{F}_0|$ -effectiveness of the  $(N, p)$  method, even without using the hypothesis that  $x$  is a  $|\tilde{F}^*|$ -regular point. Thus the following special case of Theorem 2 improves Theorem B in dropping the condition that  $x$  is a  $|\tilde{F}^*|$ -

regular point.

THEOREM C. *The  $(C, \delta)$  method is  $|\tilde{F}_0|$  - effective for every  $\delta > 0$ .*

Covering as a special case, an earlier result due to Mohanty and Mohapatra ([12], Th. 3) we prove in Theorem 3,  $|F^*|$  - effectiveness of the  $(N, p)$  method and thus demonstrate that the  $|N, p|$  of  $L^*(x)$ , is a local property of its generating function [6].

It may be observed that the proofs of some of our theorems are shorter and even more direct than the proofs given in support of the corresponding special cases by using equivalent Riesz methods instead of the  $(C)$  - methods.

3. We prove the following.

THEOREM 1. *If  $0 < \alpha < 1$  and  $\{p_n\} \in M_\alpha$ , then  $(N, p)$  is  $|\alpha|$  - effective.*

THEOREM 2. *If  $\{p_n\} \in M_0$ , then  $(N, p)$  is  $|F_0|$  and  $|\tilde{F}_0|$  - effective.*

THEOREM 3. *If  $\{p_n\} \in M_0$ , then  $(N, p)$  is  $|F^*|$  - effective.*

4. Some preliminary results. We need the following lemmas, of which Lemma 1 is the same as Theorem 6 of Das [3].

LEMMA 1. *If  $\{p_n\} \in M$ , then a necessary and sufficient condition that  $\{t_n(v)\} \in BV$ , for a given series  $\sum_n v_n$  is that*

$$\sum_{n=1}^{\infty} \frac{1}{nP_n} |\sigma_n(v)| \leq K,$$

where  $t_n(v)$  is the  $n$ -th  $(N, p)$  mean of  $\sum_n v_n$ .

LEMMA 2. *If  $\{p_n\}$  is a nonnegative monotonic nonincreasing sequence, then for any  $n$  and  $0 \leq a \leq b$*

$$\left| \sum_{k=a}^b p_k \exp i(n-k)t \right| \leq KP_{[1/t]},$$

uniformly in  $0 < t \leq \pi$ .

Lemma 2 is given in McFadden [10].

LEMMA 3. *If  $\{p_n\}$  is a positive nonincreasing sequence, then*

$$|H(n, u)| = \begin{cases} O(n^\alpha P_n), & \text{for all } u, \\ O(n^\alpha P_{[1/u]}) , & \text{for } u \geq \frac{1}{n}. \end{cases}$$

*Proof.* We write

$$\begin{aligned} \Gamma(1 - \alpha)H(n, u) &= \left\{ \int_u^{u+(1/n)} + \int_{u+(1/n)}^\pi \right\} (t - u)^{-\alpha} \frac{d}{dt} h(n, t) dt \\ &= H_1 + H_2, \end{aligned}$$

say. If  $u \geq 1/n$ , then by Abel's Lemma and Lemma 2,

$$|H_1| \leq K \int_u^{u+(1/n)} (t - u)^{-\alpha} n P_{[1/t]} dt \leq K n^\alpha P_{[1/u]},$$

since  $\{P_n\}$  is nondecreasing. Next, we have

$$\begin{aligned} |H_2| &= \left| n^\alpha \int_{u+(1/n)}^\eta \frac{d}{dt} h(n, t) dt \right|, \quad u + \frac{1}{n} < \eta < \pi, \\ &\leq K n^\alpha P_{[1/u]}, \end{aligned}$$

by virtue of Lemma 2.

This proves the second part of the lemma. The other part follows by a similar reasoning when one observes that

$$\left| \sum_{k=0}^n p_{n-k} k \exp(ikt) \right| \leq K n P_n.$$

LEMMA 4. If  $\{p_n\}$  is a positive monotonic nonincreasing sequence, then

$$|V(n, u)| = \begin{cases} O(n^\alpha u^\alpha P_n) & \text{for all } u; \\ O(n^\alpha) + O(n^\alpha u^\alpha P_{[1/u]}) & \text{for } u \geq \frac{1}{n}. \end{cases}$$

For the proof of Lemma 4, reference may be made to ([9], p. 265).

5. Proof of Theorem 1. (I)  $|F_\alpha|$  - effectiveness: We have

$$nA_n(x) = \frac{2}{\pi} \int_0^\pi \phi(t) \frac{d}{dt} \sin ntdt$$

and

$$\sigma_n(L^0(x)) = \int_0^\pi \phi(t) \frac{d}{dt} g(n, t) dt.$$

As in ([1], proof of Theorem 1), on integration by parts, we get

$$\sigma_n(L^0(x)) = \Phi_\alpha(\pi)J(n, \pi) - \phi_\alpha(\pi)V(n, \pi) + \int_0^\pi V(n, u)d\phi_\alpha(u) .$$

Thus, by Lemma 3

$$\sigma_n(L^0(x)) = O(n^\alpha) - \phi_\alpha(\pi)V(n, \pi) + \int_0^\pi V(n, u)d\phi_\alpha(u) .$$

If in particular, we suppose  $\phi(t) = 1$  for all  $t$ , in which case  $\phi_\alpha(t) = 1$  for all  $t$  and  $\sigma_n(L^0(x)) = 0$  for every  $n$ , we obtain

$$0 = O(n^\alpha) - \phi_\alpha(\pi)V(n, \pi)$$

and therefore

$$\sigma_n(L^0(x)) = O(n^\alpha) + \int_0^\pi V(n, u)d\phi_\alpha(u) .$$

Thus,

$$\begin{aligned} & \sum_{n=1}^\infty \frac{1}{nP_n} |\sigma_n(L^0(x))| \\ & \leq K \sum_{n=1}^\infty \frac{1}{n^{1-\alpha}P_n} + K \int_0^\pi \sum_{n=1}^\infty \frac{1}{nP_n} |V(n, u)| |d\phi_\alpha(u)| \leq K , \end{aligned}$$

since by hypothesis  $\int_0^\pi |d\phi_\alpha(u)| \leq K$  and by Lemma 4,

$$\sum_{n=1}^\infty \frac{1}{nP_n} |V(n, u)| \leq Ku^\alpha \sum_{n \leq 1/u} n^{\alpha-1} + K\{1 + u^\alpha P_{[1/u]}\} \sum_{n > 1/u} \frac{1}{n^{1-\alpha}P_n} \leq K ,$$

by virtue of the hypothesis that  $\{p_n\} \in M_\alpha$ .

This completes the proof of  $|F_\alpha|$  - effective part of Theorem 1, when one appeals to Lemma 1.

(II)  $|\tilde{F}_\alpha|$  - effectiveness: We have

$$nB_n(x) = -\frac{2}{\pi} \int_0^\pi \psi(t) \frac{d}{dt} \cos ntdt$$

and therefore

$$\sigma_n(\tilde{L}^0(x)) = -\int_0^\pi \psi(t) \frac{d}{dt} \tilde{g}(n, t) dt .$$

As in ([2], proof of Theorem 2), we have

$$\begin{aligned} \sigma_n(\tilde{L}^0(x)) &= -\int_0^\pi \frac{d}{dt} \tilde{g}(n, t) \left\{ \frac{1}{\Gamma(1-\alpha)} \int_0^t (t-u)^{-\alpha} d\Psi_\alpha(u) \right\} dt \\ &= -\frac{1}{\Gamma(1-\alpha)} \int_0^\pi d\Psi_\alpha(u) \int_u^\pi (t-u)^{-\alpha} \frac{d}{dt} \tilde{g}(n, t) dt \\ &= -\int_0^\pi \tilde{J}(n, u) d\Psi_\alpha(u) . \end{aligned}$$



$|\tilde{F}^\alpha|$  – *effectiveness* of the  $(N, p)$  mean now follows from Lemma 1 and the hypothesis that  $\int_0^\pi u^{-\alpha} |d\mathcal{W}_\alpha(u)| \leq K$ , when we observe that uniformly in  $0 < u \leq \pi$

$$u^\alpha \sum_{n=1}^\infty \frac{1}{nP_n} |\tilde{J}(n, u)| \leq Ku^\alpha \sum_{n \leq 1/u} n^{\alpha-1} + Ku^\alpha P_{[1/u]} \sum_{n > 1/u} \frac{1}{n^{1-\alpha} P_n} \leq K,$$

by virtue of Lemma 3 and the hypothesis that  $\{p_n\} \in M_\alpha$ .

(III)  $|F^\alpha|$  – *effectiveness*: Integrating by parts, we have

$$n^{\alpha+1} A_n(x) = \frac{2}{\pi} \int_0^\pi \phi(t) n^{\alpha+1} \cos ntdt = -\frac{2}{\pi} \int_0^\pi n^\alpha \sin ntd\phi(t).$$

Thus

$$\sum_{n=1}^\infty \frac{1}{nP_n} |\sigma_n(L^\alpha(x))| \leq \int_0^\pi \left\{ \sum_{n=1}^\infty \frac{1}{nP_n} \left| \sum_{k=0}^n p_{n-k} k^\alpha \sin kt \right| \right\} |d\phi(t)|.$$

$|F^\alpha|$  – *effectiveness*, of the  $(N, p)$  mean now follows from Lemma 1 and the hypothesis that  $\int_0^\pi t^{-\alpha} |d\phi(t)| \leq K$ , when one observes that uniformly in  $0 < t \leq \pi$

$$\begin{aligned} & t^\alpha \sum_{n=1}^\infty \frac{1}{nP_n} \left| \sum_{k=0}^n p_{n-k} k^\alpha \sin kt \right| \\ & \leq Kt^{\alpha+1} \sum_{n \leq 1/t} n^\alpha + Kt^\alpha \sum_{n > 1/t} \frac{1}{n^{1-\alpha} P_n} \max_{0 \leq \nu \leq n} \left| \sum_{k=0}^\nu p_k \sin(n-k)t \right| \\ & \leq K + Kt^\alpha P_{[1/t]} \sum_{n > 1/t} \frac{1}{n^{1-\alpha} P_n} \leq K, \end{aligned}$$

by Abel’s lemma, Lemma 2 and the hypothesis that  $\{p_n\} \in M_\alpha$ .

(IV)  $|\tilde{F}^\alpha|$  – *effectiveness*: Integrating by parts and observing that  $\psi(+0) = 0$ , we have

$$\begin{aligned} n^{\alpha+1} B_n(x) &= \frac{2}{\pi} \int_0^\pi \psi(t) n^{\alpha+1} \sin ntdt \\ &= -\frac{2}{\pi} \psi(\pi) n^\alpha \cos n\pi + \frac{2}{\pi} \int_0^\pi n^\alpha \cos ntd\psi(t). \end{aligned}$$

Thus

$$\begin{aligned} \sum_{n=1}^\infty \frac{1}{nP_n} |\sigma_n(\tilde{L}^\alpha(x))| &\leq |\psi(\pi)| \sum_{n=1}^\infty \frac{1}{nP_n} \left| \sum_{k=0}^n p_{n-k} k^\alpha \cos k\pi \right| \\ &\quad + \int_0^\pi \left\{ \sum_{n=1}^\infty \frac{1}{nP_n} \left| \sum_{k=0}^n p_{n-k} k^\alpha \cos kt \right| \right\} |d\psi(t)|. \end{aligned}$$

$|\tilde{F}^\alpha|$  – *effectiveness* now follows from Lemma 1 and the hypothesis

that  $\int_0^\pi t^{-\alpha} |d\psi(t)| \leq K$ , when one observes that uniformly in  $0 < t \leq \pi$

$$\begin{aligned} & t^\alpha \sum_{n=1}^\infty \frac{1}{nP_n} \left| \sum_{k=0}^n p_{n-k} k^\alpha \cos kt \right| \\ & \leq Kt^\alpha \sum_{n \leq 1/t} n^{\alpha-1} + Kt^\alpha \sum_{n > 1/t} \frac{1}{n^{1-\alpha} P_n} \max_{0 \leq \nu \leq n} \left| \sum_{k=0}^\nu p_k \cos(n-k)t \right| \\ & \leq K + Kt^\alpha P_{[1/t]} \sum_{n > 1/t} \frac{1}{n^{1-\alpha} P_n} \leq K, \end{aligned}$$

by Lemma 2 and the hypothesis that  $\{p_n\} \in M_\alpha$ .

(V)  $|F^*|$  - effectiveness: This follows from the result of Theorem 3, when one observes that  $\{p_n\} \in M_\alpha, \alpha < 0$ , implies  $\{p_n\} \in M_0$ .

This completes the proof of Theorem 1.

**6. Proof of Theorem 2.** It may be observed that the proof of  $|F^\alpha|$  - effectiveness, given in the preceding section remains valid even for the case  $\alpha = 0$  and therefore the  $(N, p)$  method is  $|F^0|$  or equivalently  $|F_0|$  - effective.

In order to prove  $|\tilde{F}_0|$  - effectiveness, we observe that on integration by parts we get

$$\begin{aligned} B_n(x) &= \frac{2}{\pi} \int_0^\pi \psi(t) \sin ntdt = -\frac{2}{\pi} \int_0^\pi \frac{1 - \cos nt}{n} d\psi(t) \\ &= \frac{2}{\pi} \int_0^\pi \frac{\cos nt}{n} d\psi(t), \end{aligned}$$

since  $\psi(\pi) = \psi(0) = 0$ .

Thus, we have (cf. [13])

$$\frac{\pi}{2} \sigma_n(\tilde{L}^0(x)) = \begin{cases} -\int_0^\pi \left\{ \sum_{k=0}^n p_{n-k} (1 - \cos kt) \right\} d\psi(t), \\ \text{or} \\ \int_0^\pi \left\{ \sum_{k=0}^n p_{n-k} \cos kt \right\} d\psi(t), \end{cases}$$

and

$$\begin{aligned} & \sum_{n=1}^\infty \frac{1}{nP_n} |\sigma_n(\tilde{L}^0(x))| \\ & \leq \int_0^\pi |d\psi(t)| \left\{ \sum_{n \leq 1/t} \frac{1}{nP_n} \left| \sum_{k=0}^n p_{n-k} (1 - \cos kt) \right| + \sum_{n > 1/t} \frac{1}{nP_n} \left| \sum_{k=0}^n p_{n-k} \cos kt \right| \right\} \\ & = \int_0^\pi |d\psi(t)| \{\Sigma_1 + \Sigma_2\}, \end{aligned}$$

say.  $|\tilde{F}_0|$  - effectiveness of the  $(N, p)$  method now follows from Lemma 1 and the hypothesis that  $\int_0^\pi |d\psi(t)| \leq K$ , when one observes that uniformly in  $0 < t \leq \pi$

$$\Sigma_1 \leq Kt^2 \sum_{n \leq 1/t} n \leq K,$$

since  $|1 - \cos kt| \leq k^2 t^2$  and

$$\Sigma_2 \leq KP_{[1/t]} \sum_{n > 1/t} \frac{1}{nP_n} \leq K,$$

by virtue of Lemma 2 and the hypothesis that  $\{p_n\} \in M_0$ . This completes the proof of Theorem 2 (cf. [7]).

**7. Proof of Theorem 3.** We have ([14], § 13.2)

$$nA_n^*(x) = \frac{1}{\pi} \int_0^\pi \phi^*(t) \left\{ \sin \left( n + \frac{1}{2} \right) t / \sin \frac{1}{2} t \right\} dt.$$

Thus

$$\begin{aligned} & \sum_{n=1}^{\infty} \frac{1}{nP_n} |\sigma_n(L^*(x))| \\ & \leq \int_0^\pi \left\{ \sum_{n=1}^{\infty} \frac{1}{nP_n} \left| \sum_{k=0}^n p_{n-k} \sin \left( k + \frac{1}{2} \right) t \right| \right\} \frac{|\phi^*(t)|}{t} \frac{t}{\sin \frac{1}{2} t} dt. \end{aligned}$$

$|F^*|$  - effectiveness of the  $(N, p)$  method now follows from Lemma 1 and the hypothesis that  $\int_0^\pi t^{-1} |\phi^*(t)| dt \leq K$ , when we observe that uniformly in  $0 < t \leq \pi$

$$\begin{aligned} & \sum_{n=1}^{\infty} \frac{1}{nP_n} \left| \sum_{k=0}^n p_{n-k} \sin \left( k + \frac{1}{2} \right) t \right| \\ & \leq Kt \sum_{n \leq 1/t} 1 + KP_{[1/t]} \sum_{n > 1/t} \frac{1}{nP_n} \leq K, \end{aligned}$$

by virtue of Lemma 2 and the hypothesis that  $\{p_n\} \in M_0$ . This completes the proof of Theorem 3.

**8. Remarks.** Corresponding to our Theorem 2, we have an earlier result of Hille and Tamarkin ([8], Th. II) for ordinary  $(N, p)$  summability of  $L^0(x)$  which states that under certain condition on  $\phi(t)$  the hypothesis

$$(8.1) \quad \left\{ \frac{1}{P_n} \sum_{k=0}^n \frac{P_k}{k+1} \right\} \in B$$

is both necessary and sufficient for  $(N, p)$  summability of  $L^0(x)$ , if  $\{p_n\}$  is a positive monotonic nonincreasing sequence. The intrinsic character of the hypothesis  $\{p_n\} \in M_0$  of Theorem 2, emerges from

the above result when one observes that the condition (8.1) implies that

$$P_n \sum_{k=n}^{\infty} \frac{1}{kP_k} \leq K, \quad n = 1, 2, \dots$$

This follows from a recent paper of the author ([4], p. 168).

The claim that the corresponding (C) method results, reduce to special cases of our theorems, follows when we observe that

$$\left\{ \binom{n + \beta - 1}{\beta - 1} \right\} \in M_{\alpha}, \quad 1 \geq \beta > \alpha \geq 0,$$

and appeal to a well known inclusion relation for the absolute (C)-method.

Recently  $|F_1|$  - effectiveness of  $(N, p)(C, 1)$  and  $(C, 1)(N, p)$  methods have been proved by the present author.

#### REFERENCES

1. L. S. Bosanquet, *The absolute Cesàro summability of a Fourier series*, Proc. London Math. Soc. **41** (1936), 517-528.
2. L. S. Bosanquet and J. M. Hyslop, *On the absolute summability of the Allied series of a Fourier series*, Math. Z. **42** (1937), 489-512.
3. G. Das, *Tauberian theorems for absolute Nörlund summability*, Proc. London Math. Soc. (3) **19** (1969), 357-384.
4. H. P. Dikshit, *Absolute summability of a Fourier series by Nörlund means*, Math. Z. **102** (1967), 166-170.
5. ———, *Summability of a sequence of Fourier coefficients by a triangular matrix transformation*, Proc. Amer. Math. Soc. **21** (1969), 10-20.
6. ———, *Absolute summability of a series associated with a Fourier series*, Proc. Amer. Math. Soc. **22** (1969), 316-318.
7. ———, *Absolute summability of the conjugate series of a Fourier series by Nörlund means*, Math. Ann. **184** (1970), 106-112.
8. E. Hille and J. D. Tamarkin, *On the summability of Fourier series I*, Trans. Amer. Math. Soc. **34** (1932), 757-783.
9. N. Kishore and S. N. Bhatt, *Absolute Nörlund summability of a Fourier series*, Indian J. Math. **9** (1967), 259-267.
10. L. McFadden, *Absolute Nörlund summability*, Duke Math. J. **9** (1942), 168-207.
11. R. Mohanty, *The absolute Cesàro summability of some series associated with a Fourier series and its allied series*, J. London Math. Soc. **25** (1950), 63-67.
12. R. Mohanty and S. Mohapatra, *On the absolute convergence of a series associated with a Fourier series*, Proc. Amer. Math. Soc. **7** (1956), 1049-1053.
13. T. Pati, *On the absolute Nörlund summability of the conjugate series of a Fourier series*, J. London Math. Soc. **38** (1963), 204-213.
14. E. C. Titchmarsh, *The Theory of Functions*, Oxford University Press, New York, 1957.

Received October 22, 1969.

UNIVERSITY OF JABALPUR  
JABALPUR, INDIA



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON  
Stanford University  
Stanford, California 94305

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

RICHARD PIERCE  
University of Washington  
Seattle, Washington 98105

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLE

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL WEAPONS CENTER

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

# Pacific Journal of Mathematics

Vol. 33, No. 3

May, 1970

Charles A. Akemann, <i>Approximate units and maximal abelian <math>C^*</math>-subalgebras</i> .....	543
Gail Atneosen, <i>Wild points of cellular arcs in 2-complexes in <math>E^3</math> and cellular hulls</i> .....	551
John Logan Bryant and De Witt Summers, <i>On embeddings of 1-dimensional compacta in a hyperplane in <math>E^4</math></i> .....	555
H. P. Dikshit, <i>On a class of Nörlund means and Fourier series</i> .....	559
Nancy Dykes, <i>Generalizations of realcompact spaces</i> .....	571
Hector O. Fattorini, <i>Extension and behavior at infinity of solutions of certain linear operational differential equations</i> .....	583
Neal David Glassman, <i>Cohomology of nonassociative algebras</i> .....	617
Neal Hart, <i>Ulm's theorem for Abelian groups modulo bounded groups</i> .....	635
Don Barker Hinton, <i>Continuous spectra of second-order differential operators</i> .....	641
Donald Gordon James, <i>On Witt's theorem for unimodular quadratic forms. II</i> .....	645
Melvin F. Janowitz, <i>Principal multiplicative lattices</i> .....	653
James Edgar Keesling, <i>On the equivalence of normality and compactness in hyperspaces</i> .....	657
Adalbert Kerber, <i>Zu einer Arbeit von J. L. Berggren über ambivalente Gruppen</i> .....	669
Keizō Kikuchi, <i>Various <math>m</math>-representative domains in several complex variables</i> .....	677
Jack W. Macki and James Stephen Muldowney, <i>The asymptotic behaviour of solutions to linear systems of ordinary differential equations</i> .....	693
Andy R. Magid, <i>Locally Galois algebras</i> .....	707
T. S. Ravisankar, <i>On differentially simple algebras</i> .....	725
Joseph Gail Stampfli, <i>The norm of a derivation</i> .....	737
Francis C.Y. Tang, <i>On uniqueness of central decompositions of groups</i> .....	749
Robert Charles Thompson, <i>Some matrix factorization theorems. I</i> .....	763
Robert Charles Thompson, <i>Some matrix factorization theorems. II</i> .....	811