Pacific Journal of Mathematics

SETS WHICH CAN BE MISSED BY SIDE APPROXIMATIONS TO SPHERES

JAMES W. CANNON

Vol. 34, No. 2

June 1970

SETS WHICH CAN BE MISSED BY SIDE APPROXIMATIONS TO SPHERES

J. W. CANNON

In this paper it is proved that a 2-sphere S in E^3 can be side approximated from Int S so as to miss a closed subset Xof S if and only if $(S \cup \text{Int } S) - X$ is 1-ULC.

If $X = \{p\}$ is a singleton, then the 1-ULC property described in the first paragraph reduces to a property studied by McMillan [22] in connection with piercing points p of a crumpled cube (see (0.2) below). Our work was motivated by his paper although our methods bear little relationship to his.

Our result has as immediate corollaries a number of important, well-known theorems, four of which we now mention:

(0.1) ([2. Th. 2]). A 2-sphere S in E^3 is tame if $E^3 - S$ is 1-ULC.

(0.2) ([22, Th. 1]). If C is a crumpled cube in E^3 and $p \in Bd C$, then p is a piercing point of C if and only if C - p is 1-ULC.

(0.3) ([19, Th. 3]). If S and S' are 2-spheres in E^3 and $X \subset U \subset S \cap S'$, where X is compact and U is open in S and S'. Then (*, X, S') is satisfied if and only if (*, X, S) is satisfied.

(0.4) ([8] and [10]). If S is a 2-sphere in E^3 , X is a compact subset of S, and $X = \bigcup_{i=1}^{\infty} X_i$, where each X_i is compact and satisfies $(*, X_i, S)$, then (*, X, S) is satisfied.

Properties similar to the 1-ULC property described in the first paragraph have been studied previously (e.g., properties (A, F, S) and (B, F, S) in [19]). However, the properties studied have, in general, had only restricted application to the question of determining which subsets of a 2-sphere can be missed by side approximations to that sphere and thus have been inapplicable in results such as (0.2), (0.3), and (0.4).

Precise definitions appear in §1. Our main result, mentioned in the first paragraph, is proved in §2. We also present in §2 a slightly strengthened version of our main result for use in another paper (Lemma 2.5 and Theorem 2.6). The four corollaries mentioned above are proved in §3. In §3 we also remark on ways in which our main result can be used to sharpen other results which have appeared in the literature.

The part of this paper actually necessary for the proof of Theorem (0.1) appears in five short paragraphs. This proof is considerably shorter and conceptually easier to follow than the original proof. Our proof of Theorem (0.3) is likewise much shorter than the original. The

great simplifying factor is the use of Lister's form [18, Th. 3] of Bing's Side Approximation Theorem; Lister's theorem can be used to simplify the proof of many other results on the embeddings of surfaces in 3-manifolds (cf. [12]). Except for Lister's theorem, the tools we have chosen to use are essentially those which were used in the original proofs of Theorems (0.1)-(0.4). Lemma 2.5 and Theorem 2.6 are slightly more complex than the other results.

We had not originally planned to restrict ourselves in this paper to the applications of our main theorem which appear in §3 (cf. [9, Ths. 2 and 3]), but we have found that our other applications fit nicely into more general settings which we shall study elsewhere (cf. [10] and [11]).

We thank the referee for useful comments and C. E. Burgess for discussions.

1. Property (*, X, S), definitions, and notation. Suppose that X is a closed subset of a 2-sphere S in E^3 . We say that S can be side approximated so as to miss X if the following property is satisfied:

Property (*, X, S). ([15, p. 467]). If U is a component of $E^3 - S$ and ε is a positive number, then there are a polyhedral 2-sphere S', a finite collection D_1, D_2, \dots, D_k of mutually exclusive ε -disks lying in S', and a finite collection E_1, E_2, \dots, E_n of mutually exclusive ε -disks lying in S, such that

- (1) There is an ε -homeomorphism from S onto S',
- (2) $S' igcup_{i=1}^k D_i \subset U,$
- (3) $S \cap S' \subset igcup_{i=1}^n E_i$, and
- $(4) \quad X \cap (\bigcup_{i=1}^n E_i) = \varnothing.$

If (*, X, S) is satisfied, then X is tame (*i.e.*, X lies on a tame 2-sphere in E^3) [19, Th. 6]. If X has no degenerate components, then X is tame if and only if (*, X, S) is satisfied [8, Ths. 1.1 and 5.1].

If the property defined above holds for one component U of $E^3 - S$ though perhaps not for the other, we say that (*, X, U) is satisfied. If $(*, X, \operatorname{Int} S)$ is satisfied, then X lies on the surface of a 3-cell in E^3 [21, Th. 1]. Our goal is to characterize sets satisfying $(*, X, \operatorname{Int} S)$ as those compact sets X for which $(S \cup \operatorname{Int} S) - X$ is 1-ULC. A similar result will obviously hold for $U = \operatorname{Ext} S$.

REMARK. The definition that we have given of (*, X, S) is the original formulation and not always the easiest to apply. If we wish to prove that (*, X, Int S) is satisfied, we shall use the definition we have stated. If we already know that (*, X, Int S) is satisfied and

wish to apply that fact, we shall use the following formulation due to Lister. Lister proves [18, Th. 2] that the two definitions are equivalent.

Property (*, X, Int S) (alternative definition). If $\varepsilon > 0$, then there are disjoint ε -disks D_1, D_2, \dots, D_n in S - X and an ε -homeomorphism h from S onto a polyhedral sphere h(S) in E^3 such that

- (1) $h(S) = \bigcup_{i=1}^n \operatorname{Int} h(D_i) \subset \operatorname{Int} S$,
- (2) $S \bigcup_{i=1}^{n} \operatorname{Int} D_i \subset \operatorname{Ext} h(S)$, and
- (3) $S \cap h(S) = \bigcup_{i=1}^n [\operatorname{Int} D_i \cap \operatorname{Int} h(D_i)].$

REMARK. Property $(*, \emptyset, \text{Int } S)$ is always satisfied ([4, Th. 16'] (original formulation) and [18, Th. 2] (alternative definition)).

We shall use the symbol ρ for the Euclidean metric. If A and A' are subsets of E^3 and ε is a positive number, then we define

- (1) Diam $A = \sup \rho(x, y) \ (x, y \in A)$,
- (2) $\rho(A, A') = \inf \rho(a, a') \ (a \in A, a' \in A'), \text{ and}$
- (3) $B(A, A'; \varepsilon) = \{a' \in A' | \rho(a', A) < \varepsilon\}.$ (If $A' = E^3$, we write $B(A, A'; \varepsilon) = B(A; \varepsilon).$)

The set A is an ε -set if Diam $A < \varepsilon$. A map $f: A \to E^{\mathfrak{s}}$ is called an ε -map if f moves no point as far as ε . Let \varDelta and I denote a standard disk or 2-simplex and a standard interval or 1-simplex, respectively. We say that A is 1-LC (0-LC) in A' at $x \in E^3$ if for each $\varepsilon > 0$ there is a $\delta > 0$ such that any map $f: \operatorname{Bd} \Delta \to B(x, A; \delta)$ $(f: \operatorname{Bd} I \to B(x, A; \delta))$ has a continuous extension $f^*: \Delta \to B(x, A'; \varepsilon)$ $(f^*: I \to B(x, A'; \varepsilon))$. Speaking loosely, A is 1-LC in A' at $x \in E^3$ if loops near x in A can be shrunk to a point near x in A'. We say that A is 1-ULC (0-ULC) in A' if A is 1-LC (0-LC) in A' at each point x of ClA and if for each $\varepsilon > 0$ the corresponding δ may be chosen independently of $x \in$ Cl A. If A is 1-ULC in A, then we simply say that A is 1-ULC. We say that A is weakly 1-ULC in A' if (speaking loosely again) small unknotted simple closed curves in A bound small singular disks in A'. (A simple closed curve in E^3 is said to be unknotted if it bounds a disk in E^3 .) We leave it to the reader to formalize the definition of "weakly 1-ULC".

A crumpled cube C is the union of a 2-sphere in E^3 and its interior in E^3 . The 2-sphere is called the boundary of C, denoted Bd C; and C - Bd C is called the interior of C, denoted Int C. Any homeomorphic image of C is called a crumpled cube. A point $p \in Bd C$ is said to be a piercing point of C if there is an embedding $h: C \to E^3$ such that h(Bd C) can be pierced by a tame arc at h(p).

Most of our terminology not defined explicitly in this section is fairly standard and can be found in most of the references listed. See, for example, [2], [7], and [19].

2. A 1-ULC characterization of sets satisfying (*, X, Int S). Our characterization will appear in two parts which we prove separately (Ths. 2.1 and 2.4).

THEOREM 2.1. If S is a 2-sphere in E^3 , X is a closed subset of S, and (*, X, Int S) is satisfied, then $(S \cup \text{Int } S) - X$ is 1-ULC.

Proof. Let ε be a positive number. There is a δ in the range $0 < \delta < \varepsilon/6$ having the following property: if S' is a 2-sphere in E^3 such that there is a δ -homeomorphism from S onto S', then any δ -set on S' lies on an $\varepsilon/6$ -disk on S'. Let $f: \operatorname{Bd} A \to (S \cup \operatorname{Int} S) - X$ be a map into a δ -set in $(S \cup \operatorname{Int} S) - X$. In order to show that $(S \cup \operatorname{Int} S) - X$ is 1-ULC it suffices to show that f is nullhomotopic $(f \sim 0)$ in an ε -subset of $(S \cup \operatorname{Int} S) - X$.

Because Int S is 0-ULC [24, p. 66] and S is an absolute neighborhood retract, there is a map $F: \operatorname{Bd} \Delta \times I \to (S \cup \operatorname{Int} S) - X$ such that $F_0 = f, F_1: \operatorname{Bd} \Delta \to \operatorname{Int} S$, and Diam $F(\operatorname{Bd} \Delta \times I) < \delta$ (where $F_t: \operatorname{Bd} \Delta \to E^3$ is defined by $F_t(x) = F(x, t)$). The map F_1 can be extended to a map $G: \Delta \to E^3$ such that Diam $G(\Delta) < \delta$.

Since $(*, X, \operatorname{Int} S)$ is satisfied, there are disjoint δ -disks D_1, D_2, \dots, D_n in S - X and a δ -homeomorphism h from S onto a polyhedral 2-sphere h(S) in E^3 such that (1), (2), and (3) of the alternative definition for $(*, X, \operatorname{Int} S)$ are satisfied and such that h(S) separates $G(\operatorname{Bd} \Delta) = F_1(\operatorname{Bd} \Delta)$ from X.

The intersection of $G(\varDelta)$ with h(S) is a δ -set. Hence by our choice of δ , there is an $\varepsilon/6$ -disk E on h(S) which contains $G(\varDelta) \cap h(S)$. By the Tietze extension theorem (see [7, Lemma 1] for details), there is a map $G_1: \varDelta \to E^3$ which agrees with G on the component K of

$$arDelta - G^{-1}[G(arDelta) \cap h(S)]$$

which contains $\operatorname{Bd} \varDelta$ and takes $\varDelta - K$ into E. Then

$$G_{\scriptscriptstyle 1}(arDelta)\cap S\subset igcup_{i=1}^n D_i$$
 .

Again by the Tietze theorem there is a map $G_2: \Delta \to E^3$ which agrees with G_1 on the component K' of $\Delta - G_1^{-1}[G_1(\Delta) \cap \bigcup_{i=1}^n D_i]$ which contains $\operatorname{Bd} \Delta$ and takes $\Delta - K'$ into $\bigcup_{i=1}^n D_i$.

Clearly $f \sim 0$ in $F(\operatorname{Bd} \Delta \times I) \cup G_2(\Delta)$. Thus it suffices to show that $F(\operatorname{Bd} \Delta \times I) \cup G_2(\Delta) \subset (S \subset \operatorname{Int} S) - X$ and that

$$ext{Diam}\left[F(ext{Bd}\ arDelta imes I) \cup G_{ ext{2}}(arDelta)
ight] < arepsilon$$
 .

That these conditions are satisfied is easily checked. This completes the proof of Theorem 2.1.

The second half of our characterization requires two lemmas. The first, Lemma 2.2, is a special case of [22, Lemma 2], which in turn is a special case of [14, Th. 2]. We indicate a proof for Lemma 2.2 since it is a simple special case of the more difficult general results of [14]. Lemma 2.3, a consequence of Lemma 2.2, could be generalized considerably by appeal to these more general results. This is interesting when one notes that Theorem (0.4), a conjecture by Loveland, is simply the translation of Lemma 2.3, via Theorems 2.1 and 2.4, into a statement about side approximations of 2-spheres.

LEMMA 2.2. Let B denote a closed subset of a disk D, let S be a 2-sphere in E^3 , and let X be a subset of S such that $(S \cup \text{Int } S) - X$ is 1-ULC. Let $\varepsilon > 0$ and a map $f: D \to S \cup \text{Int } S$ be given. Then there is a map $f^*: D \to S \cup \text{Int } S$ such that

$$f^*(D-B) \subset (S \cup \operatorname{Int} S) - X,$$

 $f^*|B = f|B, and$
 $ho(f^*(x), f(x)) < \varepsilon \text{ for each } x \in D.$

Indication of proof. One triangulates D-B so that the mesh of the triangulation approaches zero very rapidly near B. Then using the facts that $\operatorname{Cl}((S \cup \operatorname{Int} S) - X) = (S \cup \operatorname{Int} S), (S \cup \operatorname{Int} S) - X$ is 0-ULC [24, p. 66], and $(S \cup \operatorname{Int} S) - X$ is 1-ULC, one moves the image of the 0-, 1-, and 2-skeletons of the triangulation away from X. Note that we do not require that X be closed.

LEMMA 2.3. Let S be a 2-sphere in E^3 , and let $\{X_i\}_{i=1}^{\infty}$ be a family of closed subsets of S such that, for each i, $(S \cup \text{Int } S) - X_i$ is 1– ULC. Then $(S \cup \text{Int } S) - \bigcup_{i=1}^{\infty} X_i$ is 1–ULC.

Proof. Suppose $\varepsilon > 0$ and $x \in S \cup \text{Int } S$ given, and let Δ denote a disk. Since $(S \cup \text{Int } S) - X_1$ is 1-ULC, there is a $\delta > 0$ such that each map $f: \text{Bd } \Delta \to B(x, (S \cup \text{Int } S) - X_1; \delta)$ extends to a map

$$f^*: \varDelta \to B(x, (S \cup \operatorname{Int} S) - X_1; \varepsilon)$$
.

Let $f: \operatorname{Bd} \Delta \to B(x, (S \cup \operatorname{Int} S) - \bigcup_{i=1}^{\infty} X_i; \delta)$ be given, and let

$$f_1: \varDelta \to B(x, (S \cup \operatorname{Int} S) - X_1; \varepsilon)$$

be an extension of f. We now define positive numbers $\varepsilon_1, \varepsilon_2, \cdots$ and maps f_2, f_3, \cdots inductively as follows:

Let

$$0$$

Let $f_2: \varDelta \to (S \cup \text{Int } S) - X_2$ be such that $f_2 | \text{Bd } \varDelta = f_1 | \text{Bd } \varDelta$ and

 $\rho(f_1(p),f_2(p))<arepsilon_1$

for each $p \in A$. Such a map exists by Lemma 2.2.

For i > 1, let ε_i be in the range $0 < \varepsilon_i < 1/2 \min \{\varepsilon_{i-1}, \rho[f_i(\varDelta), X_i]\}$. Let $f_{i+1}: \varDelta \to (S \cup \operatorname{Int} S) - X_{i+1}$ be such that $f_{i+1} | \operatorname{Bd} \varDelta = f_i | \operatorname{Bd} \varDelta$ and $\rho(f_i(p), f_{i+1}(p)) < \varepsilon_i$ for each $p \in \varDelta$. Again, at each stage such a map exists by Lemma 2.2.

Because $S \cup \text{Int } S$ is a complete space, it follows that $f^* = \lim f_i$ exists, is continuous, extends f, and takes \varDelta into

$$B\!\left(x,\,(S\cup\operatorname{Int}S)\,-\,igcup_{i=1}^\infty X_i;\,arepsilon
ight)$$
 .

Since x is arbitrary and δ may be chosen independently of x, $(S \cup \text{Int } S) - \bigcup_{i=1}^{\infty} X_i$ is 1-ULC.

THEOREM 2.4. If S is a 2-sphere in E^3 , X is a closed subset of S, and $(S \cup \text{Int } S) - X$ is 1-ULC, then (*, X, Int S) is satisfied.

Proof. We first establish the theorem for the special case X = S and then show how the proof can be altered to yield the general result. There are four steps in the special case.

1. There are disjoint disks D_1, D_2, \dots, D_n in S and an ε -homeomorphism h from S onto a polyhedral sphere h(S) in E^3 such that (1), (2), and (3) of the alternative definition of $(*, \emptyset, \text{Int } S)$ are satisfied and such that Diam $[D_i \cup h(D_i)] < \varepsilon$ for each i. This is because $(*, \emptyset, \text{Int } S)$ is always satisfied.

2. By the Tietze Extension Theorem, there are maps f_1, f_2, \dots, f_n such that for each $i, f: D_i \to h(D_i) \cup \text{Int } D_i, f_i$ agrees with h on the component K_i of $D_i - h^{-1}[h(D_i) \cap D_i]$ which contains $\text{Bd } D_i$, and f_i takes $D_i - K_i$ into $\text{Int } D_i$.

3. By Lemma 2.2, since Int S is 1-ULC, if $\delta > 0$, then there is a map $g_i: D_i \to f_i(K_i) \cup B(D_i, \text{Int } S; \delta)$ which agrees with f_i on

$$K_i - f_i^{\scriptscriptstyle -1}[B(D_i,f_i(D);\delta)]$$
 .

Thus we may assume that $g_1(D_i)$, $g_2(D_2)$, \cdots , $g_n(D_n)$ are disjoint singular disks in Int S such that for each i, (1) $g_i(D_i)$ has no singularities near $g_i(\operatorname{Bd} D_i)$, (2) Diam $[g_i(D_i) \cup D_i] < \varepsilon$, and (3) $g_i(D_i) \cap h[S - \bigcup_{i=1}^n \operatorname{Int} D_i] = g_i(\operatorname{Bd} D_i)$.

4. By Dehn's Lemma [23], each singular disk $g_i(D_i)$ may be replaced by a polyhedral disk E_i in Int S such that

326

$$h\!\left[S-igcup_{i=1}^n\operatorname{Int}D_i
ight]\cupigcup_{i=1}^nE_i$$

is a polyhedral 2-sphere S' in Int S and for each i, Diam $[E_i \cup D_i] < \varepsilon$. If h' is any homeomorphism from S onto S' which extends

$$h \left| S - igcup_{i=1}^n \operatorname{Int} D_i
ight|$$
 ,

then h' is an ε -homeomorphism from S into Int S. Since ε is arbitrary, this shows that (*, S, Int S) is satisfied.

In the general case, we make the following changes:

3'. Let $\{X_i\}_{i=1}^{\infty}$ be a family of tame finite graphs on S such that $S - \bigcup_{i=1}^{\infty} X_i$ is totally disconnected. That such a family exists follows from [3, Th. 1]. If A is an arc in any X_i , then $(*, A, \operatorname{Int} S)$ is satisfied by [15, Th. 2]. Thus by Theorem 2.1, $(S \cup \operatorname{Int} S) - A$ is 1-ULC; and by Lemma 2.3, $(S \cup \operatorname{Int} S) - (\bigcup_{i=1}^{\infty} X_i \cup X)$ is 1-ULC since $\bigcup_{i=1}^{\infty} X_i \cup X$ is a countable union of closed sets A for which $(S \cup \operatorname{Int} S) - A$ is 1-ULC. Hence by Lemma 2.2, if $\delta > 0$, there is a map $g_i: D_i \to f_i(K_i) \cup B(D_i, (S \cup \operatorname{Int} S) - (\bigcup_{i=1}^{\infty} X_i \cup X); \delta)$ which agrees with f_i on $K_i - f_i^{-1}[B(D_i, f_i(D_i); \delta)]$. Thus we may assume that

$$g_1(D_1), g_2(D_2), \cdots, g_n(D_n)$$

are disjoint singular disks in $(S \cup \text{Int } S) - X$ which satisfy (1), (2), and (3) of Step 3 above and such that for each i, $g_i(D_i) \cap S$ is a 0dimensional subset of $\text{Int } D_i$.

4'. The set $g_i(D_i) \cap S$, being a closed 0-dimensional subset of $\operatorname{Int} D_i - X$, can be covered by a finite collection $E_{i1}, E_{i2}, \dots, E_{in_i}$ of disjoint disks in $\operatorname{Int} D_i - X$. Again by Dehn's Lemma, each singular disk $g_i(D_i)$ may be replaced by a polyhedral disk E_i such that

$$h \Big[S - igcup_{i=1}^n \operatorname{Int} D_i \Big] \cup igcup_{i=1}^n E_i$$

is a polyhedral 2-sphere S' in E^3 and for each i, $E_i \cap S \subset \bigcup_{j=1}^{n_i} E_{ij}$ and Diam $[E_i \cup D_i] < \varepsilon$. But since ε is arbitrary, S' and the disks $\{E_i\}$ and $\{E_{ij}\}$ are precisely what one needs in order to show that (*, X, Int S) is satisfied, where in this case we apply the original definition.

We now show that the hypotheses of Theorem 2.4 can be weakened. Our goal is Theorem 2.6. In the proof of Theorem 2.6 we need to know that a polyhedral disk in E^3 whose boundary lies in the interior of a particular 2-sphere in E^3 (possibly a wild 2-sphere in E^3) can, in a particularly nice way, almost be cut off inside the 2-sphere. This fact, stated precisely in Lemma 2.5, will be apparent to those thoroughly familiar with the methods of [5]. However, following the suggestion of the referee, we give a proof which does not rely on the methods of [5] and depends only on Bing's Side Approximation Theorem [4]. The brief outline with which the proof of Lemma 2.5 begins is due to the referee and had also been suggested to the author by C. E. Burgess.

LEMMA 2.5. Suppose that S is a 2-sphere in E^s ; U is a component of $E^s - S$; D is a polyhedral disk in E^s with $\operatorname{Bd} D \subset U$; R is a disk in S with $D \cap S \subset R$; $0 < \varepsilon < \rho(\operatorname{Bd} D, S)$; and K is the the component of D - S which contains $\operatorname{Bd} D$. Then there is a map $h: D \to (D \cap U) \cup B(R, \operatorname{Cl} U; \delta)$ such that $h | K - B(R; \varepsilon) = identity$

 $h(D) \cap S$ is 0-dimensional; and $h|D - h^{-1}(S)$ is a homeomorphism.

Proof. The plan is the following: Let S_1 be a side approximation to S from U sufficiently close to S. Cut $D_0 = D$ off on S_1 to get a new polyhedral disk D_1 . Again let S_2 be a side approximation to Sfrom U sufficiently close to S (much closer than S_1). Cut D_1 off on S_2 to obtain a new polyhedral disk D_2 . Repeat this procedure. If proper care is taken in the process, then h(D) may be taken to be $\lim D_i$. In the next few paragraphs we explain in turn how each of the conclusions of the lemma can be obtained by this procedure.

Associating a continuous function h with $\lim D_i$. Cutting D_{i-1} off on S_i can be realized as a process of removing from D_{i-1} finitely many disjoint disks ${}^{i} \Delta_1, \dots, {}^{i} \Delta_r$ in $\operatorname{Int} D_{i-1}$ and replacing them by disjoint disks ${}^{i} \Delta'_1, \dots, {}^{i} \Delta'_r$ near S_i . Thus there is a natural, though nonunique, way of defining a homeomorphism $h_i: D_{i-1} \to D_i: h_i$ differs from the identity only in that it takes a disk ${}^{i} \Delta_j$ removed in the cutting process to its replacement ${}^{i} \Delta'_j$ near S_i . A function

$$h = \lim h_i \cdots h_2 h_1 \colon D \to \lim D_i$$

will therefore exist and be continuous provided the size of the disks removed and the size of their replacements converge to 0 rapidly enough with *i*. Suppose that to this end we want the disks $\{{}^{i+1}\Delta_j\}$ and $\{{}^{i+1}\Delta'_j\}$ to have diameter less than α . This can be accomplished by using caution in the adjustments which change D_{i-1} to D_i . The procedure is tedious but straightforward. The reader may wish to omit a reading of the details which follow:

Choose β , $0 < \beta < \alpha$, such that β -sets in a β -approximation S' to S lie in α -disks in S'.

Choose γ , $0 < \gamma < \beta/3$, such that γ -sets in a γ -approximation S'' to S lie in $\beta/3$ -disks in S''.

Choose δ , $0 < \delta < \gamma$, such that δ -sets in a δ -approximation D'_{i-1} to D_{i-1} lie in γ -disks in D'_{i-1} .

Choose a δ -side approximation S_i to S from U with associated δ -disks E_1, \dots, E_n on S_i and F_1, \dots, F_m on S containing the intersections of S_i with S.

Adjust D_{i-1} slightly near S_i (by a δ -adjustment) so that the adjusted D_{i-1} misses $\bigcup E_j$ and is in general position with respect to S_i . Call the adjusted D_{i-1} by the same name as the original.

Cut D_{i-1} off on S_i to form D_i . This cutting removes disjoint disks $\{{}^{i}\Delta_{j}\}$ from Int D_{i-1} and replaces them by disks $\{{}^{i}\Delta'_{j}\}$ near S_i .

We now show that $D_i \cap S$ can be covered by a finite collection of disjoint β -disks in $\operatorname{Int} D_i$. Since $[D_{i-1}(\operatorname{adjusted})] \cap (\cup E_j) = \emptyset$, the disk-with-holes $K_i = \operatorname{cl} (D_{i-1} - \cup^i \Delta_j)$ intersects S only in $\operatorname{Int} K_i$ and if care has been taken, also only in $\cup F_j$. From this latter fact, each component of $K_i \cap S$ has diameter less than δ . Hence by our choice of δ , $K_i \cap S$ can be covered by a finite collection of disjoint γ -disks in D_{i-1} , each of which has boundary in $\operatorname{Int} K_i$. The image of each such disk under $h_i: D_{i-1} \to D_i$ has diameter less than

$$\gamma+2eta/3 .$$

A disk ${}^{i}\varDelta'_{j}$ is either contained in one of these β -disks or intersects none of them. In the latter case, note that ${}^{i}\varDelta'_{j}$ is very close homeomorphically to some disk H in S_{i} such that Bd $H \cap (\bigcup E_{k}) = \emptyset$. If ${}^{i}\varDelta'_{j}$ is close enough to H, then ${}^{i}\varDelta'_{j} \cap S$ is covered in a natural way by β -disks in Int ${}^{i}\varDelta'_{j}$ associated with the δ -disks $\{E_{k} | E_{k} \subset \text{Int } H\}$. We thus obtain finally a finite collection of disjoint β -disks in Int D_{i} covering $D_{i} \cap S$.

If S_{i+1} is a β -approximation to S that intersects D_i only in the finite union of these disjoint β -disks, then the disks $\{{}^{i+1}\Delta_j\}$ will be of diameter less than $\beta(<\alpha)$ and the disks $\{{}^{i+1}\Delta'_j\}$ of diameter less than α by our choice of β .

These considerations show that the approximations S_i and disks D_i may be chosen so that $h = \lim h_i \cdots h_2 h_1$ automatically exists and is continuous.

Making $h|K - B(R; \varepsilon)$ the identity. One simply needs to keep all the spheres S_i away from some continuum in K which contains $K - B(R; \varepsilon)$.

Keeping h(D) in $(D \cap U) \cup B(R, \operatorname{Cl} U; \varepsilon)$. Note that

 $h(D) = \lim D_i \subset \lim (S_i \cup U(S_i)) = \operatorname{Cl} U$

(where $U(S_i)$ is the complementary domain of S_i which "almost coin-

cides" with U). Then if one keeps all of the disks $\{{}^{i}\varDelta_{j}^{\prime}\}$ in $B(R, \varepsilon/2)$, which one may clearly do, it follows immediately that

$$h(D) \subset (D \cap U) \cup B(R; \varepsilon)$$
.

Making $h(D) \cap S$ 0-dimensional. As we showed in the first part of this proof, by being careful at stage i we can require at stage i+1 that $D_i \cap S$ be covered by a collection of disjoint α -disks in S. One may then require that all of the disks $\{{}^{k} \mathcal{I}'_{j}\}$ $(k \ge i+1)$ lie in an arbitrary neighborhood of the union of these α -disks. It follows then that $h(D) \cap S$ also lies very close to the union of these α -disks, hence has no component of diameter much larger than α . If our α 's converge to 0, then $h(D) \cap S$ will be 0-dimensional.

Making $h|D - h^{-1}(S)$ a homeomorphism. This is automatic since each h_i is a homeomorphism and each point mapped into U has a neighborhood moved by only finitely many of the maps h_i .

THEOREM 2.6. If S is a 2-sphere in E^3 , X is a closed subset of S, and Int S is weakly 1-ULC in $(S \cup \text{Int } S) - X$, then $(S \cup \text{Int } S) - X$ is 1-ULC.

Proof. By Theorem 2.1, it suffices to show that (*, X, Int S) is satisfied. The proof is like that of the general case of Theorem 2.4 except that Lemma 2.5 allows us to alter the second step so that our weaker hypothesis is applicable in the third step. The first and fourth steps remain unchanged.

2". There are subdisks E_1, E_2, \dots, E_n of D_1, D_2, \dots, D_n respectively such that for each i, $h(D_i) \cap D_i \subset E_i \subset \text{Int } D_i$. Choose $\alpha > 0$ such that $\alpha < 1/3\rho[E_i, (S - D_i) \cup h(S - D_i)]$ and such that

Diam $[h(D_i) \cup B(D_i; \alpha)] < \varepsilon$

for each *i*. By Lemma 2.5, there are maps f_1, f_2, \dots, f_n such that for each $i, f_i: D_i \to h(D_i) \cup B(E_i, S \cup \text{Int } S; \alpha)$, f_i agrees with *h* on the component of $D_i - h^{-1}[B(D_i, S \cup \text{Int } S; \alpha)]$ which contains Bd $D_i, f_i(D_i) \cap S$ is a 0-dimensional subset of Int D_i , and $f_i | D_i - f_i^{-1}(S)$ is a homeomorphism. Define $f: S \to E^3$ piecewise by $f | S - \bigcup_{i=1}^n D_i = h | S - \bigcup_{i=1}^n D_i$ and $f | D_i = f_i$.

3". Choose β such that

 $0 < eta < 1/3
ho(f(D_i) \cap D_i, (S-D_i) \cup f(S-D_i))$

and such that $\text{Diam}[f(D_i) \cup B(D_i; \beta)] < \varepsilon$ for each *i*. Choose γ , using

the hypothesis that Int S is weakly 1-ULC in $(S \cup Int S) - X$, such that each unknotted γ -simple closed curve in Int S bounds a singular $\beta/2$ -disk in $(S \cup Int S) - X$. We require that $0 < \gamma < \beta/2$. Let K_i denote the component of $D_i - f^{-1}(S)$ which contains Bd D_i . Since $f(D_i) \cap S$ is 0-dimensional, there is a finite collection J_1, J_2, \dots, J_k of disjoint simple closed curves in K_i such that $\bigcup_{j=1}^k J_j$ separates Bd D_i from $f^{-1}(S)$ in D_i and such that each $f(J_i)$ is a γ -loop in a γ neighborhood of the set $f(D_i) \cap D_i$. We assume that no proper subcollection of J_1, J_2, \dots, J_k separates Bd D_i from $f^{-1}(S)$ in D_i . Because $f | D_i - f^{-1}(S)$ is a homeomorphism and $K_i \subset D_i - f^{-1}(S)$, it follows from Dehn's Lemma [23] that each $f(J_i)$ is an unknotted γ -simple closed curve in Int S. Hence by the way γ was chosen, there is a map $g_i: D_i \to (S \cup \text{Int } S) - X$ such that g_i agrees with f on that component of $K_i - \bigcup_{j=1}^k J_j$ which contains Bd D_i and such that g_i takes the disk in D_i bounded by J_j into a $\beta/2$ -subset of $(S \cup \text{Int } S) - X$. Define $g: S \to E^3$ piecewise by $g | S - \bigcup_{i=1}^n D_i = f | S - \bigcup_{i=1}^n D_i$ and $g \mid D_i = g_i$.

Now proceed as in Step 3' to choose $\{X_i\}_{i=1}^{\infty}$. Apply Lemma 2.3 only to the set $(S \cup \operatorname{Int} S) - \bigcup_{i=1}^{\infty} X_i$ to find this set to be 1-ULC. Then relative to some fixed $\delta > 0$, choose $h_i: D_i \to (S \cup \operatorname{Int} S) - \bigcup_{i=1}^{\infty} X_i$ as g_i was chosen in Step 3'. Require further that $\rho(g(x), h_i(x)) < \delta$ for each $x \in D_i$. Now for $0 < \delta < \rho(g(D_i), X)$ and for δ chosen otherwise sufficiently small, it follows as in Step 3' that

$$h_1(D_1), h_2(D_2), \dots, h_n(D_n)$$

are disjoint singular disks in $(S \cup \text{Int } S) - X$ which satisfy (1), (2) and (3) of Step 3 (special case) and such that for each i, $h_i(D_i) \cap S$ is a 0-dimensional subset of $\text{Int } D_i$.

The proof can now be completed as in Step 4'.

3. Proofs of Theorems (0.1)-(0.4).

Proof of Theorem (0.1). This is essentially the content of the special case of Theorem 2.4 which we proved first. Since $E^3 - S$ is 1-ULC, Theorem 2.4 and its counterpart for sets satisfying (*, X, Ext S) imply that (*, S, S) is satisfied. That is, S can be homeomorphically approximated from each of its complementary domains. Thus S is tame [1, Th. 2.2].

Proof of Theorem (0.2). Suppose the point p is a piercing point of C, and $h: C \to E^3$ is an embedding so that $\operatorname{Bd} h(C)$ can be pierced

by a tame arc at h(p). Then by (15, see Th. 11], $(*, h(p), \operatorname{Int} h(C))$ is satisfied. By [18, Th. 7] (or from Theorems 2.1 and 2.4 of this paper) $(*, p, \operatorname{Int} C)$ is satisfied. Thus by Theorem 2.1, C - p is 1–ULC.

Conversely, if C - p is 1-ULC, let $h: C \to E^3$ be an embedding such that Bd h(C) is tame from $E^3 - h(C)$ ([16], [17], or [12]). Then h(C) - h(p) is 1-ULC, and clearly $(*, h(p), E^3 - h(C))$ is satisfied. Hence by Theorem 2.4, (*, h(p), h(Bd C)) is satisfied. Therefore by [15, see Th. 11], p is a piercing point of C.

Proof of Theorem (0.3). Since the statement of the theorem is symmetric in S and S', we assume that (*, X, S) is satisfied and prove under this assumption that (*, X, S') is satisfied. By the compactness of S', the uniform nature of E^3 , and Theorems 2.1 and 2.4, it is sufficient to show that the sets $(S' \cup \operatorname{Int} S') - X$ and $(S' \cup \operatorname{Ext} S') - X$ are 1-LC at each point $x \in S'$. If $x \in S' - X$, both sets are clearly 1-LC at x. If $x \in X$, then since S and S' share an open set containing X, both spheres are locally the same at x. Thus clearly $(S' \cup \operatorname{Int} S') - X$ and $(S' \cup \operatorname{Ext} S') - X$ are 1-LC at x if and only if $(S \cup \operatorname{Int} S) - X$ and $(S \cup \operatorname{Ext} S) - X$ are 1-LC at x. This completes the proof.

Proof of Theorem (0.4). The set $(S \cup \text{Int } S) - X$ is 1-ULC by Theorem 2.1 and Lemma 2.3. Thus since X is closed (*, X, Int S) is satisfied by Theorem 2.4. Similarly (*, X, Ext S) is satisfied.

REMARK. [6, Th. 2.1] can be shown to be a special case of Theorem (0.4). Theorems 8, 9, and 10 of [20] are stated for closed subsets G of a 2-sphere S in E^3 such that the diameters of the components of G have a positive lower bound. It follows from Theorem 2.4 that the restriction that the diameters of G have a positive lower bound is unnecessary. That is, slight modifications of Loveland's proofs yield the following theorem:

THEOREM. If G is a compact subset of a 2-sphere S in E^3 and if any of the following conditions is satisfied, then G lies on a tame 2-sphere in E^3 .

(i) The set G can be locally spanned in each complementary domain of S on tame simple closed curves.

(ii) The set G can be locally spanned in each component of $E^3 - S$ missing Cantor sets.

(iii) The sphere S can be locally spanned at each point of G from each complementary domain of S.

332

Similarly, a version of Eaton's theorem on strongly locally spherical spheres [13, Th. 3] can be stated and proved using Theorem 2.4 and modifications of Eaton's proof.

THEOREM. If G is a compact subset of a 2-sphere S in E^3 such that S is strongly locally spherical at each point of G, then G lies on a tame 2-sphere.

References

1. R. H. Bing, Conditions under which a surface in E^3 is tame, Fund. Math. 47 (1959), 105-139.

2. _____, A surface is tame if its complement is 1-ULC, Trans. Amer. Math. Soc. 101 (1961), 294-305.

3. _____, Each disk in E³ contains a tame arc, Amer. J. Math. **84** (1962), 583-590.

4. _____, Approximating surfaces from the side, Ann. of Math. (2) 77 (1963), 145-192.

5. _____, Pushing a 2-sphere into its complement, Michigan Math. J. 11 (1964), 33-45.

6. ____, Improving the side approximation theorem, Trans. Amer. Math. Soc. 116 (1965), 511-525.

7. C. E. Burgess, Characterizations of tame surfaces in E^3 , Trans. Amer. Math. Soc. **114** (1965), 80-97.

8. J. W. Cannon, Characterization of taming sets on 2-spheres, Trans. Amer. Math. Soc. 147 (1970), 289-299.

9. _____, Characterizations of sets which can be missed by side approximations of spheres, Noticer Amer. Math. Soc. 16 (1969), 174.

10. _____, Singular side approximations for 2-spheres in E^3 (in preparation)

11. _____. *-taming sets for crumpled cubes (in preparation)

12. R. J. Daverman, A new proof for the Hosay-Lininger Theorem about crumpled cubes, Proc. Amer. Math. Soc. 23 (1969), 52-54.

13. W. T. Eaton, A note about locally spherical spheres, Canad. J. Math. 21 (1969), 1001-1003.

14. S. Eilenberg and R. L. Wilder, Uniform local connectedness and contractibility, Amer. J. Math. 64 (1942), 613-622.

15. David S. Gillman, Side approximation, missing an arc, Amer. J. Math. 85 (1963), 459-476.

16. Norman Hosay, The sum of a real cube and a crumpled cube is S^3 , Notices Amer. Math. Soc. **10** (1963) 666.

17. L. L. Lininger, Some results on crumpled cubes, Trans. Amer. Math. Soc. 118 (1965), 534-549.

18. F. M. Lister, Simplifying intersections of disks in Bing's side approximation theorem, Pacific J. Math. 22 (1967), 281-295.

19. L. D. Loveland, Tame subsets of spheres in E^3 , Pacific J. Math. 19 (1966), 489-517.

20. _____, Tame surfaces and tame subsets of spheres in E^3 , Trans. Amer. Math. Soc. 123 (1966), 355-368.

21. _____, Sufficient conditions for a closed set to lie on the boundary of a 3-cell, Proc. Amer. Math. Soc. 19 (1968), 649-652.

22. D. R. McMillan, Jr., Some topological properties of piercing points, Pacific J. Math. 22 (1967), 313-322.

23. C. D. Papakyriakopoulos, On Dehn's lemma and the asphericity of knots, Ann. of

Math. (2) 66 (1957), 1-26.

24. R. L. Wilder, Topology of manifolds, Amer. Math. Soc. Colloquium Publications 32 (1949).

Received September 30, 1969, and in revised form December 5, 1969. These results were obtained while the author was a National Science Foundation Graduate Fellow at the University of Utah.

UNIVERSITY OF UTAH UTAH STATE UNIVERSITY UNIVERSITY OF WISCONSIN

334

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON Stanford University Stanford, California 94305

J. DUGUNDJI Department of Mathematics University of Southern California Los Angeles, California 90007

RICHARD ARENS

University of California Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH	B. H. NEUMANN	F. WOLE	K. YOSHIDA
	SUPPORTING	INSTITUTIONS	5
UNIVERSITY OF BRI	TISH COLUMBIA	STANFORD UN	IVERSITY
CALIFORNIA INSTIT	UTE OF TECHNOLOGY	UNIVERSITY C	OF TOKYO
UNIVERSITY OF CALIFORNIA		UNIVERSITY O)F UTAH
MONTANA STATE UNIVERSITY		WASHINGTON	STATE UNIVERSITY
UNIVERSITY OF NEW	ADA	UNIVERSITY C	F WASHINGTON
NEW MEXICO STATE	UNIVERSITY	* *	*
OREGON STATE UNI	VERSITY	AMERICAN MA	ATHEMATICAL SOCIETY
UNIVERSITY OF ORE	GON	CHEVRON RES	EARCH CORPORATION
OSAKA UNIVERSITY		TRW SYSTEMS	
UNIVERSITY OF SOU	THERN CALIFORNIA	NAVAL WEAP	ONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal. but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

RICHARD PIERCE University of Washington Seattle, Washington 98105

Pacific Journal of Mathematics Vol. 34, No. 2 June, 1970

Shair Ahmad, On the oscillation of solutions of a class of linear fourth order differential equations	289
Leonard Asimow and Alan John Ellis, <i>Facial decomposition of linearly</i>	
compact simplexes and separation of functions on cones	301
Kirby Alan Baker and Albert Robert Stralka, <i>Compact, distributive lattices of</i>	211
	311
James W. Cannon, Sets which can be missed by side approximations to	221
spheres	321
Prem Chandra, Absolute summability by Riesz means	335
Francis T. Christoph, Free topological semigroups and embedding topological	2.42
semigroups in topological groups	343
Henry Bruce Cohen and Francis E. Sullivan, <i>Projecting onto cycles in smooth</i> ,	255
reflexive Banach spaces	300
John Dauns, Power series semigroup rings	365
Robert E. Dressler, A density which counts multiplicity	371
Kent Ralph Fuller, <i>Primary rings and double centralizers</i>	379
Gary Allen Gislason, On the existence question for a family of products	385
Alan Stuart Gleit, On the structure topology of simplex spaces	389
William R. Gordon and Marvin David Marcus, An analysis of equality in	
certain matrix inequalities. I	407
Gerald William Johnson and David Lee Skoug, Operator-valued Feynman	
integrals of finite-dimensional functionals	415
(Harold) David Kahn, <i>Covering semigroups</i>	427
Keith Milo Kendig, Fibrations of analytic varieties	441
Norman Yeomans Luther, Weak denseness of nonatomic measures on perfect, locally compact spaces	453
Guillermo Owen. The four-person constant-sum games: Discriminatory	155
solutions on the main diagonal	461
Stephen Parrott Unitary dilations for commuting contractions	481
Roy Martin Pakestraw Extranal alements of the convex cone A of	401
functions $A_n = A_n = A_n = A_n$	491
Peter Lewis Repz Intersection representations of graphs by arcs	501
William Henry Puckle, Representation and series summability of complete	501
biorthogonal sequences	511
E Dennis Sentilles The strict topology on hounded sets	520
Scheron Sheleh A note on Hanf numbers	541
Sanaron Shehan, A note on Hunj numbers	541
rings	547
Imgs	547
Kenneur S. williams, Finite transformation formulae involving the Legendre	550
<i>symuot</i>	559