

Pacific Journal of Mathematics

ON PARTIAL HOMOMORPHISMS OF SEMIGROUPS

B. D. ARENDT AND C. J. STUTH

ON PARTIAL HOMOMORPHISMS OF SEMIGROUPS

B. D. ARENDT AND C. J. STUTH

Let S be a semigroup and T be a semigroup with zero ($T=T^0$). An ideal extension of S by T is a semigroup V containing S as an ideal and such that the Rees quotient V/S is isomorphic to T . A mapping α from $T^* = T - \{0\}$ into S is said to be a partial homomorphism, if $t_1, t_2 \in T^*$, $t_1 t_2 \neq 0$ implies $(t_1 t_2)\alpha = (t_1\alpha)(t_2\alpha)$. Every partial homomorphism from T^* into S gives rise to an ideal extension of S by T . Further, in certain cases every ideal extension of S by T is obtained in this way. In this paper a characterization is given for all partial homomorphisms from T^* into S .

It is not known in general when all extensions of S by T are determined by the partial homomorphisms of T^* into S . Clifford has shown this to be the case when S has an identity (see [2, §4.4]). Further results in this direction have been obtained by Warne [4] and Petrich [3]. The partial homomorphisms of a completely 0-simple semigroup into an arbitrary semigroup have been determined by Clifford [1].

An element x of a semigroup S is said to be *prime* if x does not belong to S^2 . S is said to have *unique factorization* if every nonzero element of S can be written uniquely as a product of powers of primes. Of course, if S is not commutative, we must take the order of the factors into account. We define the *kernel* of a homomorphism into a semigroup with zero to be the complete inverse image of zero.

THEOREM 1. *A [commutative] semigroup S has unique factorization if and only if S is free [commutative] or the Rees quotient of a free [commutative] semigroup.*

Proof. Suppose S has unique factorization, and let X be the set of primes of S . If $0 \notin S$, then clearly S is free [commutative] on X . So assume $0 \in S$, and let F_X be the free [commutative] semigroup on X with homomorphism ϕ from F_X onto S such that $x\phi = x$ for all $x \in X$ [2, p. 41]. Let K be the kernel of ϕ . Since S has unique factorization, ϕ must be one-to-one on $F_X - K$, so S is isomorphic to the Rees quotient F_X/K . The converse is obvious.

COROLLARY 2. *If $S = S^0$, then there exists a semigroup U with unique factorization and a homomorphism from U onto S with trivial kernel.*

Proof. There exists a free semigroup F which is homomorphic onto S with kernel K . Set $U = F/K$ and use Theorem 1.6 of [2].

THEOREM 3. *Let U be a semigroup with unique factorization and let X be the set of primes of U . Let S be any semigroup. Then any mapping α from X into S can be extended to a partial homomorphism of U^* into S .*

Proof. Omitted.

We denote by π_α the equivalence relation induced by a mapping α on its domain and use \leq for the usual partial ordering of relations on a set.

THEOREM 4. *Let $T = T^0$ and S be semigroups. By Corollary 2 there exists a semigroup U with unique factorization and a homomorphism ϕ from U onto T with trivial kernel. Let α be any partial homomorphism from U^* into S such that $\pi_\phi \leq \pi_\alpha$ on U^* and define $\alpha': T^* \rightarrow S$ as follows. If $y \in T^*$ then there exists an $x \in U^*$ such that $y = x\phi$ and we define $y\alpha' = x\alpha$. Then α' is a partial homomorphism from T^* into S . Conversely every partial homomorphism of T^* into S is determined in this manner. Finally, the mapping $\alpha \rightarrow \alpha'$ is one-to-one.*

Proof. α' is well defined since $\pi_\phi \leq \pi_\alpha$ on U^* , and it is a partial homomorphism since α is. Conversely, if α' is a partial homomorphism from T^* into S , then define $x\alpha = x\phi\alpha'$ for $x \in U^*$. If $x_1, x_2 \in U^*$ with $x_1x_2 \neq 0$, then $(x_1x_2)\phi\alpha' = ((x_1\phi)(x_2\phi))\alpha'$ and this in turn is equal to $(x_1\phi\alpha')(x_2\phi\alpha')$ since ϕ has trivial kernel. Thus α is a partial homomorphism from U^* into S such that $\pi_\phi \leq \pi_\alpha$ on U^* .

Now let α, β be partial homomorphisms from U^* into S such that $\pi_\phi \leq \pi_\alpha, \pi_\phi \leq \pi_\beta$ on U^* and $\alpha' = \beta'$. Thus for all $x \in U^*$, $x\phi\alpha' = x\phi\beta' \Rightarrow x\alpha = x\beta$ so $\alpha = \beta$ and the mapping is one-to-one.

BIBLIOGRAPHY

1. A. H. Clifford, *Partial homomorphic images of Brandt groupoids*, Proc. Amer. Math. Soc. **16** (1965), 538-544.
2. A. H. Clifford and G. B. Preston, *Algebraic theory of semigroups*, vol. 1, Math. Surveys No. 7, Amer. Math. Soc., Providence, R.I., 1961.
3. Mario Petrich, *On extensions of semigroups determined by partial homomorphisms*, Nederl. Akad. Wetensch. Proc. (A) **69**; Indag. Math. **28** (1966), 49-51.
4. R. J. Warne, *Extensions of completely 0-simple semigroups by completely 0-simple semigroups*, Proc. Amer. Math. Soc. **17** (1966), 524-526.

Received October 7, 1969. The first author was partially supported by a University of Missouri Research Grant. The second author was partially supported by NSF Grant No. 9618.

THE UNIVERSITY OF MISSOURI
COLUMBIA, MISSOURI

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

B. D. Arendt and C. J. Stuth, <i>On the structure of commutative periodic semigroups</i>	1
B. D. Arendt and C. J. Stuth, <i>On partial homomorphisms of semigroups</i>	7
Leonard Asimov, <i>Extensions of continuous affine functions</i>	11
Claude Elias Billigheimer, <i>Regular boundary problems for a five-term recurrence relation</i>	23
Edwin Ogilvie Buchman and F. A. Valentine, <i>A characterization of the parallelepiped in E^n</i>	53
Victor P. Camillo, <i>A note on commutative injective rings</i>	59
Larry Jean Cummings, <i>Decomposable symmetric tensors</i>	65
J. E. H. Elliott, <i>On matrices with a restricted number of diagonal values</i> ...	79
Garth Ian Gaudry, <i>Bad behavior and inclusion results for multipliers of type (p, q)</i>	83
Frances F. Gulick, <i>Derivations and actions</i>	95
Langdon Frank Harris, <i>On subgroups of prime power index</i>	117
Jutta Hausen, <i>The hypo residuum of the automorphism group of an abelian p-group</i>	127
R. Hrycay, <i>Noncontinuous multifunctions</i>	141
A. Jeanne LaDuke, <i>On a certain generalization of p spaces</i>	155
Marion-Josephine Lim, <i>Rank preservers of skew-symmetric matrices</i>	169
John Hathway Lindsey, II, <i>On a six dimensional projective representation of the Hall-Janko group</i>	175
Roger McCann, <i>Transversally perturbed planar dynamical systems</i>	187
Theodore Windle Palmer, <i>Real C^*-algebras</i>	195
Don David Porter, <i>Symplectic bordism, Stiefel-Whitney numbers, and a Novikov resolution</i>	205
Tilak Raj Prabhakar, <i>On a set of polynomials suggested by Laguerre polynomials</i>	213
B. L. S. Prakasa Rao, <i>Infinitely divisible characteristic functionals on locally convex topological vector spaces</i>	221
John Robert Reay, <i>Caratheodory theorems in convex product structures</i>	227
Allan M. Sinclair, <i>Eigenvalues in the boundary of the numerical range</i>	231
David R. Stone, <i>Torsion-free and divisible modules over matrix rings</i>	235
William Jennings Wickless, <i>A characterization of the nil radical of a ring</i>	255