

Pacific Journal of Mathematics

**ALGEBRAS WITH MINIMAL LEFT IDEALS WHICH ARE
HILBERT SPACES**

BRUCE ALAN BARNES

ALGEBRAS WITH MINIMAL LEFT IDEALS WHICH ARE HILBERT SPACES

BRUCE A. BARNES

This paper gives a necessary and sufficient condition that certain topological algebras A (normed algebras and algebras which are inner product spaces) be left (right) annihilator algebras. It is assumed that the socle of A is dense in A and that a proper involution $*$ is defined on the socle. Then the necessary and sufficient condition is essentially that the minimal left (right) ideals of A be complete in the norm on A and be a Hilbert space in an equivalent norm.

We prove a useful preliminary result in § 2. In § 3 we deal with the question of when a normed algebra A is a left or right annihilator algebra. In § 4 we consider this same question when A is a topological algebra in a topology defined by an inner product. This section is motivated by the work of P. Saworotnow and B. Yood on such algebras (see, for example, [5] and [9]). In the final section we generalize the well known result of Bonsall and Goldie that B^* annihilator algebras are dual.

Notation and terminology. A is always a complex algebra. S_A denotes the socle of A , when this exists. If E is a subset of A , let $L(E)$ and $R(E)$ denote the left and right annihilator of E respectively ($L(E) = \{u \in A \mid uv = 0 \text{ for all } v \in E\}$). A is a left (right) annihilator algebra if for every proper closed right (left) ideal M of A $L(M) \neq 0$ ($R(M) \neq 0$) and $L(A) = 0$ ($R(A) = 0$). A left (right) ideal M of A is a left (right) annihilator ideal if $M = L(E)$ ($M = R(E)$) for some subset E of A . If A is semi-simple, A is a modular annihilator algebra if A/S_A is a radical algebra; see [8]. Annihilator and dual algebras are defined and discussed in [4, pp.96-107].

An involution $*$ defined on A (or S_A) is proper if $uu^* = 0$ implies $u = 0$. u is a self-adjoint if $u = u^*$. We denote the set of all self-adjoint minimal idempotents of A by H . If $*$ is proper on S_A , then minimal left (right) ideals of A will have the form Ah (hA), $h \in H$, by [4, Lemma 4.10.1, p. 261].

Let \mathcal{H} be a Hilbert space. $\mathcal{B}(\mathcal{H})$ is the algebra of all bounded operators on \mathcal{H} , $\mathcal{F}(\mathcal{H})$ is the subalgebra of $\mathcal{B}(\mathcal{H})$ consisting of all operators which have finite dimensional range, and $\mathcal{C}(\mathcal{H})$ is the algebra of compact operators on \mathcal{H} . If $T \in \mathcal{B}(\mathcal{H})$, we denote the operator bound of T as $|T|$. Given $u, v \in \mathcal{H}$, we define an operator $(u|v)$ on \mathcal{H} by $(u|v)(w) = (w, u)v$ for all $w \in \mathcal{H}$. More generally

if X is a normed linear space and X' is the normed dual of X , given $x \in X$ and $f \in X'$ we define an operator $(f|x)$ on X by $(f|x)(y) = f(y)x$ for all $y \in X$.

2. Preliminary results. Let \mathcal{H} be a Hilbert space. Assume that B is a subalgebra of $\mathcal{B}(\mathcal{H})$ with $\mathcal{F}(\mathcal{H}) \subset B$. Furthermore, assume that B is a topological linear space with a topology \mathcal{T} such that

(i) The maps $x \rightarrow xy$ and $x \rightarrow yx$ are continuous on B for all $y \in B$;

(ii) $\mathcal{F}(\mathcal{H})$ is dense in B in the topology \mathcal{T} ;

(iii) If $\{u_n\} \subset \mathcal{H}$ and $u_n \rightarrow 0$ in \mathcal{H} , then $(w|u_n) \rightarrow 0$ in \mathcal{T} for any $w \in \mathcal{H}$.

For \mathcal{K} a closed subspace of \mathcal{H} , define

$$\mathcal{R}(\mathcal{K}) = \{T \in B \mid T(\mathcal{H}) \subset \mathcal{K}\}.$$

THEOREM 2.1. *Assume B is as given above. Then B is a left annihilator algebra. Also every right annihilator ideal of B is of the form $\mathcal{R}(\mathcal{K})$ for some closed subspace \mathcal{K} of \mathcal{H} . If $T \in \overline{TB}$ for all $T \in B$, then every closed right ideal of B is a right annihilator ideal.*

Proof. Assume that N is a closed right ideal of B . Let

$$\mathcal{J} = \{Tu \mid T \in N, u \in \mathcal{H}\}.$$

Assume that $w = T(u) + S(v)$ where $u, v \in \mathcal{H}$ and $T, S \in N$. Assume that $u \neq 0$, and let $\lambda = \|u\|_2^{-2}$ ($\|\cdot\|_2$ the norm on \mathcal{H}). Then

$$(1/\lambda)S(u|v) \in N$$

and $(T + (1/\lambda)S(u|v))(u) = w$. This proves that \mathcal{J} is a subspace of \mathcal{H} . Let $\mathcal{K} = \overline{\mathcal{J}}$. The proof of [4, Lemma 2.8.24, p. 104] implies that $R(L(N)) = \mathcal{R}(\mathcal{K})$. If $v \in \mathcal{K}$, then there exists $\{u_n\} \subset \mathcal{H}$ and $\{T_n\} \subset N$ such that $T_n(u_n) = v_n \rightarrow v$ in \mathcal{H} . Then given any $w \in \mathcal{H}$, $(w|v_n) = T_n(w|u_n) \in N$ for all n . By (iii) $(w|v_n) \rightarrow (w|v)$ in the topology \mathcal{T} . Thus whenever $v \in \mathcal{K}$ and $w \in \mathcal{H}$, $(w|v) \in N$. Using this result, the proof of [4, Lemma 2.8.26, p. 105] implies that

$$R(L(N)) \cdot B \subset N.$$

Therefore if $L(N) = 0$, $B^2 \subset N$, and it follows that $\mathcal{F}(\mathcal{H}) \subset N$. Then by (ii) $N = B$. This proves that B is a left annihilator algebra.

If $T \in \overline{TB}$ for all $T \in B$, then whenever $T \in R(L(N))$, $T \in \overline{TB} \subset \overline{R(L(N)) \cdot B} \subset N$. Therefore $N = R(L(N))$, so that N is a right annihilator ideal.

A theorem similar to Theorem 2.1 can be proved concerning the left ideals of B . Assume that $\mathcal{F}(\mathcal{H}) \subset B \subset \mathcal{B}(\mathcal{H})$ and that B satisfies (i), (ii), and

(iv) If $\{u_n\} \subset \mathcal{H}$ and $u_n \rightarrow 0$ in \mathcal{H} , then $(u_n|v) \rightarrow 0$ in the topology \mathcal{F} on B for all $v \in \mathcal{H}$

Define $B^* = \{T^* | T \in B\}$. Topologize B^* with the topology

$$\mathcal{F}^* = \{U^* | U \in \mathcal{F}\}.$$

Then $\mathcal{F}(\mathcal{H}) \subset B^* \subset \mathcal{B}(\mathcal{H})$ and B^* satisfies (i) and (ii). But also by (iv) and the fact that $(v|w)^* = (w|v)$ for all $v, w \in \mathcal{H}$, B^* satisfies (iii). Then the conclusions of Theorem 2.1 hold for B^* . Therefore B^* is a left annihilator algebra and every right annihilator ideal is of the form $\{T \in B^* | T(\mathcal{H}) \subset \mathcal{N}\}$ for some closed subspace \mathcal{N} of \mathcal{H} . Let N be a proper closed left ideal of B . Then N^* is a proper closed right ideal of B^* . Therefore there exists $T \in B, T \neq 0$, such that $T^*N^* = 0$. Then $R(N) \neq 0$. Now assume that N is a left annihilator ideal of B . Then N^* is a right annihilator ideal of B^* which implies that $N^* = \{T \in B^* | T(\mathcal{H}) \subset \mathcal{N}^\perp\}$ for \mathcal{N} some closed subspace of \mathcal{H} . Then it is not difficult to verify that

$$N = \{T \in B | T(\mathcal{N}) = 0\}.$$

Finally if $T \in \overline{BT}$ for all $T \in B$, then $T \in \overline{TB}^*$ for all $T \in B^*$. This implies that when $T \in \overline{BT}$ for all $T \in B$, then every closed left ideal of B is a left annihilator ideal (by Theorem 2.1 again).

Combining these remarks and Theorem 2.1 we have the following result.

THEOREM 2.2. *Assume that $\mathcal{F}(\mathcal{H}) \subset B \subset \mathcal{B}(\mathcal{H})$ and that B satisfies (i)–(iv). Then B is an annihilator algebra. If in addition $T \in \overline{TB}$ and $T \in \overline{BT}$ for all $T \in B$, then B is dual.*

3. Normed algebras. We assume throughout this section that A is a semi-simple modular annihilator algebra, that there is a proper involution $*$ defined on S_A , and that A is a normed algebra with norm $\|\cdot\|$. Recall that H denotes the set of self-adjoint minimal idempotents of A . When $h \in H$, we define a functional f_h on S_A by the rule $f_h(u)h = huh$. By the proof of [7, Th. 5.2, p. 358] we have that f_h is a positive hermitian functional on S_A . We introduce an inner product on the minimal left ideal Ah by the usual definition, $(uh, vh) = f_h((vh)^*uh)$, $u, v \in A$. We call this inner product the canonical inner product on Ah and denote the corresponding norm by

$|\cdot|_2$. We define a *-representation of S_A on the inner product space Ah by $u \rightarrow T_u^h$, $u \in S_A$, where $T_u^h(vh) = uvh$ for all $v \in A$. As shown in the proof of [7, Th. 5.2, p. 358], the operators T_u^h are bounded on Ah . Also by [7, Lemma 7.1, p. 358] T_u^h has finite dimensional range on Ah for all $u \in S_A$. In a similar fashion a canonical inner product can be introduced on the minimal right ideal hA , and a *-representation of S_A can be constructed into $\mathcal{B}(hA)$.

Since S_A is a modular annihilator algebra with proper involution $*$, then by [1, (1.3), p. 6] there is a unique norm $|\cdot|$ on S_A with the property that $|uu^*| = |u|^2$ for all $u \in S_A$. We call $|\cdot|$ the operator norm on S_A .

THEOREM 3.1. *Assume that A is a left (right) annihilator algebra in the norm $\|\cdot\|$. Also assume that there exists $K > 0$ such that $K\|u\| \geq |u|$ for all $u \in S_A$. Then for any $h \in H$, Ah (hA) is a Hilbert space in the canonical norm $|\cdot|_2$, and $\|\cdot\|$ and $|\cdot|_2$ are equivalent on Ah (hA).*

Proof. We consider only the case where A is a left annihilator algebra. Also it is sufficient to prove the theorem when A is primitive. For in the general case given $h \in H$, Ah is a minimal left ideal of some minimal closed two sided ideal M of A . Then M is primitive and by the proof of [4, Th. 2.8.12, p. 99] M is a left annihilator algebra. Therefore assume that A is primitive. We shall show that S_A is a left annihilator algebra. If N is a proper closed right ideal of S_A , then \bar{N} , the closure of N in A , is a proper closed right ideal of A . Then $L(\bar{N}) \neq 0$, and therefore there exists a minimal idempotent $e \in L(\bar{N})$. Then $e \in S_A$ and $eN = 0$. Thus S_A is a left annihilator algebra.

Assume $h \in H$. Note that $|uh|^2 = |(uh)^*uh| = |uh \frac{1}{2} h| = |uh \frac{1}{2}|$ so that $|\cdot|$ and $|\cdot|_2$ coincide on Ah . By hypothesis $K\|u\| \geq |u|$ for all $u \in S_A$, and therefore $K\|uh\| \geq |uh|_2$ for all $u \in A$. We prove that $\|\cdot\|$ and $|\cdot|_2$ are equivalent on Ah . Since A is primitive, the representation $u \rightarrow T_u^h$ of S_A on Ah is faithful. Let $\mathcal{F} = \{T_u^h | u \in S_A\}$. By the proof of [4, Lemma 2.8.20, p. 101] $(f|x) \in \mathcal{F}$ whenever f is a continuous linear functional on Ah with respect to $\|\cdot\|$ and $x \in Ah$. It follows that any such functional f must be continuous on Ah with respect to $|\cdot|_2$. Let V be the normed dual of Ah with respect to $\|\cdot\|$, and let B be the unit ball in Ah with respect to $|\cdot|_2$. For any $f \in V$, $\sup_{x \in B} |f(x)| < +\infty$. Then by the Uniform Boundedness Theorem applied to the set B ; $\sup_{x \in B} \sup_{\|f\| \leq 1} |f(x)| \leq J$ for some finite number J . It follows that $\|x\| \leq J|x|_2$ for all $x \in Ah$. Therefore $\|\cdot\|$ and $|\cdot|_2$ are equivalent on Ah .

It remains to be shown that Ah is a Hilbert space in the norm

$|\cdot|_2$. Since $K\|u\| \geq |u|$ for all $u \in S_A$, S_A is a left annihilator algebra with respect to $|\cdot|$. Let \mathcal{H} be the Hilbert space completion on Ah . Given $w \in \mathcal{H}$, we define $f(x) = (x, w)$ for $x \in \mathcal{H}$. Choose $uh \in Ah$ such that $|uh|_2 = 1$. $(f|uh) \in \mathcal{F}$ by the proof of [4, Lemma 2.8.20, p. 101]. Therefore $(f|uh)^* \in \mathcal{F}$. For any $x \in Ah$,

$$(x, w) = ((f|uh)x, uh) = (x, (f|uh)^*uh). \text{ Therefore}$$

$$w = (f|uh)^*(uh) \in Ah.$$

Thus $\mathcal{H} = Ah$.

Using the previous result, we give an example of a norm on $\mathcal{F}(\mathcal{H})$ in which $\mathcal{F}(\mathcal{H})$ is not a left annihilator algebra. Let \mathcal{H} be an infinite dimensional Hilbert space and denote the norm on \mathcal{H} by $|\cdot|_2$. Let $\|\cdot\|$ be any norm on \mathcal{H} such that $|x|_2 \leq \|x\|$ for all $x \in \mathcal{H}$, and $|\cdot|_2$ and $\|\cdot\|$ are inequivalent on \mathcal{H} . If f is any discontinuous linear functional on \mathcal{H} , then $\|x\| = |x|_2 + |f(x)|$ is an example of such a norm. Every functional on \mathcal{H} continuous with respect to $|\cdot|_2$ is continuous with respect to $\|\cdot\|$. It follows that every operator $T \in \mathcal{F}(\mathcal{H})$ is bounded in the norm

$$\|T\| = \sup_{\|x\| \leq 1} \|Tx\|.$$

We note that there exists $K > 0$ such that $K\|T\| \geq |T|$ ($|\cdot|$ the operator norm on $\mathcal{F}(\mathcal{H})$) by [4, Th. 2.4.17, p. 69]. Now fix $u \in \mathcal{H}$ such that $|u|_2 = 1$. Let N be the minimal left ideal of $\mathcal{F}(\mathcal{H})$ defined by $N = \{(u|v) | v \in \mathcal{H}\}$. $v \rightarrow (u|v)$ is an isometry of \mathcal{H} in the norm $|\cdot|_2$ onto N in the operator norm since $|(u|v)| = |u|_2|v|_2 = |v|_2$. To verify that $\mathcal{F}(\mathcal{H})$ is not a left annihilator algebra in the norm $\|\cdot\|$, it is sufficient to prove that the map $v \rightarrow (u|v)$ is a bicontinuous map from \mathcal{H} in the norm $\|\cdot\|$ onto N in the norm $\|\cdot\|$. For then $\|\cdot\|$ and $|\cdot|$ are inequivalent on N , and therefore Theorem 3.1 gives the result.

$$\|(u|v)\| = \sup_{\|x\| \leq 1} \|(u|v)(x)\| = \sup_{\|x\| \leq 1} |(x, u)| \|v\| \leq \|v\|,$$

and

$$\|(u|v)\| \geq \|(u|v)(u/\|u\|)\| = (1/\|u\|)\|v\|.$$

This completes the example.

Now we prove a converse of Theorem 3.1.

THEOREM 3.2. *Assume that S_A is dense in A . Assume that for every $h \in H$ Ah (hA) is a Hilbert space in the norm $|\cdot|_2$, and that $|\cdot|_2$ and $\|\cdot\|$ are equivalent on Ah (hA). Then A is a left (right)*

annihilator algebra. If in addition $u \in \overline{uA}$ ($u \in \overline{Au}$) for all $u \in A$, then every closed right (left) ideal of A is a right (left) annihilator ideal.

Proof. We assume that for every $h \in H$ Ah is a Hilbert space in the norm $|\cdot|_2$, and that $|\cdot|_2$ and $\|\cdot\|$ are equivalent on Ah . First suppose that A is primitive. Given $h \in H$, then $u \rightarrow T_u^h$ is a faithful $*$ -representation of S_A on the Hilbert space Ah . Given any $u, v, w \in A$, $T_{(u_h)(v_h)^*}^h(wh) = (wh, v_h)(uh) = (v_h|uh)(wh)$. Therefore all the operators of the form $(v_h|uh)$ are in the image of the representation $w \rightarrow T_w^h$. It follows that $\mathcal{F}(Ah)$ is in the image of this representation. By [4, Th. 2.4.17, p. 69] there exists $K > 0$ such that $K\|u\| \geq |T_u^h|$ for all $u \in S_A$. Then since S_A is dense in A , there is a unique extension of the representation $u \rightarrow T_u^h$ of S_A to a representation $u \rightarrow T_u$ of A onto a subalgebra B of $\mathcal{B}(Ah)$. Therefore

$$\mathcal{F}(Ah) \subset B \subset \mathcal{B}(Ah).$$

We consider B normed by $\|\cdot\|$ in the natural way, $\|T_u\| = \|u\|$ for $u \in A$. B clearly has properties (i) and (ii) listed previous to Theorem 2.1. If $|u_n h|_2 \rightarrow 0$, then by hypothesis $\|u_n h\| \rightarrow 0$, and therefore

$$\|(wh|u_n h)\| = \|T_{(u_n h)(wh)^*}\| \rightarrow 0$$

for any $w \in A$. This proves that B also satisfies (iii). By Theorem 2.1, B , and hence A , is a left annihilator algebra. If in addition $u \in \overline{uA}$ for all $u \in A$, then again by Theorem 2.1, every closed right ideal of A is a right annihilator ideal. This proves the theorem when A is primitive. In the general case let $\{M_\alpha | \alpha \in I\}$ be the set of all minimal closed two sided ideals of A . M_α is primitive for each $\alpha \in I$, and therefore the theorem holds for each M_α . Since A has dense socle, A is the topological sum of the $M_\alpha, \alpha \in I$. Then by the proof of [4, Th. 2.8.29, p. 106], the theorem holds for A .

4. Algebras which are inner product spaces. Throughout this section we assume that A is a semi-simple modular annihilator algebra which is an inner product space with inner product (\cdot, \cdot) . Also we assume that the maps $x \rightarrow xy$ and $x \rightarrow yx$ are continuous on A for all $y \in A$. An element x has a left (right) adjoint if there exists $w \in A$ such that $(xy, z) = (y, wz)$ ($(yx, z) = (x, zw)$) for all $y, z \in A$. If $x \in A$ has a left (right) adjoint, then it is unique. Assume that every element $u \in S_A$ has a left adjoint which we denote by u^* . Suppose that $u^*u = 0$. By [1, (2.2), p. 6] there exists an idempotent $e \in A$ such that $u = ue$. Then $0 = (u^*u, e) = (u, u)$ so that $u = 0$. This verifies that $*$ must be proper on S_A . Similarly if every element in

S_A has a right adjoint, then this adjoint must be proper on S_A . We denote the norm determined on A by the inner product by $|\cdot|_2$.

THEOREM 4.1. *Assume that every element $u \in S_A$ has a left (right) adjoint u^* and that A is a left (right) annihilator algebra. Then for every $h \in H$, Ah (hA) is a Hilbert space in the norm $|\cdot|_2$, and $|\cdot|_2$ and $|\cdot|_2$ are equivalent on Ah (hA).*

Proof. We prove the “left” part of the theorem only. As in the proof of Theorem 3.1, it is sufficient to prove the theorem when A is primitive. Therefore assume A is primitive. Given $h \in H$,

$$(uh, vh) = ((vh)^*uh, h) = (uh, vh)|h|_2^2$$

for all $u, v \in A$. Therefore $|\cdot|_2$ and $|\cdot|_2$ are equivalent on Ah . $u \rightarrow T_u^h$ is a faithful representation of S_A on Ah . Let $\mathcal{F} = \{T_u^h | u \in S_A\}$. By the same argument as in the proof of Theorem 3.1, S_A is a left annihilator algebra with respect to $|\cdot|_2$. Then by the proof of [4, Lemma 2.8.20, p. 101] $(f|uh) \in \mathcal{F}$ for all $u \in A$ and all functionals f continuous on Ah with respect to $|\cdot|_2$. Then the argument in the last paragraph of the proof of Theorem 3.1 implies that Ah is a Hilbert space in the norm $|\cdot|_2$.

Now we prove a result in the other direction.

THEOREM 4.2. *Assume that every element $u \in S_A$ has a left (right) adjoint u^* . Assume that A has dense socle in the norm $|\cdot|_2$ and that for every $h \in H$, Ah (hA) is a Hilbert space in the norm $|\cdot|_2$. Then A is a left (right) annihilator algebra. If in addition $u \in \overline{uA}$ ($u \in \overline{Au}$) for all $u \in A$, then every closed right (left) ideal of A is a right (left) annihilator ideal.*

Proof. We prove the “left” part of the theorem only. It is sufficient to prove that the theorem holds for each minimal closed two sided ideal M of A . For then by the proof of [4, Th. 2.8.29, p. 106] the result follows for A . Therefore assume that M is a minimal closed two sided ideal of A . Choose $h \in H \cap M$. Then $u \rightarrow T_u^h$ is a faithful representation of M on the Hilbert space Ah . T_u^h is a bounded operator on Ah since $u \rightarrow ux$ is a continuous map on A . Let $B = \{T_u^h | u \in M\}$. We norm B by $|T_u^h|_2 = |u|_2$ for $u \in M$, Given wh and vh , then $T_{(uh)(vh)^*}^h \in B$, and

$$T_{(uh)(vh)^*}^h(wh) = (wh, vh)wh = (vh|uh)(wh)$$

for all $wh \in Ah$. Therefore $\mathcal{F}(Ah) \subset B$. B satisfies properties (i)

and (ii) given previous to Theorem 2.1 by hypothesis. Also as noted in the proof of Theorem 4.1, $\|uh\|_2^2 = \|uh\|_2^2 \|h\|_2^2$ for all $u \in A$. Therefore if $\|u_n h\|_2 \rightarrow 0$, then $\|u_n h\|_2 \rightarrow 0$, so that for any $v \in A$, $\|u_n h(vh)^*\|_2 \rightarrow 0$. It follows that $\|(vh|u_n h)\|_2 = \|T_{(u_n h)(vh)^*}^h\|_2 \rightarrow 0$. Therefore B satisfies (iii). Then Theorem 2.1 applies and this completes the proof.

We apply the previous theorems to right-modular complemented algebras as defined by B. Yood [9, p. 261]. Let A be an algebra with an inner product (\cdot, \cdot) . A is a right-modular complemented algebra if

- (a) the maps $x \rightarrow xy$ and $x \rightarrow yx$ are continuous for all $y \in A$,
- (b) any right or left ideal I for which $I^\perp = \{0\}$ is dense in A (where $I^\perp = \{x \in A \mid (x, y) = 0 \text{ for all } y \in I\}$),
- (c) the intersection of the closed modular maximal right ideals of A is $\{0\}$, and M^\perp is a right ideal for each closed modular maximal right ideal M .

We prove the following theorem.

THEOREM 4.3. *Assume that A is a modular annihilator algebra and a right-modular complemented algebra. Then A is an annihilator algebra if and only if every minimal left or right ideal of A is a Hilbert space in the norm determined by the inner product.*

Proof. First note that A is semi-simple by property (c). Since A is a modular annihilator algebra, then by [8, Lemma 3.3, p. 38] every modular maximal right ideal M of A is of the form $(1 - e)A$ where e is a minimal idempotent of A . Then by (a) M is closed. Similarly every modular maximal left ideal of A is closed. Also by [9, Th. 2.1, p. 262] K^\perp is a right (left) ideal for all right (left) ideals K of A .

Assume that every minimal left or right ideal of A is a Hilbert space in the norm determined by the inner product. Given K a minimal right ideal of A , then $N = K^\perp$ is a right ideal. Also $N + K$ is dense by (b). Since K is complete, it follows that $N + K = A$. Therefore N is a modular maximal right ideal of A . By the proof of [8, Th. 4.5, p. 44] every element of $N^\perp = K$ has a left adjoint. Since K was an arbitrary minimal right ideal, then every element in S_A has a left adjoint. A similar proof shows that every element of S_A has a right adjoint. A has dense socle by (b). Therefore by Theorem 4.2, A is an annihilator algebra.

Now assume that A is an annihilator algebra. Take K minimal right ideal of A . Then $N = K^\perp$ is a proper closed right ideal of A . Since A is an annihilator algebra, there exists a modular maximal right ideal M such that $N \subset M$. $K + N$ is a dense right ideal of A

by (b). Assume that $x \in K^{\perp\perp}$. Then there exists $\{x_n\} \subset N$ and $\{y_n\} \subset K$ such that $x_n + y_n \rightarrow x$. Then

$$|x_n|_2 = |(x_n + y_n - x, x_n / |x_n|_2)| \leq |x_n + y_n - x|_2 \rightarrow 0.$$

Therefore $y_n \rightarrow x$ and since K is closed, $x \in K$. It follows that $K = K^{\perp\perp}$. Now $K^\perp \subset M$, and therefore $M^\perp \subset K$. Since M^\perp is a nonzero right ideal of A , $M^\perp = K$. Then every element in K has a left adjoint by the proof of [8, Th. 4.5, p. 44]. It follows that every element in S_A has a left adjoint, and by a similar proof every element in S_A has a right adjoint. Then Theorem 4.1 implies that every minimal left or right ideal of A is a Hilbert space in the norm determined by the inner product.

5. Algebras dual in the operator norm. A well known theorem of Bonsall and Goldie states that an annihilator B^* -algebra is dual. This was generalized by B. Yood who proved that any modular annihilator B^* -algebra is dual; see [8, Th. 4.1, p. 42]. In this section we generalize this result still further. We assume throughout that A is a modular annihilator algebra with an involution $*$ and a norm $|\cdot|$ with the property that $|u^*u| = |u|^2$ for all $u \in A$ (such a norm always exists on A when A is a normed algebra and $*$ is proper by [7, Th. 5.2, p. 358]). We call $|\cdot|$ the operator norm on A .

THEOREM 5.1. *Assume that A has the properties given above. Then if every minimal left ideal of A is complete in the operator norm, A is dual.*

We prove three lemmas.

LEMMA 5.2. *If every minimal left ideal of A is complete in the operator norm, then there is an isometric $*$ -representation $u \rightarrow T_u$ of A onto a subalgebra B of the compact operators on a Hilbert space \mathcal{H} with the following properties:*

(1) \mathcal{H} is the Hilbert space direct sum of a set of closed subspaces $\mathcal{H}_\alpha, \alpha \in I$ where I is some index set.

(2) If $T \in B$, then T is reduced by each $\mathcal{H}_\alpha, \alpha \in I$ (i.e.,

$$T(\mathcal{H}_\alpha) \subset \mathcal{H}_\alpha \text{ and } T(\mathcal{H}_\alpha^\perp) \subset \mathcal{H}_\alpha^\perp$$

all $\alpha \in I$).

(3) If $T \in \mathcal{F}(\mathcal{H})$ and T is reduced by \mathcal{H}_α for all $\alpha \in I$, then $T \in B$.

(4) $B \cap \mathcal{F}(\mathcal{H})$ is dense in B .

Proof. Let $\{M_\alpha | \alpha \in I\}$ be the set of minimal two sided ideals of

A, I some index set. For each $\alpha \in I$, choose an element $h_\alpha \in H \cap M_\alpha$. Let $\mathcal{H}_\alpha = Ah_\alpha$. Ah_α is an inner product space in the canonical inner product. Also $|uh_\alpha|^2 = |(uh_\alpha)^*(uh_\alpha)| = |uh_\alpha|_2^2$. Therefore $|\cdot|_2$ coincides with $|\cdot|$ on Ah_α . Therefore \mathcal{H}_α is a Hilbert space. Let \mathcal{H} be the Hilbert space direct sum of the $\mathcal{H}_\alpha, \alpha \in I$. For each α we have a $*$ -representation $u \rightarrow T_u^{h_\alpha}$ of A on $Ah_\alpha = \mathcal{H}_\alpha$. $|T_u^{h_\alpha}| \leq |u|$ for all $u \in A, \alpha \in I$. Then we define $u \rightarrow T_u$ a representation of A on \mathcal{H} in the usual fashion, $T_u(\sum_{\alpha \in I} v_\alpha h_\alpha) = \sum_{\alpha \in I} T_u^{h_\alpha}(v_\alpha h_\alpha)$. $u \rightarrow T_u$ is a faithful $*$ -representation of A onto a subalgebra B of $\mathcal{B}(\mathcal{H})$. By [1, (1.3), p. 6] $|u| = |T_u|$ for all $u \in A$. T_u has finite dimensional range for all $u \in S_A$ by [7, Lemma 5.1, p. 358]. Also the socle of A is dense in A by the proof of [2, Lemma 2.6, p. 287]. It follows that $\mathcal{F}(\mathcal{H}) \cap B$ must be dense in B and that $B \subset \mathcal{C}(\mathcal{H})$. It remains to prove (3). By Theorem 3.2 A is a left annihilator algebra, and by the proof of that theorem $\mathcal{F}(\mathcal{H}_\alpha) \subset \{T_u^{h_\alpha} | u \in M_\alpha\}$. Assume that $T \in \mathcal{F}(\mathcal{H}), T(\mathcal{H}_\alpha) \subset \mathcal{H}_\alpha$, and $T(\mathcal{H}_\alpha^\perp) \subset \mathcal{H}_\alpha^\perp$ for all $\alpha \in I$. Then $T(\mathcal{H}_\alpha) = 0$ for all but a finite number of $\alpha \in I, \alpha_1, \alpha_2, \dots, \alpha_n$. Then there exists $u_k \in M_{\alpha_k}, 1 \leq k \leq n$, such that $T_{u_k}^{h_{\alpha_k}}(x) = T(x)$ for all $x \in \mathcal{H}_\alpha$. Let $u = u_1 + \dots + u_n$. Then $T_u(x) = T(x)$ for all $x \in \mathcal{H}$. This proves (3).

LEMMA 5.3. *Let B be as in Lemma 5.2. Then $T \in \overline{TB}$ and $T \in \overline{BT}$ for all $T \in B$.*

Proof. Assume that $T \in B$. Then T^*T is a compact operator on the Hilbert space \mathcal{H} . Let $\{\lambda_k\}$ be the sequence of distinct nonzero eigenvalues of T^*T . Let $\{E_k\}$ be the sequence of projections onto the corresponding eigenspaces. For all $\alpha \in I$ denote by F_α the projection onto the subspace \mathcal{H}_α . By hypothesis $F_\alpha T^*T = T^*T F_\alpha$ for all $\alpha \in I$. It follows that $F_\alpha E_k = E_k F_\alpha$ for all $\alpha \in I$ and all k . By (3) of Lemma 5.2 $E_k \in B$ for all k . Then $|T - \sum_{k=1}^N T E_k|^2 = |(T - \sum_{k=1}^N T E_k)^*(T - \sum_{k=1}^N T E_k)| = |T^*T - \sum_{k=1}^N \lambda_k E_k|$. Since $T^*T = \sum_{k=1}^{+\infty} \lambda_k E_k$ by the Spectral Theorem for compact operators, then $T(\sum_{k=1}^N E_k) \rightarrow T$ as $N \rightarrow +\infty$. This proves $T \in \overline{TB}$. A similar argument using TT^* in place of T^*T shows that $T \in \overline{BT}$.

LEMMA 5.4. *Assume that \mathcal{H} is a Hilbert space. Then $\mathcal{F}(\mathcal{H})$ is dual in the operator norm.*

Proof. Assume that M is a closed right ideal of $\mathcal{F}(\mathcal{H})$, and let $N = M + L(M)^*$. N is a right ideal of $\mathcal{F}(\mathcal{H})$. Let

$$\mathcal{I} = \{Tu | T \in N, u \in \mathcal{H}\}.$$

As in the proof of Theorem 2.1, \mathcal{I} is a subspace of \mathcal{H} . If $w \perp \mathcal{I}$,

then for every $u \in \mathcal{H}$ and $T \in N$, $(w|w)T(u) = (Tu, w)w = 0$. Therefore $(w|w)N = 0$. But then $(w|w)M = 0$ and $L(M)(w|w) = 0$. Therefore $|w|_2^2(w|w) = (w|w)^2 = 0$ so that $w = 0$. This proves that \mathcal{L} is dense in \mathcal{H} . Assume $v, w \in \mathcal{H}$. Choose $\{u_n\} \subset \mathcal{H}$ and $\{T_n\} \subset N$ such that $T_n(u_n) = v_n \rightarrow v$. Then $T_n(w|u_n) = (w|v_n) \rightarrow (w|v)$ so that $(w|v) \in N$. Therefore $\mathcal{F}(\mathcal{H}) = N$. Take $T \in R(L(M))$. $T = T_1 + T_2$ where $T_1 \in M$ and $T_2 \in L(M)^*$. Then $T_2^*T = 0$ and $T_2^*T_1 = 0$. Thus $T_2^*T_2 = 0$ which implies $T_2 = 0$. It follows that $R(L(M)) = M$. If M is a closed left ideal of $\mathcal{F}(\mathcal{H})$, then $L(R(M)) = M$ by taking involutions. Therefore $\mathcal{F}(\mathcal{H})$ is dual.

Now we complete the proof of Theorem 5.1. By Lemma 5.2. it is enough to prove that an algebra B with the properties listed in that lemma is dual. Let F_α be the projection of \mathcal{H} onto \mathcal{H}_α for all $\alpha \in I$. Set $S_\alpha = \{T \in \mathcal{F}(\mathcal{H}) \mid TF_\alpha = F_\alpha T = T\}$. By Lemma 5.2 $S_\alpha \subset B$. Furthermore $\mathcal{F}(\mathcal{H}_\alpha)$ is isometrically isomorphic to S_α . Therefore S_α is dual by Lemma 5.4. Also S_α is a two sided ideal of B for each $\alpha \in I$, and B is the topological sum of the S_α , $\alpha \in I$. By Lemma 5.3 $T \in \overline{TB}$ and $T \in \overline{BT}$ for all $T \in B$. Then it follows from the proof of [4, Th. 2.8.29, p. 106] that B is dual.

REFERENCES

1. B. Barnes, *Subalgebras of modular annihilator algebras*, Proc. Camb. Phil. Soc. **66** (1969), 5-12.
2. ———, *Algebras with the spectral expansion property*, Illinois J. Math. **11** (1967), 284-290.
3. I. Kaplansky, *Dual rings*, Ann. of Math. **49** (1948), 689-701.
4. C. Rickart, *General Theory of Banach Algebras*, Princeton, 1960.
5. P. P. Saworotnow, *A generalization of the notion of H^* -algebra*, Proc. Amer. Math. Soc. **8** (1957), 49-55.
6. B. Yood, *Homomorphisms on normed algebras*, Pacific J. Math. **8** (1958), 373-381.
7. ———, *Faithful $*$ -representations of normed algebras*, Pacific J. Math. **10** (1960), 345-363.
8. ———, *Ideals in topological rings*, Canad. J. Math. **16** (1964), 28-45.
9. ———, *On algebras which are pre-Hilbert spaces*, Duke Math. J. **36** (1969), 261-272.

Received January 29, 1970. This research was partially supported by a grant from the Graduate School of the University of Oregon.

THE UNIVERSITY OF OREGON
EUGENE, OREGON

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLE

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics

Vol. 35, No. 3

November, 1970

John D. Arrison and Michael Rich, <i>On nearly commutative degree one algebras</i>	533
Bruce Alan Barnes, <i>Algebras with minimal left ideals which are Hilbert spaces</i>	537
Robert F. Brown, <i>An elementary proof of the uniqueness of the fixed point index</i>	549
Ronn L. Carpenter, <i>Principal ideals in F-algebras</i>	559
Chen Chung Chang and Yiannis (John) Nicolas Moschovakis, <i>The Suslin-Kleene theorem for V_κ with cofinality $(\kappa) = \omega$</i>	565
Theodore Seio Chihara, <i>The derived set of the spectrum of a distribution function</i>	571
Tae Geun Cho, <i>On the Choquet boundary for a nonclosed subspace of $C(S)$</i>	575
Richard Brian Darst, <i>The Lebesgue decomposition, Radon-Nikodym derivative, conditional expectation, and martingale convergence for lattices of sets</i>	581
David E. Fields, <i>Dimension theory in power series rings</i>	601
Michael Lawrence Fredman, <i>Congruence formulas obtained by counting irreducibles</i>	613
John Eric Gilbert, <i>On the ideal structure of some algebras of analytic functions</i>	625
G. Goss and Giovanni Viglino, <i>Some topological properties weaker than compactness</i>	635
George Grätzer and J. Sichler, <i>On the endomorphism semigroup (and category) of bounded lattices</i>	639
R. C. Lacher, <i>Cell-like mappings. II</i>	649
Shiva Narain Lal, <i>On a theorem of M. Izumi and S. Izumi</i>	661
Howard Barrow Lambert, <i>Differential mappings on a vector space</i>	669
Richard G. Levin and Takayuki Tamura, <i>Notes on commutative power joined semigroups</i>	673
Robert Edward Lewand and Kevin Mor McCrimmon, <i>Macdonald's theorem for quadratic Jordan algebras</i>	681
J. A. Marti, <i>On some types of completeness in topological vector spaces</i>	707
Walter J. Meyer, <i>Characterization of the Steiner point</i>	717
Saad H. Mohamed, <i>Rings whose homomorphic images are q-rings</i>	727
Thomas V. O'Brien and William Lawrence Reddy, <i>Each compact orientable surface of positive genus admits an expansive homeomorphism</i>	737
Robert James Plemmons and M. T. West, <i>On the semigroup of binary relations</i>	743
Calvin R. Putnam, <i>Unbounded inverses of hyponormal operators</i>	755
William T. Reid, <i>Some remarks on special disconjugacy criteria for differential systems</i>	763
C. Ambrose Rogers, <i>The convex generation of convex Borel sets in euclidean space</i>	773
S. Saran, <i>A general theorem for bilinear generating functions</i>	783
S. W. Smith, <i>Cone relationships of biorthogonal systems</i>	787
Wolmer Vasconcelos, <i>On commutative endomorphism rings</i>	795
Vernon Emil Zander, <i>Products of finitely additive set functions from Orlicz spaces</i>	799
G. Sankaranarayanan and C. Suyambulingom, <i>Correction to: "Some renewal theorems concerning a sequence of correlated random variables"</i>	805
Joseph Zaks, <i>Correction to: "Trivially extending decompositions of E^n"</i>	805
Dong Hoon Lee, <i>Correction to: "The adjoint group of Lie groups"</i>	805
James Edward Ward, <i>Correction to: "Two-groups and Jordan algebras"</i>	806