

Pacific Journal of Mathematics

**THE SUSLIN-KLEENE THEOREM FOR V_κ WITH COFINALITY
 $(\kappa) = \omega$**

CHEN CHUNG CHANG AND YIANNIS (JOHN) NICOLAS MOSCHOVAKIS

THE SUSLIN-KLEENE THEOREM FOR V_κ WITH COFINALITY(κ) = ω

C. C. CHANG AND YANNIS N. MOSCHOVAKIS

It is easy to extend to arbitrary structures $\mathfrak{A} = \langle A, R_1, \dots, R_l f_1, \dots, f_m \rangle$ the concepts of Π_1^1 and inductively definable relations, which are familiar for the structure of the integers. The second author showed in a recent paper that these two concepts coincide for countable \mathfrak{A} that satisfy certain mild definability conditions—this is a generalization of the classical Suslin-Kleene theorem. Here we generalize the Suslin-Kleene theorem in a different direction.

MAIN RESULT. Let V_κ be the set of sets of rank less than κ , i.e., $V_0 = \phi$, $V_{\xi+1} = \text{power of } V_\xi$, $V_\kappa = \bigcup_{\xi < \kappa} V_\xi$, if κ is limit. The classes of inductively definable and Π_1^1 relations on the structure $\mathscr{V}_\kappa = \langle V_\kappa, \in \upharpoonright V_\kappa \rangle$ ($\kappa \geq \omega$) coincide if and only if κ is a limit ordinal with cofinality ω .

This implies several corollaries about the class of Π_1^1 relations on V_κ , when cofinality(κ) = ω , e.g., that it has the reduction property.

The nontrivial part of the theorem is the implication $\Pi_1^1 \Rightarrow \text{inductively definable}$ for \mathscr{V}_κ with cofinality(κ) = ω .

1. **Proof of the main result.** We assume familiarity with [7], whose notation we shall use.

Notice first that for each $\kappa \geq \omega$, \mathscr{V}_κ is an *acceptable* structure, in the sense of [7]. This is immediate for limit κ , by taking the ordinary set-theoretic pair and the standard ω for the integers within V_κ . For successor κ the proof is by induction; let $\kappa = \lambda + 1$, let $(,)_\lambda$ be a definable pair in \mathscr{V}_λ , for x_1, \dots, x_n in V_κ put

$$\langle x_1, \dots, x_n \rangle = \{(1, u)_\lambda : u \in x_1\} \cup \dots \cup \{(n, u)_\lambda : u \in x_n\}.$$

These n -tuple functions are definable in \mathscr{V}_κ and using them one can easily define a pair for \mathscr{V}_κ and also show that first-order definability on \mathscr{V}_κ is preserved under inductive definitions.

Since \mathscr{V}_κ is acceptable, the inductively definable relations on \mathscr{V}_κ are Π_1^1 by the argument given in §3 of [7]. Also, if $\kappa \geq \omega$ and κ is a successor or cofinality(κ) > ω , then the relation

$$S \in WF \Leftrightarrow \text{there is no sequence } u_0, u_1, \dots, \text{ so that} \\
(n)[(u_n, u_{n+1}) \in S]$$

is first-order definable on \mathscr{V}_κ , so that by the usual analysis of trans-

finite inductions "from within", each inductively definable relation is Σ_1^1 , and hence these relations do not exhaust Π_1^1 . To complete the proof we must show that if cofinality $(\kappa) = \omega$, then each Π_1^1 relation on \mathcal{V}_κ is inductively definable.

Let $P(x) \Leftrightarrow (\alpha)(Ey)Q(\alpha, y, x)$ be a typical Π_1^1 relation, where $Q(\alpha, y, x)$ is defined by the simple, quantifier-free formula $Q(\alpha, y, x)$ of \mathcal{L}^2 , let $t_1(x, y), \dots, t_d(x, y)$ be the finitely many terms s such that the term $\alpha(s)$ occurs in $Q(\alpha, y, x)$, let $t_1(x, y), \dots, t_d(x, y)$ be the functions on V_κ that these terms define, choose $Q^*(z, y, x)$ as in §4 of [7] so that

$$(1) \quad Q(\alpha, y, x) \Leftrightarrow Q^*(z, y, x)$$

whenever

$$(2) \quad \text{Seq}(z) \ \& \ K(z) = d \ \& \ (i)_{1 \leq i \leq d} [(z)_i = \alpha(t_i(x, y))].$$

We shall define for each $x \in V_\kappa$ a game $\mathcal{G}(x)$ so that when cofinality $(\kappa) = \omega$,

$$P(x) \Leftrightarrow \text{I has a winning strategy in } \mathcal{G}(x).$$

In the game $\mathcal{G}(x)$, player I chooses $a_1 (a_1 \in V_\kappa)$, player II chooses a pair $b_1, c_1 (b_1, c_1 \in V_\kappa)$, then I chooses a_2 , then II chooses b_2, c_2 , etc. We say that *the outlook is good for player II at step k* , when $a_1, \dots, a_k, b_1, c_1, \dots, b_k, c_k$ have been played, if the following conditions are satisfied.

(i) For each $i \leq k$, b_i is a function with domain ω and range power (a_i) , so that

$$\bigcup_{j \in \omega} b_i(j) = a_i.$$

(Thus II decomposes a_i into an ω -sequence of sets.)

(ii) c_k is a function with domain $\{(i, j) : i, j \leq k\}$ which assigns to each pair (i, j) a function $f_{i,j}^k$ with domain $(f_{i,j}^k) = b_i(j)$.

(iii) The union

$$f^k = \bigcup_{i,j \leq k} f_{i,j}^k$$

is a function.

(iv) There is no element $y \in V_\kappa$ such that all $t_1(x, y), \dots, t_d(x, y)$ are in the domain of f^k and such that (1) holds when we choose z so that (2) holds with $f^k(t_i(x, y))$ substituted for $\alpha(t_i(x, y))$, $i = 1, \dots, d$.

At the end of the game, player II wins if the outlook is good for him at every step k , otherwise player I wins.

LEMMA 1. *If cofinality $(\kappa) = \omega$ and player I has a winning strategy in $\mathcal{G}(x)$, then $(\alpha)(Ey)Q(\alpha, y, x)$.*

Proof. Given a function α on V_κ to V_κ , consider the game where I plays a_1, a_2, \dots following his winning strategy and II plays as follows. Since cofinality $(\kappa) = \omega$, we can choose a countable sequence v_1, v_2, \dots of elements of V_κ such that

$$V_\kappa = \bigcup_{j \in \omega} v_j.$$

For each k , the function $\alpha \upharpoonright a_k$ is a subset of V_κ and it can be decomposed into a countable union of subfunctions which are elements of V_κ ,

$$\alpha \upharpoonright a_k = \bigcup_{j \in \omega} ((\alpha \upharpoonright a_k) \cap v_j).$$

At step k , II chooses a b_k so that

$$b_k(j) = \text{domain}((\alpha \upharpoonright a_k) \cap v_j) \quad (j \leq k)$$

and a c_k so that

$$f_{i,j}^k = (a \upharpoonright \alpha_i) \cap v_j \quad (i, j, \leq k).$$

It is now clear that at each k , conditions (i), (ii), (iii) above are satisfied. Since I wins the game, there must be a k at which condition (iv) fails. For that k we have $Q^*(z, y, x)$ for some y and some z that codes a subfunction of α , so that by (1) we have $Q(\alpha, y, x)$ and the proof is complete.

LEMMA 2. *If cofinality $(\kappa) = \omega$ and $(\alpha)(Ey)Q(\alpha, y, x)$, then I has a winning strategy in $\mathcal{G}(x)$.*

Proof. Let I simply play $a_k = v_k$, where the v_j are elements of V_κ such that $V_\kappa = \bigcup_{j \in \omega} v_j$. Any winning sequence of plays for II determines a completely defined function α on V_κ to V_κ , so by hypothesis there is some y , so that $Q(\alpha, y, x)$. Now $y \in a_i$, for some i , and for large enough j , all $t_1(x, y), \dots, t_d(x, y)$ must be elements of $b_i(1) \cup \dots \cup b_i(j)$. It is then clear that the outlook is not good for II at step $k = \max(i, j)$, since condition (iv) will fail at that k .

Proof of the main result from these two lemmas is just like the proof in §5 of [7] and we shall omit it. The key points are that the game $\mathcal{G}(x)$ is *open* (i.e., if I wins, then he knows it at some point k of the game) and that conditions (i)-(iv) are first-order definable on \mathcal{V}_κ .

The result can be easily relativized to relations on *functions* on V_κ to V_κ as in §6 of [7]. One can also imitate the argument of §7 of [7] to show that the result cannot be proved by the classical method of representing \prod_1^1 relations via the property of well-foundedness.

We start with some λ with cofinality $(\lambda) > \omega$ and then use the Montague-Vaught method of [4] to find a $\kappa < \lambda$, with cofinality $(\kappa) = \omega$ and such that for some $C' \subseteq \text{power}(V_\lambda)$, the structure $\langle V_\kappa, C', \varepsilon \rangle$ is an elementary substructure of $\langle V_\lambda, \text{power}(V_\lambda), \varepsilon \rangle$. It is then easy to show that some Π_1^1 relation $P(x)$ on \mathcal{V}_κ is not of the form

$$P(x) \Leftrightarrow \lambda uv Q(x, u, v) \text{ is well-founded,}$$

with first-order definable $Q(x, u, v)$.

2. Corollaries and comments. Let Γ be a class of relations on some acceptable structure \mathfrak{A} . We say that Γ is *parametrized* if there is a binary relation $G(z, x)$ in Γ , so that each unary relation $P(x)$ in Γ is of the form

$$P(x) \Leftrightarrow G(z_0, x)$$

for some fixed z_0 in the domain of the structure. It is easy to verify that the classes of Π_1^1, Σ_1^1 and inductively definable relations on an acceptable structure are parametrized.

Suppose Γ is parametrized by $G(z, x)$. Put

Prewellordering (Γ) \Leftrightarrow there is a function Ψ on (the extension of) G into some ordinal κ and relations \leq and $\dot{\leq}$ in Γ and $\neg \Gamma$ (= the class of negations of relations in Γ) respectively, so that

$$\begin{aligned} G(z, x) &\Leftrightarrow (u)(v)[(u, v) \dot{\leq} (z, x) \Leftrightarrow (u, v) \leq (z, x) \\ &\Leftrightarrow [G(u, v) \ \& \ \Psi(u, v) \leq \Psi(z, x)]] . \end{aligned}$$

It is well-known that if Γ satisfies reasonable closure conditions, then *Prewellordering* (Γ) implies that Γ satisfies many interesting structure properties -e.g., see [5], [3], [6]. One of them is

Reduction (Γ). Given relations $P(x), Q(x)$ in Γ , there exist relations $P_1(x), Q_1(x)$ in Γ such that

$$\begin{aligned} P_1(x) &\Leftrightarrow P(x), Q_1(x) \Leftrightarrow Q(x), \\ P(x) \vee Q(x) &\Leftrightarrow P_1(x) \vee Q_1(x), \\ (x) \neg [P_1(x) \ \& \ Q_1(x)] . \end{aligned}$$

Others include the existence of a hierarchy on $\Gamma \cap \neg \Gamma$, where Γ now must satisfy fairly strong closure properties.

Our main result here together with the results in [5] gives

$$\text{cofinality}(\kappa) = \omega \Rightarrow \text{Prewellordering}(\Pi_1^1(\mathcal{V}_\kappa)),$$

where $\Pi_1^1(\mathcal{V}_\kappa)$ is the class of Π_1^1 relations on \mathcal{V}_κ . Since $\Pi_1^1(\mathcal{V}_\kappa)$

satisfies all the required closure properties, this further gives *Reduction* ($\prod_1^1(\mathcal{V}_\kappa)$) and the existence of a hierarchy on $\Delta_1^1(\mathcal{V}_\kappa) = \prod_1^1(\mathcal{V}_\kappa) \cap \Sigma_1^1(\mathcal{V}_\kappa)$.

The classical arguments of Gödel and Addison [1], [2] suffice to show

$$[\text{Axiom of Constructibility \& } \kappa \text{ a successor or cofinality } (\kappa) > \omega] \Rightarrow \text{Prewellordering } (\Sigma_1^1(\mathcal{V}_\kappa)).$$

However we do not know how to settle *Prewellordering* ($\prod_1^1(\mathcal{V}_\kappa)$) or *Prewellordering* ($\Sigma_1^1(\mathcal{V}_\kappa)$) when κ is a successor or cofinality $(\kappa) > \omega$ in Zermelo-Fraenkel set theory or in extensions of that theory by strong axioms which do not restrict our conception of arbitrary set. The problem has been attacked without success by some people for the case $\kappa = \omega + 1$, corresponding to the class of \prod_1^2 or Σ_1^2 relations on the continuum in type-theoretic notation. We suspect that it may be easier to settle for limit κ with cofinality $(\kappa) > \omega$, perhaps for κ satisfying strong axioms of infinity. An optimist would hope that for each κ , one of $\prod_1^1(\mathcal{V}_\kappa)$ or $\Sigma_1^1(\mathcal{V}_\kappa)$ must satisfy the prewellordering property.

REFERENCES

1. J. W. Addison, *Separation principles in the hierarchies of classical and effective descriptive set theory*, Fund. Math. **46** (1959), 123-135.
2. ———, *Some consequences of the axiom of constructibility*, Fund. Math. **46** (1959), 337-357.
3. J. W. Addison and Yiannis N. Moschovakis, *Some consequences of the axiom of definable determinateness*, Proc. Nat. Acad. Sci. U.S.A. **59** (1968), 708-712.
4. R. M. Montague and R. L. Vaught, *Natural models for set theory*, Fund. Math. **47** (1959), 219-242.
5. Yiannis N. Moschovakis, *Abstract first order computability, II*, Trans. Amer. Math. Soc. **138** (1969), 465-504.
6. ———, *Determinacy and prewellorderings of the continuum*, Mathematical Logic and Foundations of Set Theory, Amsterdam (1970), 24-63.
7. ———, *The Suslin-Kleene theorem for countable structures* (to appear in Duke Math. J.)

Received September 26, 1969 and in revised form May 1, 1970. Preparation of this paper was sponsored in part by NSF Grant GP-8827.

UNIVERSITY OF CALIFORNIA, LOS ANGELES

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLE

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics

Vol. 35, No. 3

November, 1970

John D. Arrison and Michael Rich, <i>On nearly commutative degree one algebras</i>	533
Bruce Alan Barnes, <i>Algebras with minimal left ideals which are Hilbert spaces</i>	537
Robert F. Brown, <i>An elementary proof of the uniqueness of the fixed point index</i>	549
Ronn L. Carpenter, <i>Principal ideals in F-algebras</i>	559
Chen Chung Chang and Yiannis (John) Nicolas Moschovakis, <i>The Suslin-Kleene theorem for V_κ with cofinality $(\kappa) = \omega$</i>	565
Theodore Seio Chihara, <i>The derived set of the spectrum of a distribution function</i>	571
Tae Geun Cho, <i>On the Choquet boundary for a nonclosed subspace of $C(S)$</i>	575
Richard Brian Darst, <i>The Lebesgue decomposition, Radon-Nikodym derivative, conditional expectation, and martingale convergence for lattices of sets</i>	581
David E. Fields, <i>Dimension theory in power series rings</i>	601
Michael Lawrence Fredman, <i>Congruence formulas obtained by counting irreducibles</i>	613
John Eric Gilbert, <i>On the ideal structure of some algebras of analytic functions</i>	625
G. Goss and Giovanni Viglino, <i>Some topological properties weaker than compactness</i>	635
George Grätzer and J. Sichler, <i>On the endomorphism semigroup (and category) of bounded lattices</i>	639
R. C. Lacher, <i>Cell-like mappings. II</i>	649
Shiva Narain Lal, <i>On a theorem of M. Izumi and S. Izumi</i>	661
Howard Barrow Lambert, <i>Differential mappings on a vector space</i>	669
Richard G. Levin and Takayuki Tamura, <i>Notes on commutative power joined semigroups</i>	673
Robert Edward Lewand and Kevin Mor McCrimmon, <i>Macdonald's theorem for quadratic Jordan algebras</i>	681
J. A. Marti, <i>On some types of completeness in topological vector spaces</i>	707
Walter J. Meyer, <i>Characterization of the Steiner point</i>	717
Saad H. Mohamed, <i>Rings whose homomorphic images are q-rings</i>	727
Thomas V. O'Brien and William Lawrence Reddy, <i>Each compact orientable surface of positive genus admits an expansive homeomorphism</i>	737
Robert James Plemmons and M. T. West, <i>On the semigroup of binary relations</i>	743
Calvin R. Putnam, <i>Unbounded inverses of hyponormal operators</i>	755
William T. Reid, <i>Some remarks on special disconjugacy criteria for differential systems</i>	763
C. Ambrose Rogers, <i>The convex generation of convex Borel sets in euclidean space</i>	773
S. Saran, <i>A general theorem for bilinear generating functions</i>	783
S. W. Smith, <i>Cone relationships of biorthogonal systems</i>	787
Wolmer Vasconcelos, <i>On commutative endomorphism rings</i>	795
Vernon Emil Zander, <i>Products of finitely additive set functions from Orlicz spaces</i>	799
G. Sankaranarayanan and C. Suyambulingom, <i>Correction to: "Some renewal theorems concerning a sequence of correlated random variables"</i>	805
Joseph Zaks, <i>Correction to: "Trivially extending decompositions of E^n"</i>	805
Dong Hoon Lee, <i>Correction to: "The adjoint group of Lie groups"</i>	805
James Edward Ward, <i>Correction to: "Two-groups and Jordan algebras"</i>	806