

Pacific Journal of Mathematics

A GENERAL THEOREM FOR BILINEAR GENERATING FUNCTIONS

S. SARAN

A GENERAL THEOREM FOR BILINEAR GENERATING FUNCTIONS

S. SARAN

The following theorem was proved by Chatterjea for ultra-spherical polynomials:

$$\text{If } F(x, t) = \sum_{m=0}^{\infty} a_m t^m P_m^\lambda(x)$$

then

$$\bar{\rho}^{2\lambda} F\left(\frac{x-t}{\rho}, \frac{ty}{\rho}\right) = \sum_{r=0}^{\infty} t^r b_r(y) P_r^\lambda(x)$$

where

$$b_r(y) = \sum_{m=0}^{\infty} (v_m) a_m y^m \text{ and } \rho = (1 - 2xt + t^2)^{1/2}.$$

The object of this paper is to show that a general theorem for any polynomial satisfying certain conditions can be given so as to include the above case, and may be applicable in obtaining new generating functions for other polynomials also.

THEOREM. If

$$(1.1) \quad f_n(x) = \mu(n)G(x)D^n\{g(x)\}$$

where $g(x)$ and $G(x)$ are independent of n , and

$$(1.2) \quad F(x, t) = \sum_{m=0}^{\infty} a_m t^m f_m(x)$$

then

$$\frac{G(x)F(x-t, ty)}{G(x-t)} = \sum_{r=0}^{\infty} \frac{(-t)^r}{\mu(r)r!} b_r(y) f_r(x)$$

where

$$b_r(y) = \sum_{m=0}^r (-r)_m \mu(m) a_m y^m.$$

2. Proof of the theorem. Writing ty for t in (1.2) and using (1.1) we get

$$[G(x)]^{-1} F(x, ty) = \sum_{m=0}^{\infty} a_m t^m y^m \mu(m) D^m [g(x)].$$

Applying the operator \bar{e}^{tD} where $D \equiv d/dx$ on both sides, we get, since

$$\bar{e}^{tD} f(x) = f(x-t)$$

$$\begin{aligned} \frac{F(x-t, ty)}{G(x-t)} &= \sum_{m=0}^{\infty} a_m t^m y^m \mu(m) \bar{e}^{tD} D^m [g(x)] \\ &= \sum_{m=0}^{\infty} a_m t^m y^m \mu(m) \sum_{r=0}^{\infty} \frac{(-tD)^r}{r!} D^m [g(x)] \\ &= \sum_{m=0}^{\infty} a_m y^m \mu(m) \sum_{r=0}^{\infty} \frac{(-)^r t^{m+r}}{r!} D^{m+r} [g(x)] \\ &= \sum_{r=0}^{\infty} \frac{(-)^r t^r}{r!} D^r [g(x)] \sum_{m=0}^{\infty} (-r)_m a_m t^m y^m . \end{aligned}$$

This on using (1.1) proves the theorem.

3. Some applications of theorem. (i) First we consider the generating function given by Brafman

$$(1 - 2xt)^{-c} {}_2F_0\left(\frac{1}{2}C, \frac{1}{2}C + \frac{1}{2}, -; \frac{-4t^2}{(1 - 2xt)^2}\right) \cong \sum_{n=0}^{\infty} \frac{(c)_n H_n(x) t^n}{n!} .$$

If we take $(a)_m = (c)_m/m!$, $f_m(x) = H_m(x)$, $G(x) = \exp(x^2)$, $\mu(n) = (-)^n$ from Rodrigues formula [5] we obtain

$$\begin{aligned} [1 + 2yt(x-t)]^{-c} \exp(2xt - t^2) {}_2F_0\left(\frac{1}{2}c, \frac{1}{2}c + 1, -; \frac{-4y^2 t^2}{(1 + 2xyt - 2yt^2)^2}\right) \\ \cong \sum_{n=0}^{\infty} \frac{{}_2F_0(-n, c, -; y) H_n(x) t^n}{n!} , \end{aligned}$$

a relation proved by Brafman [1] and Rainville [5] in different ways.

(ii) We now consider the generating function given by Carlitz [2]

$$(3.1) \quad [1 + \frac{1}{2}(x+1)t]^{\alpha} [1 + \frac{1}{2}(x-1)t]^{\beta} = \sum_{n=0}^{\infty} P_n^{\alpha-n, \beta-n}(x) t^n .$$

Writing tz for t , multiplying both sides by $\bar{e}^z z^{c-1}$ and integrating w.r.t. z we get

$$\begin{aligned} \sum_{n=0}^{\infty} (c)_n P_n^{\alpha-n, \beta-n}(x) t^n \\ = \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{(-\alpha)_m (-\beta)_n (c)_{m+n}}{m! n!} [\frac{1}{2}(x+1)t]^m [\frac{1}{2}(x-1)t]^n . \end{aligned}$$

Similarly, writing t/p for t , multiplying both sides by $e^p p^{-D}/2\pi i$ and evaluating over the contour C given by Hankel [4] we get

$$(3.2) \quad \begin{aligned} F^{(1)}(c, D; -\alpha, -\beta; \frac{1}{2}(x+1)t, \frac{1}{2}(x-1)t) \\ = \sum_{n=0}^{\infty} \frac{(c)_n}{(D)_n} P_n^{\alpha-n, \beta-n}(x) t^n , \end{aligned}$$

where $F^{(1)}$ is Appell's hypergeometric function of the first kind [5]

defined by

$$F^{(1)}(\alpha, b; c, d; x, y) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(\alpha)_{m+n}(c)_m(d)_n}{(b)_{m+n}m!n!} x^m y^n \quad |x| < 1, |y| < 1.$$

Assuming that $f_n^{\alpha-n, \beta-n}(x) = (1/(1-x^2))p_n^{\alpha-n, \beta-n}(x)$ (3.2) can be put in the form

$$(3.3) \quad \begin{aligned} & F^{(1)}\left(c, D; -\alpha, -\beta; \frac{-t}{2(x-1)}, \frac{-t}{2(x+1)}\right) \\ &= \sum_{n=0}^{\infty} \frac{(c)_n}{(D)_n} f_n^{\alpha-n, \beta-n}(x) t^n. \end{aligned}$$

The differential formula for $f_n^{\alpha-n, \beta-n}(x)$ is [5]

$$(3.4) \quad f_n^{\alpha-n, \beta-n}(x) = \frac{(-)^n}{2^n n!} (1-x)^{-\alpha} (1+x)^{-\beta} D^n [(1-x)^\alpha (1+x)^\beta].$$

Thus comparing (3.3) and (3.4) with (1.1) and (1.2) we get

$$a_m = \frac{(c)_m}{(D)_m}, \mu(n) = \frac{(-)^n}{2^n n!}, G(x) = (1-x)^\alpha (1+x)^\beta$$

and therefore our theorem gives, after replacing t by $t(1-x^2)$,

$$(3.5) \quad \begin{aligned} & [1 + \frac{1}{2}t(x+1)]^\alpha [1 + \frac{1}{2}t(x-1)]^\beta \\ & \times F^{(1)}\left(C, D; -\alpha, -\beta; \frac{ty(x+1)}{2+t(x+1)}, \frac{ty(x-1)}{2+t(x-1)}\right) \\ &= \sum_{n=0}^{\infty} {}_2F_1(-n, c; D; -y) p_n^{\alpha-n, \beta-n}(x) t^n. \end{aligned}$$

Since $p_n^{\alpha-n, \beta-n}(x)$ can be reduced to $L_n^{\alpha-n}(x)$ by replacing x by $1-2x/\beta$ and letting β tending to infinity, the above result can be put in the following interesting form

$$(3.6) \quad \begin{aligned} & (1+t)^\alpha \exp(xt) \phi_1\left(C, -\alpha, D; -xty, \frac{ty}{t+1}\right) \\ &= \sum_{n=0}^{\infty} {}_2F_1(-n, C; D; -y) L_n^{\alpha-n}(x) \end{aligned}$$

where ϕ_1 is a confluent form of $F^{(1)}$ given by

$$\phi_1(\alpha, \beta, \gamma; x, y) = \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{(\alpha)_{m+n}(\beta)_n}{(\gamma)_{m+n}m!n!} x^m y^n, \quad |x| < 1.$$

This again reduces to the following result due to Srivastava [6]

$$(3.7) \quad \begin{aligned} & (1+t)^\alpha \exp(-yt) \phi_3\left(-\alpha, D; \frac{yt}{t+1}, xyt\right) \\ &= \sum_{n=0}^{\infty} \binom{D-1+n}{n}^{-1} L_n^{D-1}(y) L_n^{\alpha-n}(x) t^n \end{aligned}$$

where ϕ_3 is a confluent form of ϕ_1 . This is obtained by replacing y by $-(y/C)$ in (3.4) and letting C tend to infinity.

4. **Generalization of (3.2).** The result (3.1) is capable of generalization by the repeated application of the method indicated in proving (3.2). One can easily write it as

$$(4.1) \quad \begin{aligned} &F_{0,0}^{r,1}(|a|_r; -\alpha, -\beta; \frac{1}{2}(x+1)t, \frac{1}{2}(x-1)t) \\ &= \sum_{n=0}^{\infty} \frac{(a_1)_n \cdots (a_r)_n}{(b_1)_n \cdots (b_s)_n} p_n^{\alpha-n, \beta-n}(x) t^n \end{aligned}$$

in the notation of Srivastava and Saran [7] for Kampé de Fériet function.

Thus (3.5) can be generalized into

$$(4.2) \quad \begin{aligned} &[1 + \frac{1}{2}t(x+1)]^\alpha [1 + \frac{1}{2}t(x-1)]^\beta \\ &\times F_{s,0}^{r,1}(|a|_r; -\alpha, -\beta; \frac{yt(x+1)}{2+t(x+1)}, \frac{yt(x-1)}{2+t(x-1)}) \\ &= \sum_{n=0}^{\infty} {}_{r+1}F_s(-n, a_r; b_s; -y) p_n^{\alpha-n, \beta-n}(x) t^n . \end{aligned}$$

REFERENCES

1. F. Brafman, *Some generating functions for Laguerre and Hermite polynomials*, *Canad. J. Math.* **9** (1957), 180-187.
2. L. Carlitz, *A bilinear generating function for the Jacobi polynomials*, *Boll. Un. Math. Italy* (3) **18** (1963), 87.
3. S. K. Chatterjea, *A bilateral generating function for the Ultraspherical polynomials*, *Pacific J. Math.* **29** (1969), 73-76.
4. E. T. Copson, *An Introduction to Theory of Function of a Complex Variable*, Oxford University Press, (1935), 226.
5. E. D. Rainville, *Special functions*, Macmillan Co., New York, 1960.
6. H. M. Srivastava, *An extension of the Hille-Hardy formula*, *Math. Comp.* **23** (1969), 310.
7. G. P. Srivastava and S. Saran, *A theorem on Kampé de Fériet function*, *Proc. Camb. Phil. Soc.* **64** (1968), 435.

Received April 7, 1970.

PUNJABI UNIVERSITY
PATIALA, (INDIA)

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLE

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

John D. Arrison and Michael Rich, <i>On nearly commutative degree one algebras</i>	533
Bruce Alan Barnes, <i>Algebras with minimal left ideals which are Hilbert spaces</i>	537
Robert F. Brown, <i>An elementary proof of the uniqueness of the fixed point index</i>	549
Ronn L. Carpenter, <i>Principal ideals in F-algebras</i>	559
Chen Chung Chang and Yiannis (John) Nicolas Moschovakis, <i>The Suslin-Kleene theorem for V_κ with cofinality $(\kappa) = \omega$</i>	565
Theodore Seio Chihara, <i>The derived set of the spectrum of a distribution function</i>	571
Tae Geun Cho, <i>On the Choquet boundary for a nonclosed subspace of $C(S)$</i>	575
Richard Brian Darst, <i>The Lebesgue decomposition, Radon-Nikodym derivative, conditional expectation, and martingale convergence for lattices of sets</i>	581
David E. Fields, <i>Dimension theory in power series rings</i>	601
Michael Lawrence Fredman, <i>Congruence formulas obtained by counting irreducibles</i>	613
John Eric Gilbert, <i>On the ideal structure of some algebras of analytic functions</i>	625
G. Goss and Giovanni Viglino, <i>Some topological properties weaker than compactness</i>	635
George Grätzer and J. Sichler, <i>On the endomorphism semigroup (and category) of bounded lattices</i>	639
R. C. Lacher, <i>Cell-like mappings. II</i>	649
Shiva Narain Lal, <i>On a theorem of M. Izumi and S. Izumi</i>	661
Howard Barrow Lambert, <i>Differential mappings on a vector space</i>	669
Richard G. Levin and Takayuki Tamura, <i>Notes on commutative power joined semigroups</i>	673
Robert Edward Lewand and Kevin Mor McCrimmon, <i>Macdonald's theorem for quadratic Jordan algebras</i>	681
J. A. Marti, <i>On some types of completeness in topological vector spaces</i>	707
Walter J. Meyer, <i>Characterization of the Steiner point</i>	717
Saad H. Mohamed, <i>Rings whose homomorphic images are q-rings</i>	727
Thomas V. O'Brien and William Lawrence Reddy, <i>Each compact orientable surface of positive genus admits an expansive homeomorphism</i>	737
Robert James Plemmons and M. T. West, <i>On the semigroup of binary relations</i>	743
Calvin R. Putnam, <i>Unbounded inverses of hyponormal operators</i>	755
William T. Reid, <i>Some remarks on special disconjugacy criteria for differential systems</i>	763
C. Ambrose Rogers, <i>The convex generation of convex Borel sets in euclidean space</i>	773
S. Saran, <i>A general theorem for bilinear generating functions</i>	783
S. W. Smith, <i>Cone relationships of biorthogonal systems</i>	787
Wolmer Vasconcelos, <i>On commutative endomorphism rings</i>	795
Vernon Emil Zander, <i>Products of finitely additive set functions from Orlicz spaces</i>	799
G. Sankaranarayanan and C. Suyambulingom, <i>Correction to: "Some renewal theorems concerning a sequence of correlated random variables"</i>	805
Joseph Zaks, <i>Correction to: "Trivially extending decompositions of E^n"</i>	805
Dong Hoon Lee, <i>Correction to: "The adjoint group of Lie groups"</i>	805
James Edward Ward, <i>Correction to: "Two-groups and Jordan algebras"</i>	806