

Pacific Journal of Mathematics

CHARACTERS AND ORTHOGONALITY IN FROBENIUS ALGEBRAS

TIMOTHY V. FOSSUM

CHARACTERS AND ORTHOGONALITY IN FROBENIUS ALGEBRAS

T. V. FOSSUM

Matrix-theoretical proofs of orthogonality relations for the coefficients of representations of Frobenius algebras have been extensively developed in the literature. This paper has grown out of a desire to prove some of these orthogonality relations in a matrix-free, module-theoretic manner. Most of the attention is focused on characters, even though some of the results diverge from this restriction.

Sections 1 and 2 set the stage for the development of the principal theorems in § 3. There we derive the orthogonality relations and demonstrate a relation between characters and certain homogeneous modules over Frobenius algebras. In § 4 we apply these results to obtain some information about the character of the left regular module.

All rings are assumed to have an identity, and all modules are assumed to be unital. The Jacobson radical of a ring A will be written $J(A)$, or simply J .

Let A be a ring, M a simple left A -module, and let $A_M = \{a \in A: aM = 0\}$. Then A_M is a two-sided ideal of A , the *annihilator* of M . If L is a left A -module, $\text{Soc}_M(L)$ will denote the M -*socle* of L , *i.e.*, the sum of all submodules of L isomorphic to M . We say L is M -*homogeneous* if $\text{Soc}_M(L) = L$. In particular if A is left artinian, then A_M is a maximal two-sided ideal of A , and L is M -homogeneous if and only if $A_M L = 0$. We let S_M denote the M -socle $\text{Soc}_M(A)$ of A . A *block* of A is an indecomposable ring direct summand of A [2, § 55]. We say M *belongs to* the block B of A if M is a composition factor of B regarded as a left A -module.

Let K be a field. If V is a vector space over K , let $(V:K)$ denote the K -dimension of V . Now assume A is a finite dimensional K -algebra. Then $A^* = \text{Hom}_K(A, K)$ is an (A, A) -bimodule, where for $a \in A$ and $\lambda \in A^*$ we define

$$\begin{cases} (a\lambda)(x) = \lambda(xa) \\ (\lambda a)(x) = \lambda(ax) \end{cases} \quad (x \in A)$$

If $\lambda \in A^*$, we say λ is a *class function* if $\lambda(ab) = \lambda(ba)$ for all $a, b \in A$. Let $\text{cf}(A)$ denote the set of class functions in A^* . Observe that $\lambda \in \text{cf}(A)$ if and only if $a\lambda = \lambda a$ for all $a \in A$. We say $\chi \in A^*$ is an *A-character* if χ is the character of a (finite dimensional) left A -module. Clearly all A -characters belong to $\text{cf}(A)$. The A -character χ is said to be *irreducible* if it is the character of a simple left A -module.

DEFINITION. Let A be a finite-dimensional K -algebra, K a field, and let M be a simple left A -module. We say an element $\lambda \in A^*$ belongs to M if $\lambda(A_M) = 0$. One checks that if λ belongs to M , then $a\lambda$ and λa belong to M for any $a \in A$. If χ is the character of M , then χ clearly belongs to M .

All other notation may be found in [2] or [3].

1. Socles of QF rings. A ring A is said to be a QF (quasi-Frobenius) ring if A is (left and right) artinian and ${}_A A$ is injective [2, (58.6)].

LEMMA 1.1. *Let A be a QF ring, M a simple left A -module, and $S_M = \text{Soc}_M(A)$. Then S_M is a nonzero simple two-sided ideal of A .*

Proof. By [2, (58.13)], M is isomorphic to a minimal left ideal I in A , so $S_M \neq 0$. Moreover $S_M = \sum \{f(I) : f \in \text{Hom}_A(I, A)\}$ since $I \cong M$. By assumption ${}_A A$ is injective and from the "injective test lemma" [2, (57.14)] it follows that each $f \in \text{Hom}_A(I, A)$ is right multiplication by some $a \in A$; that is, $f(I) = Ia \subseteq IA$. Therefore $S_M \subseteq IA$. But plainly $IA \subseteq S_M$, and the lemma follows.

LEMMA 1.2. *Let A be a QF ring, M a simple left A -module, and $S_M = \text{Soc}_M(A)$. Then the following are equivalent.*

- (a) S_M is a simple block of A to which M belongs
- (b) $S_M^2 \neq 0$
- (c) M is projective.

Proof. That (a) implies (b) is obvious. For (b) implies (c), suppose M is not projective. By [2, (58.12)] M is isomorphic to the "bottom" constituent I of a principal indecomposable left ideal U , and as such, $I \subseteq JU$ where J is the radical of A . Hence $I \subseteq J$, and since $I \subseteq S_M$, $S_M \subseteq J$. But S_M is M -homogeneous and $J \subseteq A_M$ so therefore

$$S_M^2 \subseteq A_M S_M = 0.$$

To show that (c) implies (a), notice that if M is projective, then M is injective [2, (58.14)] and therefore M is a direct summand of every module with M as a composition factor. Therefore every principal indecomposable module linked to M is isomorphic to M , so S_M is a block of A [2, § 55]. From (1.1) it follows that S_M is simple, completing the proof.

2. Frobenius algebras and class functions. Let A be a finite-dimensional K -algebra, K a field. We say that A is a Frobenius algebra if $A \cong A^* = \text{Hom}_K(A, K)$ as left A -modules. To make this

connection more explicit, we say that the pair (A, φ) is a *Frobenius algebra* if $\varphi: A \rightarrow A^*$ is a left A -isomorphism (see [4] and [2, (61.1)].) By [2, (61.1), (61.2)] a Frobenius algebra is necessarily a *QF* ring; hence the results of §1 apply.

Let (A, φ) be a Frobenius algebra. If $(a_i), (b_i)$ is an ordered pair of bases for A such that $\varphi(b_i)(a_j) = \delta_{ij}$ for all i, j , we say the bases $(a_i), (b_i)$ are φ -dual. If (a_i) is any basis for (A, φ) , there exists a second basis (b_i) such that $(a_i), (b_i)$ are φ -dual. (In most cases we are given an explicit isomorphism φ and certain φ -dual bases, although for an arbitrary Frobenius algebra A there may be many different choices for φ or φ -dual bases.)

LEMMA 2.1. *Let $(a_i), (b_i)$ be φ -dual bases for the Frobenius algebra (A, φ) , and assume $\lambda \in A^*$. Then $\varphi^{-1}(\lambda) = \sum_i \lambda(a_i)b_i$.*

Proof. The proof is left to the reader.

LEMMA 2.2. *Let (A, φ) be a Frobenius algebra. Then the mapping $\tau = \tau_\varphi: A \rightarrow A$ given by $\tau(a) = \varphi^{-1}(\varphi(1)a)$ is a K -algebra automorphism of A .*

Proof. Observe that $\varphi(\tau(a)) = \varphi(1)a$ for each $a \in A$. Because φ is a left A -homomorphism and A^* is an (A, A) -bimodule, it follows easily that

$$(2.3) \quad \varphi(x\tau(a)) = \varphi(x)a, \quad (x, a \in A).$$

If $x = \varphi^{-1}(\lambda)$ for $\lambda \in A^*$, taking φ^{-1} on both sides of (2.3) gives

$$(2.4) \quad \varphi^{-1}(\lambda)\tau(a) = \varphi^{-1}(\lambda)a, \quad (a \in A).$$

Now for $a, b \in A$, $\varphi(\tau(ab)) = \varphi(1)ab = \varphi(\tau(a))b = \varphi(\tau(a)\tau(b))$ by (2.3), and therefore since φ is one-to-one, $\tau(ab) = \tau(a)\tau(b)$. If $\tau(a) = 0$, then $0 = \varphi(\tau(a)) = \varphi(1)a$, so $\varphi(b)a = b(\varphi(1)a) = 0$ for all $b \in A$. But then $0 = (\varphi(b)a)(1) = \varphi(b)(a)$ for all $b \in A$, and it follows that $a = 0$ since $\varphi(A) = A^*$. Therefore τ is one-to-one. Verification of the remaining properties of τ is left to the reader.

The reader may observe that $\tau = \tau_\varphi$ is the inverse of “Nakayama’s automorphism” [5, Th. 1].

DEFINITION. Let (A, φ) be a Frobenius algebra with automorphism $\tau = \tau_\varphi$ as in (2.2). Define $Z_\tau(A) = \{a \in A: ba = a\tau(b) \text{ for all } b \in A\}$.

LEMMA 2.5. *Let (A, φ) be a Frobenius algebra with automorphism $\tau = \tau_\varphi$ as in (2.2). Then $\varphi^{-1}(\text{cf}(A)) = Z_\tau(A)$.*

Proof. If $a \in Z_\tau(A)$, then $ba = a\tau(b)$ for all $b \in A$. Then by (2.3), $b\varphi(a) = \varphi(ba) = \varphi(a\tau(b)) = \varphi(a)b$ for all $b \in B$, so $\varphi(a) \in \text{cf}(A)$. The converse is similar.

COROLLARY 2.6. *Let (A, φ) , τ be as in (2.5). If $\lambda \in \text{cf}(A)$ and $\lambda^* = \varphi^{-1}(\lambda)$, then $A\lambda^* = \lambda^*A$.*

Proof. By (2.5), $\lambda^* \in Z_\tau(A)$. Since τ is an automorphism of A , $A\lambda^* = \lambda^*\tau(A) = \lambda^*A$.

3. Orthogonality relations, socles and characters. *Throughout this section let (A, φ) be a Frobenius algebra. Recall that if M is a simple left A -module, then $A_M = \{a \in A : aM = 0\}$, and for $\lambda \in A^*$, λ belongs to M if $\lambda(A_M) = 0$.*

LEMMA 3.1. *Let M be a simple left A -module, and assume λ belongs to M , $\lambda \in A^*$. Set $\lambda^* = \varphi^{-1}(\lambda)$. Then $A_M\lambda^* = 0$. In particular if N is any left A -module, then $A\lambda^*N$ is M -homogeneous.*

Proof. If $b \in A_M$, then $b\lambda = 0$ because $(b\lambda)(A) = \lambda(Ab) \subseteq \lambda(A_M) = 0$. Therefore $b\lambda^* = \varphi^{-1}(b\lambda) = 0$, $b \in A_M$, and so $A_M\lambda^* = 0$. For N a left A -module, $A_M(A\lambda^*N) = A_M\lambda^*N = 0$, so $A\lambda^*N$ is M -homogeneous.

THEOREM 3.2. (Orthogonality relations). *Let M and N be simple left A -modules, and assume λ, μ belong to M, N , respectively, where $\lambda, \mu \in A^*$. Set $\lambda^* = \varphi^{-1}(\lambda)$. If $\lambda^*N \neq 0$, then $M \cong N$. In particular if $\mu(\lambda^*) \neq 0$, then $M \cong N$.*

Proof. Suppose $\lambda^*N \neq 0$. Since N is simple, $A\lambda^*N = N$. By (3.1), N must then be M -homogeneous, i.e., $M \cong N$. In particular if $\mu(\lambda^*) \neq 0$, then plainly $\lambda^*N \neq 0$, so by above $M \cong N$.

COROLLARY 3.3. *Let M, N , λ, μ be as in (3.2), and set*

$$\lambda^* = \varphi^{-1}(\lambda), \mu^* = \varphi^{-1}(\mu).$$

If $\lambda^\mu^* \neq 0$, then $M \simeq N$.*

Proof. Assume $\lambda^*\mu^* \neq 0$. Then $\lambda^*\mu \neq 0$, and therefore for some $a \in A$, $0 \neq (\lambda^*\mu)(a) = \mu(a\lambda^*) = (\mu a)(\lambda^*)$. But μa belongs to N , so by (3.2), $M \cong N$.

COROLLARY 3.4. (Orthogonality relations for characters). *Let M, N be simple left A -modules with characters χ and ζ , respectively. Set $\chi^* = \varphi^{-1}(\chi)$. If $\zeta(\chi^*) \neq 0$, then $M \cong N$ and $\chi = \zeta$. In particular if $(a_i), (b_i)$ are φ -dual bases for A , and if $\sum_i \chi(a_i)\zeta(b_i) \neq 0$, then $M \cong N$ and $\chi = \zeta$.*

Proof. The first part follows directly from (3.2) since the character of a simple module belongs to that module. Moreover modules uniquely determine their characters. For the second part we need only apply this and (2.1).

THEOREM 3.5. *Let χ be the character of the simple left A -module M , and assume $\chi \neq 0$. Set $\chi^* = \varphi^{-1}(\chi)$. Then $A\chi^* = \text{Soc}_M(A) = S_M$. In particular either $(A\chi^*)^2 = 0$, or $A\chi^*$ is a simple block of A .*

Proof. Since $\chi \in \text{cf}(A)$, (2.6) implies that $A\chi^* = \chi^*A = A\chi^*A$. By (3.1), $A\chi^*$ is M -homogeneous, so $0 \neq A\chi^* \subseteq S_M$. Since S_M is a simple two-sided ideal (1.1), $A\chi^* = S_M$. The second assertion follows from this and (1.2).

COROLLARY 3.6. *Let χ^* be as in (3.5). If $(\chi^*)^2 = 0$, then $S_M^2 = 0$. If $(\chi^*)^2 \neq 0$, then χ^* is a unit in the simple block $A\chi^* = S_M$.*

Proof. By (3.5), $A\chi^* = S_M$. If $(\chi^*)^2 = 0$, then by (2.6),

$$S_M^2 = A\chi^*A\chi^* = A(\chi^*)^2 = 0.$$

On the other hand if $(\chi^*)^2 \neq 0$, then $S_M^2 \neq 0$, so by (1.2) $S_M = A\chi^*$ is a simple block of A in which χ^* is clearly a unit.

COROLLARY 3.7. *Let χ, χ^* be as in (3.5). If $\chi(\chi^*) \neq 0$, then χ^* is a unit in the block $A\chi^* = S_M$.*

Proof. If $\chi(\chi^*) \neq 0$, then $\chi^*\chi \neq 0$ (in A^*), so by applying φ^{-1} we have $(\chi^*)^2 = \chi^*\varphi^{-1}(\chi) = \varphi^{-1}(\chi^*\chi) \neq 0$. By (3.6), χ^* is a unit in S_M .

REMARK. The converse to (3.7) is false. For example let F be a field of characteristic $p \neq 0$, and let F_p be the complete ring of p -by- p matrices over F . Then F_p is a Frobenius algebra with isomorphism φ given by $\varphi(X)(Y) = \text{trace}(YX)$, $X, Y \in F_p$. One checks that if χ is the character of the unique simple left F_p -module M , then $\chi^* = \varphi^{-1}(\chi) = I_p$ where I_p is the p -by- p identity matrix. Then $\chi(\chi^*) = p = 0$ since F has characteristic p , but χ^* is clearly a unit in F_p .

The procedure for finding the socle of the left regular module ${}_A A$ of a Frobenius algebra (3.5) may also be applied to arbitrary left A -modules and in particular to principal indecomposable modules [2, (54.3)].

PROPOSITION 3.8. *Let M, χ^* be as in (3.5), and assume N is any left A -module. If $(\chi^*)^2 \neq 0$, then $\chi^*N = \text{Soc}_M(N)$. If $(\chi^*)^2 = 0$, then $\chi^*N \subseteq \text{Soc}_M(JN)$ where J is the radical of A .*

Proof. Assume $(\chi^*)^2 \neq 0$. Then $S_M = \chi^*A$ is the simple block of A to which M belongs (3.6), and $S_M + A_M = A$. If L is any M -homogeneous left A -module, then $L = AL = S_M L = \chi^*L$. By (3.1), $A\chi^*N = \chi^*N$ is M -homogeneous, so $\chi^*N \subseteq \text{Soc}_M(N)$. But by above $\text{Soc}_M(N) = \chi^* \text{Soc}_M(N) \subseteq \chi^*N$. Therefore $\chi^*N = \text{Soc}_M(N)$. Now if $(\chi^*)^2 = 0$, then $\chi^* \in J = J(A)$ (since $\chi^* \in S_M$ and $S_M^2 = 0$ by (3.6)) so that $\chi^*N \subseteq JN$. But χ^*N is M -homogeneous by (3.1), and therefore $\chi^*N \subseteq \text{Soc}_M(JN)$.

EXAMPLE. Equality does not necessarily occur in the relation $\chi^*N \subseteq \text{Soc}_M(JN)$ when $(\chi^*)^2 = 0$. For example let G be an abelian group of order a prime $p > 2$, and let F be a field of characteristic p . Then the group-algebra FG is a commutative local ring which is clearly a Frobenius algebra [2, Remark 2, p. 440] with isomorphism φ , say. Moreover if J denotes the radical of FG , then $J^p = 0$ and $J^{p-1} = \text{Soc}(FG)$.

Let M be a simple left FG -module with character χ . Then all simple left FG -modules are isomorphic to M , and in particular

$$J^{p-1} = \text{Soc}_M(FG) \cong M.$$

If $\chi^* = \varphi^{-1}(\chi)$, then $\chi^* \in \text{Soc}_M(FG) = J^{p-1}$, and $(\chi^*)^2 = 0$. Let $N = J$, viewed as a left FG -module. Then $\chi^*N \subseteq J^{p-1}J = J^p = 0$, so $\chi^*N = 0$. But $\text{Soc}_M(JN) = \text{Soc}_M(J^2) = J^{p-1} \neq 0$. Hence $\chi^*N \neq \text{Soc}_M(JN)$.

THEOREM 3.9. *Let f be a primitive idempotent in A , and let M be a simple left A -module with character $\chi \neq 0$. Set $\chi^* = \varphi^{-1}(\chi)$. Then $\chi^*f \neq 0$ if and only if $\chi^*Af = \text{Soc}(Af) \cong M$.*

Proof. By [2, (58.12)], $\text{Soc}(Af)$ is simple. If $\chi^*f \neq 0$, then $0 \neq A\chi^*Af = \chi^*Af \subseteq \text{Soc}_M(Af) \subseteq \text{Soc}(Af)$ by (2.6) and (3.1), so

$$\chi^*Af = \text{Soc}(Af) \cong M.$$

Conversely if $\chi^*Af = \text{Soc}(Af)$, then $A\chi^*f = \chi^*Af \neq 0$ by (2.6), and therefore $\chi^*f \neq 0$.

4. Applications. *In this section (A, φ) is a Frobenius algebra, and ρ is the character of the left regular module ${}_A A$.*

LEMMA 4.1. *Let $(a_i), (b_i)$ be φ -dual bases for (A, φ) . Then*

$$\varphi^{-1}(\rho) = \sum_i a_i b_i.$$

Proof. This proof is identical to that of [3, (3.3)].

Let M_1, \dots, M_s be a complete set of (nonisomorphic) simple left A -modules with characters χ_1, \dots, χ_s , respectively. Let m_i denote the multiplicity of M_i as a composition factor of ${}_A A$. It follows that $\rho = \sum_{i=1}^s m_i \chi_i$. If we set $\rho^* = \varphi^{-1}(\rho)$ and $\chi_i^* = \varphi^{-1}(\chi_i)$ for $1 \leq i \leq s$, then

$$(4.2) \quad \rho^* = \sum_{i=1}^s m_i \chi_i^* .$$

Now assume χ_k is any irreducible A -character, $1 \leq k \leq s$. For any $c \in A$, $c\chi_k$ belongs to M_k , and by (4.2),

$$(4.3) \quad \chi_k(\rho^*c) = (c\chi_k)(\rho^*) = \sum_{i=1}^s m_i (c\chi_k)(\chi_i^*) .$$

But by (3.2), $(c\chi_k)(\chi_i^*) = 0$ unless $M_i \cong M_k$, i.e., unless $i = k$. Therefore (4.3) implies that $\chi_k(\rho^*c) = m_k (c\chi_k)(\chi_k^*) = m_k \chi_k(\chi_k^*c)$. Together with (2.1) and (4.1) this gives the following (compare [3, (3.12)]).

THEOREM 4.4. *Let ρ be the character of the left regular module ${}_A A$, and let M be a simple left A -module with character χ . Set $\chi^* = \varphi^{-1}(\chi)$. Assume M appears as a composition factor of ${}_A A$ with multiplicity m . Then for any $c \in A$,*

$$\chi(\rho^*c) = m\chi(\chi^*c) .$$

In particular if $(a_i), (b_i)$ are φ -dual bases for (A, φ) , then for any $c \in A$,

$$\chi\left(\sum_i a_i b_i c\right) = m \sum_i \chi(a_i) \chi(b_i c) .$$

COROLLARY 4.5. *Let G be a finite group of order $|G|$, and let K be a field of characteristic p such that p does not divide $|G|$. Assume that K is a splitting field for KG . Then for any irreducible KG -character χ and for any $h \in G$,*

$$|G|\chi(h) = \chi(1) \sum_{g \in G} \chi(g)\chi(g^{-1}h) .$$

Proof. (Compare [2, (31.6)].) Let M be the simple left KG -module with character χ . Since KG is semi-simple (Maschke) and K -split, M appears exactly $(M:K)$ times as a composition factor of KG , and $\chi(1) = (M:K)$ in K . Finally KG is a Frobenius algebra with respect to the particular isomorphism φ for which $(g), (g^{-1})$ are φ -dual bases. By (4.4), $|G|\chi(h) = \chi(\sum_{g \in G} gg^{-1}h) = \chi(1) \sum_{g \in G} \chi(g)\chi(g^{-1}h)$ as desired.

Assume that B is a simple, finite-dimensional K -algebra, K a field, and let M be a simple left B -module. Set $D = \text{Hom}_B(M, M)$.

Then D is a division ring, finite-dimensional over K , and B is isomorphic to D_m , the full ring of m -by- m matrices over D , for some positive integer m . It follows that M appears m times as a composition factor of ${}_B B$, and that $(M:K)m = (D:K)m^2$. Therefore $(M:K) = (D:K)m$. We combine these remarks with (4.4) to obtain the following.

COROLLARY 4.6. *Assume (A, φ) is a Frobenius algebra over a field K of characteristic zero, and let M be a simple projective left A -module with character χ . Set $\chi^* = \varphi^{-1}(\chi)$, $\rho^* = \varphi^{-1}(\rho)$, and $D = \text{Hom}_A(M, M)$. Then for some $c \in A$, $\chi(\rho^*c) \neq 0$, and*

$$(D:K) = \chi(\rho^*c)^{-1}\chi(1)\chi(\chi^*c).$$

In particular if $(a_i), (b_i)$ are φ -dual bases for (A, φ) , then for some $c \in A$, $\chi(\sum_i a_i b_i c) \neq 0$ and for any such c ,

$$(D:K) = \chi\left(\sum_i a_i b_i c\right)^{-1} \chi(1) \sum_i \chi(a_i)\chi(b_i c).$$

Proof. Because M is projective and A is a QF ring, the M -socle S_M of A is a simple block of A to which M belongs by (1.2). Moreover $\chi(1) = (M:K)$ in K , and since K has characteristic zero, $\chi \neq 0$. By (3.6), $S_M = \chi^*A$ and $A = S_M + A_M$, so $0 \neq \chi(A) = \chi(S_M) = \chi(\chi^*A)$. Therefore for some $c \in A$, $\chi(\chi^*c) \neq 0$. But (4.4) shows that $\chi(\rho^*c) = m\chi(\chi^*c)$ where m is the multiplicity of M as a composition factor of ${}_A A$, and therefore $\chi(\rho^*c) \neq 0$. The remarks above the statement of this corollary show that $(M:K) = (D:K)m$. Combining these results, we have that

$$(D:K) = \chi(\rho^*c)^{-1}\chi(1)\chi(\chi^*c).$$

The last statement follows from this, (2.1) and (4.1).

REMARK. The hypothesis that M be projective is essential in the above corollary, for if M is not projective (equivalently, if $(\chi^*)^2 = 0$) one can show from (1.2) and (3.6) that $\chi(\chi^*c) = 0$ for all $c \in A$.

COROLLARY 4.7. *Hypotheses as in (4.6). If $(a_i), (b_i)$ are φ -dual bases for A such that $\sum_i a_i b_i = (A:K)$, then*

$$(D:K) = (A:K)^{-1} \sum_i \chi(a_i)\chi(b_i).$$

Proof. Certainly $\chi(\rho^*) = \chi(\sum_i a_i b_i) = (A:K)\chi(1) \neq 0$. Therefore we may take $c = 1$ in the proof of (4.6), and the conclusion follows.

REMARKS. One might be surprised to observe that the hypotheses of (4.7) forces (A, φ) to be a semi-simple symmetric algebra [3]. For we know that $\rho \in \text{cf}(A)$, so by (2.5) and (4.1), $(A:K) = \sum_i a_i b_i = \varphi^{-1}(\rho) \in Z_\tau(A)$. But then $1 \in Z_\tau(A)$ and therefore for any $b \in A$, $b = b \cdot 1 = 1 \cdot \tau(b) = \tau(b)$. We conclude that τ is the identity automorphism which forces φ to be an (A, A) -bimodule isomorphism (2.3). Therefore (A, φ) is a symmetric algebra. Semi-simplicity follows from [3, (1.7)].

If G is a finite group, K a field of characteristic zero which is a splitting field for KG (e.g., K algebraically closed), then the sequence (2.1), (3.1), (3.2), (3.4), (4.1), (4.4) and (4.5) gives an elementary matrix-free proof of the orthogonality relations for irreducible characters of G .

The references below include a partial bibliography of papers relating to orthogonality relations for Frobenius algebras.

REFERENCES

1. R. Brauer, *On hypercomplex arithmetic and a theorem of Speiser*, Festschrift für Speiser, Zurich, 1945.
2. C. W. Curtis and I. Reiner, *Representation Theory of Finite Groups and Associative Algebras*, Interscience, New York, 1962.
3. T. V. Fossum, *Characters and centers of symmetric algebras*, J. of Algebra **16** (1970), 4-13.
4. T. Nakayama, *On Frobeniusean algebras I*, Math. Ann. **40** (1939), 611-633.
5. ———, *On Frobeniusean algebras II*, Math. Ann. **42** (1941) 1-21.
6. ———, *Orthogonality relations for Frobenius and quasi-Frobenius-algebras*, Proc. Amer. Math. Soc. **3** (1952), 183-195.
7. C. Nesbitt and R. M. Thrall, *Some ring theorems with applications to modular representations*, Ann. of Math. **47** (1946) 551-567.
8. M. Osima, *On the Schur relations for the representations of a Frobenius algebra*, J. Math. Soc. of Japan **4** (1952) 1-13.
9. ———, *Supplementary remarks on the Schur relations for a Frobenius algebra*, J. Math. Soc. Japan **5** (1953) 24-28.

Received November 14, 1969.

THE UNIVERSITY OF UTAH

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

C. R. HOBBY
University of Washington
Seattle, Washington 98105

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Norman Larrabee Alling, <i>Analytic and harmonic obstruction on nonorientable Klein surfaces</i>	1
Shimshon A. Amitsur, <i>Embeddings in matrix rings</i>	21
William Louis Armacost, <i>The Frobenius reciprocity theorem and essentially bounded induced representations</i>	31
Kenneth Paul Baclawski and Kenneth Kapp, <i>Topisms and induced non-associative systems</i>	45
George M. Bergman, <i>The index of a group in a semigroup</i>	55
Simeon M. Berman, <i>Excursions above high levels for stationary Gaussian processes</i>	63
Peter Southcott Bullen, <i>A criterion for n-convexity</i>	81
W. Homer Carlisle, III, <i>Residual finiteness of finitely generated commutative semigroups</i>	99
Roger Clement Crocker, <i>On the sum of a prime and of two powers of two</i>	103
David Eisenbud and Phillip Alan Griffith, <i>The structure of serial rings</i>	109
Timothy V. Fossum, <i>Characters and orthogonality in Frobenius algebras</i>	123
Hugh Gordon, <i>Rings of functions determined by zero-sets</i>	133
William Ray Hare, Jr. and John Willis Kenelly, <i>Characterizations of Radon partitions</i>	159
Philip Hartman, <i>On third order, nonlinear, singular boundary value problems</i>	165
David Michael Henry, <i>Conditions for countable bases in spaces of countable and point-countable type</i>	181
James R. Holub, <i>Hilbertian operators and reflexive tensor products</i>	185
Robert P. Kaufman, <i>Lacunary series and probability</i>	195
Erwin Kreyszig, <i>On Bergman operators for partial differential equations in two variables</i>	201
Chin-pi Lu, <i>Local rings with noetherian filtrations</i>	209
Louis Edward Narens, <i>A nonstandard proof of the Jordan curve theorem</i>	219
S. P. Philipp, Victor Lenard Shapiro and William Hall Sills, <i>The Abel summability of conjugate multiple Fourier-Stieltjes integrals</i>	231
Joseph Earl Valentine and Stanley G. Wayment, <i>Wilson angles in linear normed spaces</i>	239
Hoyt D. Warner, <i>Finite primes in simple algebras</i>	245
Horst Günter Zimmer, <i>An elementary proof of the Riemann hypothesis for an elliptic curve over a finite field</i>	267