# Pacific Journal of Mathematics

SPECTRAL THEORY OF MONOTONE HAMMERSTEIN OPERATORS

CHARLES VERNON COFFMAN

Vol. 36, No. 2

December 1971

# SPECTRAL THEORY OF MONOTONE HAMMERSTEIN OPERATORS

CHARLES V. COFFMAN

Consider the linear integral equation,

(1) 
$$y(t) = \mu \int_{a} K(t, s) p(s) y(s) ds$$
,

where K(s, t) is a real-valued symmetric positive definite kernel and p(s) is a positive function. Let L denote the inverse of the integral operator  $u \rightarrow \int_{a} K(\cdot, s) u(s) ds$ , and for a function y in the domain of L,  $y \neq 0$ , (all functions are assumed to be real valued) define the Rayleigh quotient J(y) for (1) by,

$$J(y) = \int_{\mathcal{Q}} y(t) \left[ Ly \right](t) dt / \int_{\mathcal{Q}} p(t) y^2(t) dt .$$

If  $y_1 \neq 0$  and  $y_1$  is in the domain of L and if

$$y_2 = \int_{a} K(\cdot, s) p(s) y_1(s) ds$$
,

then several applications of the Schwarz inequality show that,

$$J(y_2) \leq J(y_1)$$
.

with equality only if  $y_1$  is an eigenfunction of (1). On the basis of this fact, when the integral operator in (1) is compact, one can develop the complete spectral theory of (1).

In this paper it is shown that the approach indicated above for the study of (1) has a simple and natural extension for the study of the nonlinear integral equation.

(2) 
$$y(\mathbf{t}) = \mu \int_{\mathcal{Q}} K(t, s) f(s, y(s)) ds$$

where K(t, s) is as above and f(t, y) is an odd function of y,

$$f(t, y) = -f(t, -y),$$

and satisfies,

$$yf(t, y) > 0$$
,  $y$ 

and

$$f(t, y_2) \ge f(t, y_1)$$
,  $y_2 \ge y_1$ .

 $\neq$  0,

The problem of minimizing the Rayleigh quotient J(y) for (1) can be generalized to either of the dual variational problems,

$$\int_{\Omega} y(t)[Ly](t)dt = \min, \qquad \int_{\Omega} \int_{0}^{y(t)} f(t, \eta)d\eta dt = \text{const.},$$
$$\int_{\Omega} \int_{0}^{y(t)} f(t, \eta)d\eta dt = \max, \qquad \int_{\Omega} y(t)[Ly](t)dt = \text{const.}$$

By studying the first of these two variational problems, we shall prove here a variant of a theorem of Sobolev, [17], concerning (2); Sobolev's work treats the second of the above variational problems. For a discussion of Sobolev's theorem and related results see [8].

The Lyusternik-Schnirelman theory upon which the theorem of Sobolev is based has undergone considerable development in recent years, see [14], [16], with fruitful applications to the theory of nonlinear elliptic boundary value problems, see [1], [2]. The methods employed here differ substantially and in several respects from the standard methods of the Lyusternik-Schnirelman theory. We believe that for the problem at hand these methods are simpler and more natural, and therefore should be of interest. The central idea of using an iteration operator in the variational study of a nonlinear problem was suggested by the work in Moore and Nehari [11], and Nehari [12], [13]. The notion of "genus" which we use was introduced by Krasnosel'skii, and is treated in [8]: see also [4].

2. Let X be a real infinite dimensional Banach space,  $X^*$  its dual space, and let the value of a linear functional  $y \in X^*$  on an element  $x \in X$  be denoted (y, x). Let  $A: X \to X^*$  be a compact linear operator, take

(3) 
$$Y = \overline{AX},$$

and assume that A is symmetric,

(4) 
$$(Ax_1, x_2) = (Ax_2, x_1)$$
.  $x_1, x_2 \in X$ ,

and positive definite,

(5) 
$$(Ax, x) > 0$$
,  $x \in X \setminus \{0\}$ .

Note that if X is reflexive then (5) implies that  $Y = X^*$ .

Let  $\Phi: Y \to X$  be a continuous nonlinear mapping and assume that  $\Phi$  is the gradient (Fréchet derivative) of a real valued even functional  $\gamma(y)$  on Y satisfying  $\gamma(0) = 0$ . (The gradient of a functional on Y is actually an operator from Y to Y\*, however, X is canonically isometric to a subspace of Y\* so that  $\Phi$  determines in an obvious way a mapping of Y into Y\*, it is this mapping which, properly speaking, is the gradient of  $\gamma$ ; clearly this problem does not arise when x is reflexive.) The fact that  $\gamma$  is even implies that  $\Phi$  is odd,

(6) 
$$\Phi(y) = -\Phi(-y) ,$$

we assume also that  $\Phi$  is positive definite,

(7) 
$$(y, \varphi(y)) > 0$$
,  $y \in Y \setminus \{0\}$ ,

and monotone,

$$(y_2 - y_1, \ \Phi(y_2) - \Phi(y_1)) \ge 0, \qquad y_1, \ y_2 \in Y.$$

This last is equivalent to assuming that  $\gamma$  is convex, [19].

The result which we shall prove is the following.

THEOREM 1. Under the above assumptions concerning A and  $\Phi$ , the eigenvalue problem,

(9) 
$$y = \mu A \Phi(y) ,$$

has infinitely many eigenvectors satisfying,

(10) 
$$\gamma(y) = c$$

for every c > 0.

3. We begin the proof of the theorem stated above by establishing several results concerning the linear operator A. If we let  $\mathscr{R}$  denote the range of A then, by (3),  $\mathscr{R}$  is dense in Y, and by (5), A has an inverse  $L: \mathscr{R} \to X$ . From (4), (5) and the Schwarz inequality there follows

(9) 
$$(Ax_1, x_2)^2 \leq (Ax_1, x_1) (Ax_2, x_2)$$
.

and thus, since

$$||Ax||_{X^*} = \sup \{(Ax, x'): ||x'||_X \leq 1\},\$$

we have

$$|| Ax ||_{X^*} \leq || A || (Ax, x)$$
,

and for  $y \in \mathcal{R}$ , we have,

(10) 
$$|| y ||_{X^*}^2 \leq || A || (y, Ly) .$$

If we complete  $\mathscr{R}$  with respect to the inner product  $\langle y, y \rangle = (y, Ly)$  we obtain a Hilbert space  $H \subseteq Y$  with

(11) 
$$||i\eta||_{X^*}^2 \leq ||A|| ||\eta||_{H^*}^2$$
,

where *i* denotes the inclusion mapping  $H \subseteq X^*$ . It is clear that for  $\eta, \eta' \in H$  and  $i\eta = Ax \in \mathscr{R}$ ,

(12) 
$$\langle \eta', \eta \rangle = (i\eta', x)$$
.

Hence if we define  $B: X \rightarrow H$  by iBx = Ax, then

$$|| Bx ||_{\scriptscriptstyle H}^{\scriptscriptstyle 2} = \langle Bx, Bx \rangle = (Ax, x) \leq || A || \, || \, x \, ||_{\scriptscriptstyle X}^{\scriptscriptstyle 2}$$
 ,

so that B is a continuous linear mapping of X into H. Moreover, upon taking  $\eta = Bx$  in (12) we get

$$\langle \eta', Bx \rangle = (i\eta', x)$$
,

from which it follows that

(13)  $B^* = i$ .

We next show that the compactness of A imples the compactness of both B and i.

LEMMA 2. The mapping  $B: X \rightarrow H$  is compact.

*Proof.* Let  $\{x_n\}$  be a bounded sequence in X. There is no loss of generality in assuming that  $\{Ax_n\}$  converges strongly in  $X^*$ , since A is compact, but then,

$$|| B(x_n - x_m) ||^2 = \langle B(x_n - x_m), B(x_n - x_m) \rangle,$$
  
=  $(A(x_n - x_m), (x_n - x_m)) \rightarrow 0,$ 

as  $n, m \to \infty$ , and thus B is compact.

By standard results, the compactness of B implies the compactness of  $i = B^*$ , thus we have the following.

LEMMA 3. The imbedding  $i: H \subseteq X^*$  is compact.

4. We turn our attention now to the nonlinear operator  $\Phi$ , and observe that, for  $y_1, y_2 \in Y$  we have

$$\gamma(y_2) - \gamma(y_1) = \int_0^1 (y_2 - y_1, \Phi(y_1 + t(y_2 - y_1))) dt$$

(see [18]), but by (8),

$$(y_2-y_1, arPhi(y_1+t(y_2-y_1))) \geqq (y_2-y_1, arPhi(y_1))$$
 ,

so that

(14) 
$$\gamma(y_2) - \gamma(y_1) \ge (y_2 - y_1, \ \Phi(y_1)) \ .$$

Next we define  $\varphi: H \to R, \Psi: H \to H$ , by

(15) 
$$\varphi = \gamma \cdot i$$
,  $\Psi = B \cdot \Phi \cdot i;$ 

:306

then the continuity of  $\gamma$  and  $\Phi$  and the compactness of *i* imply the following result.

LEMMA 4. The functional  $\varphi$  is weakly continuous and  $\Psi$  is completely continuous.

From (7), (15), (14) and (13) we get

(16) 
$$\langle \eta, \Psi(\eta) \rangle > 0$$
,  $\eta \in H \setminus \{0\}$ ,

and

(17) 
$$\langle \eta' - \eta, \Psi(\eta) \rangle \leq \varphi(\eta') - \varphi(\eta) , \qquad \eta', \eta \in H.$$

LEMMA 5. Let c > 0, then for  $\eta \in H \setminus \{0\}$ , there exists a unique a > 0 such that  $\varphi(\alpha \eta) = c$ . Moreover,  $\alpha$  is a continuous function of  $\eta$  on  $H \setminus \{0\}$ .

*Proof.* By hypothesis,  $\gamma(0) = 0$ , thus, from (15) it follows that  $\varphi(0) = 0$ . By taking  $\eta' = \alpha \eta$  in (17) we get

(18) 
$$\varphi(\alpha\eta) \ge \varphi(\eta) + (\alpha - 1) < \eta, \Psi(\eta) > ,$$

hence (16) implies that  $\varphi(\alpha \eta) \to \infty$  as  $\alpha \to \infty$ , for  $\eta \neq 0$ . The existence of  $\alpha$  then follows from the continuity of  $\varphi$ ; uniqueness follows from (16) and (18). The continuity follows from the continuity of  $\varphi$  and the uniqueness of  $\alpha$ .

Let c > 0 be fixed, put

(19) 
$$\Sigma = \Sigma_c = \{ \eta \in H \colon \varphi(\eta) = c \},\$$

and define  $\sigma: H \setminus \{0\} \to \Sigma_c$  by

(20) 
$$\sigma(\eta) = \alpha \Psi(\eta) ,$$

where  $\alpha > 0$  is chosen so that  $\sigma(\eta) \in \Sigma$ .

LEMMA 6. The mapping  $\sigma: H \setminus \{0\} \to \Sigma$  is odd and strongly continuous (in the sense of [18], i.e.  $\sigma$  maps weakly convergent sequences in  $H \setminus \{0\}$  to strongly convergent sequences). If  $\eta \in \Sigma$ , then

$$|| \sigma(\eta) ||_{H} \leq || \eta ||_{H}.$$

with equality only if  $\eta$  is an eigenvector of

(22) 
$$\eta = \mu \Psi(\eta) \; .$$

*Proof.* The oddness and strong continuity of  $\sigma$  follow respectively from (6), and the definition of  $\Psi$ , and Lemmas 4 and 5. If  $\eta' = \sigma(\eta)$ ,

then since  $\varphi(\eta') = \varphi(\eta)$ , (17) implies

$$\langle \eta' - \eta, \Psi(\eta) \rangle \leq 0$$
,

but by (20), this is equivalent to

$$\langle \eta' - \eta, \eta' 
angle \leq 0$$
 ,

or

$$\langle \eta', \eta' 
angle \leq \langle \eta, \eta' 
angle \leq \langle \eta, \eta 
angle^{_{1/2}} \langle \eta', \eta' 
angle^{_{1/2}}$$
 ,

which implies (21). From the way in which the Schwarz inequality was used, it is clear that equality can hold in (21) only if  $\eta'$  and  $\eta$  are proportional, i.e. only if  $\eta$  is an eigenvector of (22).

5. Let S denote the class of closed subsets of  $H\setminus\{0\}$  which are symmetric through the origin. For a non-empty set  $F \in S$ , the genus of F,  $\rho(F)$ , is the supremum of the set of nonnegative integers n such that every odd continuous map of F into  $\mathbb{R}^{n-1}$  has a zero in F; here we understand  $\mathbb{R}^0 = \{0\}$ . The genus of the empty set is zero.

Below, the letter 'F', with or without subscript, will always denote a set in the class S. The genus has the following properties.

1. If there exists an odd continuous map  $h: F_1 \to F_2$ , then  $\rho(F_1) \leq \rho(F_2)$ , in particular, if  $F_1 \subseteq F_2$ , then  $\rho(F_1) \leq \rho(F_2)$ .

2.  $\rho(F_1 \cup F_2) \leq \rho(F_1) + \rho(F_2)$ .

3. If F is compact then  $\rho(F) < \infty$  and F has a neighborhood U such that  $\bar{U} \in S$  and  $\rho(\bar{U}) = \rho(F)$ .

4. If  $\{F_n\}$  is a decreasing sequence of compact sets and if  $F = \bigcap_{n=1}^{\infty} F_n$  then  $F \in S$  and

$$\rho(F) = \lim_{n \to \infty} \rho(F_n)$$
.

5. If there exists an odd homeomorphism of F onto the *n*-sphere then  $\rho(F) = n + 1$ .

For a proof of the above assertions see [4].

6. In this section we complete the proof of Theorem 1. Let the number c > 0 be fixed, and let  $\Sigma$  be defined by (19). We shall call a set F admissible if F is compact,  $F \in S$ , and  $F \subseteq \Sigma$ . The class of admissible sets will be denoted by  $\mathscr{F}$ , and we shall take, for  $n = 1, 2, \cdots$ ,

$$\mathscr{F}_n = \{F \in \mathscr{F} : \rho(F) \ge n\}.$$

LEMMA 7. For any positive integer n, the class  $\mathscr{F}_n$  is nonempty.

*Proof.* Let M be a subspace of H of dimension n, then it follows from Lemma 5 that the mapping  $\eta \to ||\eta||_{H}^{-1}\eta$  on  $\Sigma \cap M$  is a homeomorphism onto the unit sphere in M. It thus follows from property 5 of the genus that  $\rho(\Sigma \cap M) = n$ , and thus  $\Sigma \cap M \in \mathscr{F}_{n}$ .

We now define the numbers  $\lambda_n = \lambda_n(c)$  by,

(23)  
$$\lambda_n = \frac{1}{2c} \inf_{F \in \mathscr{F}_n} \max_{\eta \in F} ||\eta||_H^2,$$
$$= \frac{1}{8c} \inf_{F \in \mathscr{F}_n} [\operatorname{diam} F]^2,$$

for  $n \ge 1$ , and for convenience we define  $\lambda_0 = 0$ . Since  $\Sigma$  is closed and does not contain zero, and because of the definition (23), it is clear that sequence  $\lambda_1, \lambda_2, \cdots$ , is a nondecreasing sequence of positive numbers. We say that  $\lambda_n$  has *multiplicity* m if,

$$\lambda_{k-1} < \lambda_k = \lambda_{k+1} = \cdots = \lambda_{k+m-1} < \lambda_{k+m}$$
 ,

for some  $k \ge 1$ , where  $k \le n \le k + m - 1$ . We also make the definitions, for  $\lambda \ge 0$ ,

$$egin{aligned} \Sigma\left(\lambda
ight) &=\left\{\eta\in\Sigma\colon||\:\eta\:||_{H}^{2}\leq2c\lambda
ight\}\,,\ \Sigma^{\scriptscriptstyle 1}(\lambda) &=\sigma(\Sigma(\lambda))\;, \end{aligned}$$

and note that because of the weak continuity of  $\varphi$  (Lemma 4),  $\Sigma(\lambda)$  is weakly closed and hence since it is bounded it is weakly compact. It follows then from Lemma 6 that  $\Sigma^{1}(\lambda)$  is compact in the strong topology and,

(24) 
$$\Sigma^{\scriptscriptstyle 1}(\lambda) \subset \Sigma(\lambda)$$
 ,

clearly  $\Sigma^{1}(\lambda)$  is symmetric and thus

(25) 
$$\Sigma^{1}(\lambda) \in \mathscr{F}$$
,  $(\operatorname{diam} \Sigma^{1}(\lambda))^{2} \leq 8\lambda c$ .

On the other hand, by property 1 of the genus, since  $\sigma$  is odd, and because of (24),

$$\rho(\Sigma^{1}(\lambda)) = \rho(\Sigma(\lambda))$$

Since  $\Sigma^{1}(\lambda)$  is compact, it follows from property 3 that  $\rho(\Sigma(\lambda)) < \infty$  for any  $\lambda > 0$ .

If we introduce the "spectral function"

$$au(\lambda)=
ho(arsigma^{_1}(\lambda))$$
 ,

then  $\tau(\lambda)$  is a monotone integer valued function of  $\lambda$  for  $\lambda \ge 0$ ,

$$au(\lambda)=0$$
 ,  $0\leq\lambda<\lambda_{1}$  ,

and since,

$$\Sigma^{1}(\lambda) = \bigcap_{\mu > \lambda} \Sigma^{1}(\mu)$$
 ,

it follows from property 4 that  $\tau(\lambda)$  is right continuous. Finally from (23) and (24) and the right continuity of  $\tau$  it follows that,

(26) 
$$\tau(\lambda) = n$$
,  $\lambda_n \leq \lambda < \lambda_{n+1}$ .

Thus the discontinuities of  $\tau(\lambda)$  occur only at the numbers  $\lambda_n$  given by (23) and

(27) 
$$\tau(\lambda_n) - \tau(\lambda_n - 0) =$$
multiplicity of  $\lambda_n$ .

It follows from (26) that no finite  $\lambda$ -interval can contain more than finitely many of the numbers  $\lambda_n$ , thus

(28) 
$$\lim_{n\to\infty}\lambda_n = \infty$$

Now let  $E_n$  denote the set of solutions  $\eta$  in  $\Sigma = \Sigma_c$  of (22) such that  $||\eta||_H^2 = 2c\lambda_n$ , (clearly  $E_n$  is admissible and  $E_n \subseteq \Sigma^1(\lambda_n)$ ), and choose, by property 3, a neighborhood U of  $E_n$  such that  $\overline{U} \in S$  and  $\rho(\overline{U}) = \rho(E_n)$ . It then follows from Lemma 6 that

$$[\text{diam } (\sigma(\Sigma^{1}(\lambda_{n}) \setminus U))]^{2} < 8c\lambda_{n},$$

so that from the definition of  $\tau$  and (27),

$$egin{aligned} &
ho(\sigma(arsigma^{1}(\lambda_{n})igwed U)) &\leq au(\gamma_{n}\,-\,0) \ &\leq au(\lambda_{n})\,-\, ext{mult.} ext{ of } \lambda_{n} \;. \end{aligned}$$

Using this last inequality, property 1 of the genus, and the definition of  $\tau$ , we obtain,

$$\rho(\Sigma^{1}(\lambda_{n}) \setminus U) \leq \rho(\Sigma^{1}(\lambda_{n})) - \text{mult. of } \lambda_{n}$$
.

It then follows from property 2 of the genus that

$$ho(E_n) = 
ho(\overline{U}) \ge ext{mult. of } \lambda_n$$
 ,

in particular,  $E_n$  is not empty. We have thus proved the following.

(\*) The eigenvalue problem (22) has infinitely many eigenvectors  $\eta$  satisfying

(29) 
$$\varphi(\eta) = c$$
.

The set  $E_n(c)$  of eigenvectors  $\eta$  of (22) satisfying (29) and

 $||\gamma||_{H}^{2}=2c\lambda_{n}$ ,

where  $\lambda_n$  is given by (23), is a set of genus  $\geq multiplicity$  of  $\lambda_n$ .

It is clear from (15) and the definition of *B* that if  $\eta$  is an eigenvector of (22) satisfying (29) then  $y = i\eta$  is an eigenvector of (9) and satisfies (10). Thus (\*) implies Theorem 1. Moreover, the following is valid.

$$(**)$$
 The set of eigenvectors y of (9) which satisfy (10) and

$$(30) (y, Ly) = 2c\lambda_n(c) ,$$

is a set of genus  $\geq$  multiplicity of  $\lambda_n$ . The numbers  $\lambda_n(c)$  can be determined as follows

(31) 
$$\lambda_n(c) = \frac{1}{2c} \inf_{G \in \mathscr{G}_n} \sup_{y \in G} (y, Ly) ,$$

where  $\mathcal{G}_n$  is the class of symmetric subsets G of  $\mathscr{R}\setminus\{0\}$  which are closed in Y, have genus  $\geq n$  and satisfy

$$G \subseteq \{y \in Y: \gamma(y) = c\}$$
 .

Here the genus is to be understood to be relative to the Y-topology.

*Proof.* First we observe that the H and Y topologies coincide on compact subsets of H. Thus the genus of  $E_n$  relative to the H topology and the genus of  $i(E_n)$  relative to the Y topology are the same. Since the set of eigenvectors of (9) satisfying (10) and (30) is just  $i(E_n)$ , the first assertion above is proved. To prove the second assertion, let  $G \in \mathcal{G}_n$  and let  $\lambda = 1/2c \sup_{y \in G} (y, Ly) < \infty$ , then G = i(F), where F is a closed symmetric subset of H and  $F \subseteq \Sigma(\lambda)$ . From (15) and (20), and the continuity of  $\alpha$  in (20) it follows that  $\sigma(F)$  is the image, under an odd completely continuous transformation, of G. Thus  $\sigma(F)$  is compact and has genus  $\geq n$ . However, we clearly have  $[\operatorname{diam} (\sigma(F))]^2 \leq 8c\lambda$  and thus it follows from (23) that

$$\lambda_n \leq \frac{1}{2c} \inf_{G \in \mathscr{G}_n} \sup_{y \in G} (y, Ly) .$$

Making use of the observation at the beginning of the proof we conclude that  $i(\sigma(\Sigma^1(\lambda_n))) \in \mathcal{G}_n$ , and this, together with the above inequality, implies (31).

REMARK. The existence of an infinity of eigenfunctions y of the problem (9) satisfying

$$(y, Ly) = c$$
,

### CHARLES V. COFFMAN

for an arbitrary c > 0, follows by applying Theorem 4.3 of [8] to the operator  $\Psi$ , provided that  $\Psi$  is uniformly differentiable on bounded sets; it is not required that  $\Phi$  satisfy (8). The theorem of sobolev, [17], quoted earlier, follows from this result. For a discussion of the multiplicity of solutions in this case, when the associated critical values are repeated, see [3].

7. It is of interest to show that the above results do include the complete spectral theory of (9) in the linear case. Thus suppose that  $\Phi$  is linear and observe that in this case,

$$\gamma(y)=\int_{0}^{1}(y,\, arPhi(ty))dt=rac{1}{2}(y,\, arPhi(y))\;.$$

Thus, for  $y, h \in Y$ ,

$$egin{aligned} &\gamma(y+h)\ &=rac{1}{2}\left(y,\,arPsi(y)
ight)+rac{1}{2}\left(h,\,arPsi(y)
ight)+rac{1}{2}\left(y,\,arPsi(h)
ight)+rac{1}{2}\left(h,\,arPsi(h)
ight)\,, \end{aligned}$$

while on the other hand the definition of the gradient gives

$$egin{aligned} &\gamma(y\,+\,h) = \,\gamma(y)\,+\,(h,\,arPhi(y)\,+\,o(||\,h\,||)\,, \ &= rac{1}{2}\,(y,\,arPhi(y))\,+\,(h,\,arPhi(y))\,+\,o(||\,h\,||)\,\,, \end{aligned}$$

as  $||h|| \rightarrow 0$ . Comparison of these two formulas shows that  $\Phi$  is symmetric,

$$(y_1, \Phi(y_2)) = (y_2, \Phi(y_1)) , \qquad y_1, y_2 \in Y$$
 .

Using this together with (7), (15) and Lemma 2 we conclude that  $\Psi$  is a compact self-adjoint positive definite operator and

$$arphi(\eta)=rac{1}{2}ig\langle\eta,arPsi_\eta
angle$$
 ;

see the discussion of the operator  $\chi_1$  in §8 below. Let  $0 < \mu_1 \leq \mu_2 \cdots$ denote the characteristic values of  $\Psi$  and let  $M_1 \subseteq M_2 \subseteq \cdots$  be the corresponding sequence of invariant subspaces for  $\Psi$ , i.e.,  $M_n$  is spanned by those eigenvectors of  $\Psi$  corresponding to the first *n* characteristic values. Suppose that  $\mu_n < \mu_{n+1}$  and for some c > 0 let  $\Sigma_c(\lambda)$ be defined as in §6.

If  $\mu_n \leq \lambda < \mu_{n+1}$  then for  $\eta \in \Sigma_c(\lambda)$ ,

$$egin{aligned} &|| \eta \, ||_{H}^{2} \leq 2 \lambda c = \lambda ig\langle \eta, \, \varPsi(\eta) ig
angle \ &< \mu_{n+1} ig\langle \eta, \, \varPsi(\eta) ig
angle \, . \end{aligned}$$

Thus if  $P_n$  denotes the orthogonal projection of H onto  $M_n$  then

 $P_n\eta \neq 0$  for  $\eta \in \Sigma_c(\lambda)$ , since

$$|| \, \eta \, ||_{_{H}}^{_{2}} \geq \mu_{n+1} ig\langle \eta, \, \varPsi(\eta) ig
angle$$

for  $\eta$  in the kernel of  $P_n$ . It follows that the map

$$\eta \longrightarrow (2c)^{1/2} \langle P_n \eta, \Psi(\eta) \rangle^{-1/2} P_n \eta$$

maps  $\Sigma_{\mathfrak{c}}(\lambda)$  onto  $\Sigma_{\mathfrak{c}} \cap M_n$ . Since this mapping is odd and continuous on  $\Sigma_{\mathfrak{c}}(\lambda)$  it follows that  $n = \rho(\Sigma_{\mathfrak{c}} \cap M_n) \ge \rho(\Sigma_{\mathfrak{c}}(\lambda))$ . On the other hand, for  $\eta \in M_n$ ,

$$|| \eta ||_{H}^{2} \leq \mu_{n} \langle \eta, \Psi(\eta) \rangle$$

and thus

$$\Sigma_{c} \cap M \subseteq \Sigma_{c}(\mu_{n}) \subseteq \Sigma_{c}(\lambda),$$

from which we conclude that

$$ho({\Sigma}_{c}(\lambda)) = n$$
,  $\mu_{n} \leq \lambda < \mu_{n+1}$ .

Combined with the results of § 6, this yields the following.

(\*\*\*) For the linear problem and for any c > 0, the number  $\lambda_n(c)$  is just the  $n^{th}$  characteristic value, in increasing order, of the operator  $\Psi$ .

We require also the following result concerning the linear case.

LEMMA 8. Let F be a compact symmetric subset of  $H \setminus \{0\}$ , with  $\rho(F) \geq n$ , then

$$\max_{\eta \in F} rac{||\eta||_{H}^{2}}{\langle \eta, \Psi \eta \rangle} \geq \mu_{n} \; ,$$

where  $\mu_n$  is the  $n^{th}$  characteristic value, in increasing order, of the linear operator  $\Psi$ .

*Proof.* Let F' be the image of F under  $\eta \to (2c)^{1/2} \langle \eta, \Psi \eta \rangle^{-1/2} \eta$ , where c > 0. Then  $F' \subseteq \Sigma_c$  and  $\rho(F') \ge n$ , thus by (\*\*\*),

$$\max_{\eta \, \in \, F} rac{||\, \eta \, ||_{H}^{2}}{\langle \eta, \, \varPsi \, \eta 
angle} = rac{1}{2c} \max_{\eta \, \in \, F'} \, ||\, \eta \, ||_{H}^{2} \geqq \mu_{n} \; .$$

8. In this section we shall derive a "principle of linearization" for the problem (9); cf. [Ch. VI,  $\S$  2, 8]. What we shall prove is the following.

(†) Let  $\Phi$  have a Fréchet derivative  $\chi$  at y = 0, and suppose that the continuous linear operator  $\chi: Y \to X$  is symmetric,

$$(32) (y_1, \chi y_2) = (y_2, \chi y_1), y_1, y_2 \in Y,$$

and positive defininite,

$$(33) (y,\chi y) > 0, y \in Y \setminus \{0\}.$$

Let  $\kappa_1, \kappa_2, \kappa_3, \cdots$ , be the characteristic values, arranged in increasing order, of the linear operator  $A\chi$ , then the numbers  $\lambda_n(c)$ , given by (31) satisfy

(34) 
$$\lim_{c\to 0} \lambda_n(c) = \kappa_n .$$

Moreover, given  $\varepsilon > 0$ , there exists  $c_1(\varepsilon) > 0$  such that for  $0 < c < c_1$ , the characteristic numbers  $\mu$  corresponding to those eigenvectors  $\eta$ of (9) which satisfy (10) and (30) all lie in the interval

$$(\kappa_n-\varepsilon, \kappa_n+\varepsilon)$$
.

Finally, if  $\tilde{N}_n$  is the eigenspace of A $\chi$  corresponding to the characteristic number  $\kappa_n$ , and if  $\tilde{E}_n(c)$  is the set of eigenvectors of (9) satisfying (10) and (30), then

$$\lim_{c\to 0} \max_{y \in \widetilde{E}_n(c)} \frac{\operatorname{dist}(y, N_n)}{||y||} = 0.$$

REMARK. The above assertion implies that, in the terminology of [8], each characteristic value of  $A\chi$  is a bifurcation point for the problem (9), compare Theorem 2.2, [8, p. 332].

Proof of (†). From the definition of the Fréchet derivative

$$\Phi(y) = \chi y + \omega(y), \qquad \qquad y \in Y,$$

where

$$|| \omega(y) || = o(|| y ||)$$
, as  $y \longrightarrow 0$ .

Thus, for  $y \in Y$ ,

$$egin{aligned} &\gamma(y) = \int_{_0}^{_1} (y, \, \varPhi(ty)) dt, \ &= rac{1}{2} \left( y, \, \chi y 
ight) + o(||\, y\, ||^2) \;, \qquad & ext{as} \; \; y \longrightarrow 0 \;, \end{aligned}$$

and consequently, from (15), (12) and (13), for  $\eta \in H$ ,

$$arphi(\eta) = rac{1}{2} \langle \eta, \chi_1 \eta 
angle + o(|| \, i\eta \, ||^2) \; , \qquad ext{ as } \eta \longrightarrow 0 \; ,$$

where

$$\chi_1 = B\chi i$$
.

If we make the natural identification of X with a subspace of  $Y^*$ , then (32) implies that  $\chi^* \supseteq \chi$ . Since the range of *i* is contained in the domain of  $\chi$  it follows, using (13), that the operator  $\chi_1 = B\chi i$  is self-adjoint. Since

$$\langle \eta, \chi_{\scriptscriptstyle 1} \eta 
angle = (i\eta, \chi i \eta)$$
 ,

it follows from (33) that  $\chi_1$  is positive definite, and from Lemma 2 it follows that  $\chi_1$  is compact. Clearly the characteristic values  $\{\kappa_n\}$  of  $A\chi$  are also characteristic values of  $\chi_1$ , thus,  $\chi_1$  has an invariant subspace M of dimension n such that

$$(36) || \eta ||_{H}^{2} \leq \kappa_{n} \langle \eta, \chi_{1} \eta \rangle, \eta \in M.$$

Let  $\varepsilon > 0$  be given and, on the strength of (35), choose  $c_1 > 0$  such that

(37) 
$$\left| \varphi(\eta) - \frac{1}{2} \langle \eta, \chi_1 \eta \rangle \right| \leq \frac{1}{2} \kappa_n^{-1} \varepsilon || \eta ||_H^2$$

for

(38) 
$$||\eta||_{H}^{2} \leq 2 (1 - \varepsilon)^{-1} \kappa_{n} c_{1}$$
.

Form (36) and (37) we then have

$$arphi(\eta) \geqq rac{1}{2} \kappa_n^{-1} (1-arepsilon) || \ \eta \ ||_H^2 \ , \qquad \eta \in M, \ || \ \eta \ ||_H^2 \leqq 2 (1-arepsilon)^{-1} \kappa_n c' \ ,$$

from which, it follows, by the use of (18), that for 0 < c < c',

$$rac{1}{8} \left[ ext{diam} \left( M \cap \varSigma(c) 
ight) 
ight]^2 \leq \kappa_n (1 - arepsilon)^{-1} c \; .$$

Hence, from (23) and since  $\rho(M \cap \Sigma(c)) = n$ ,

$$\lambda_n(c) \leq \kappa_n(1-\varepsilon)^{-1}$$
.

Letting  $\varepsilon$  tend to zero, we obtain finally,

(39) 
$$\lim_{a\to 0} \sup \lambda_n(c) \leq \kappa_n.$$

Suppose now that

(40) 
$$\lim_{c\to 0} \inf \lambda_n(c) < \kappa < \kappa_n ,$$

and then choose  $\varepsilon > 0$  so that

$$(41) (1-\varepsilon)^{-1} \kappa < \kappa_n .$$

With this choice of  $\varepsilon$ , let c' > 0 be chosen as above so that (37) holds when  $\eta$  satisfies (38). In view of the above supposition, there exists a number c and a set F such that 0 < c < c', F is admissible for the given choice of c,  $\rho(F) \ge n$  and

(42) 
$$\frac{1}{2c} \max_{\eta \in F} ||\eta||_{H}^{2} < \kappa .$$

From (37), (42) and the fact that  $F \subseteq \Sigma_c$ , there follows, for  $\eta \in F$ ,

$$rac{1}{2}\langle\eta,\chi_{1}\eta
angle\geq c(1-arepsilon)\;.$$

Hence, by (42) and (41), (since  $\kappa < \kappa_n$ ),

$$egin{aligned} \max_{\eta \, \epsilon \, F} \, rac{|| \, \eta \, ||_{H}^{2}}{\left< \eta, \, \chi_{1} \eta \right>} & \leq rac{1}{2c(1-arepsilon)} \, \max_{\eta \, \epsilon \, F} \, || \, \eta \, ||_{H}^{2} \; , \ & \leq rac{1}{1-arepsilon} \, \kappa \; , \ & < \kappa_{n} \; . \end{aligned}$$

In view of Lemma 8 this is impossible, we conclude therefore that (40) cannot hold, and thus from (39) we have (34).

Since for an eigenvector  $\eta$  of (22) the corresponding characteristic value  $\mu$  is given by

$$\mu = rac{||\,\eta\,||_{_H}^2}{ig\langle \eta,\, arpsilon(\eta) ig
angle} \ ,$$

we have, for  $\eta \in \Sigma_{e}$ 

$$\mu = rac{arphi(\eta)}{rac{1}{2} ig\langle \eta, arphi(\eta) ig
angle} \cdot rac{1}{2c} \, || \, \eta \, ||^{\scriptscriptstyle 2}_{\scriptscriptstyle H} \; ,$$

thus if  $\eta \in E_n(c)$ ,

(43) 
$$\mu = \frac{\varphi(\eta)\lambda_n(c)}{\frac{1}{2}\langle \eta, \Psi(\eta) \rangle} .$$

By definition

$$[\operatorname{diam} E_n(c)]^2 = 8c\lambda_n(c)$$
,

so that from (35) and (34), for  $\eta \in E_n(c)$ , the characteristic value  $\mu$ 

which is given by (43), satisfies

$$\mu = \mu(\eta) = \kappa_n(1 + o(c))$$
, as  $c \longrightarrow 0$ ,

from which follows the second assertion of  $(\dagger)$ ; notice that

$$\widetilde{E}_n(c) = i(E_n(c))$$
.

The final assertion of  $(\dagger)$  follows from the second assertion of  $(\dagger)$  and a result from [8, pp. 194–195].

9. We conclude with several examples to which the above results are applicable. First, for the verification of condition (5), the following result will be useful.

LEMMA 9. The compact symmetric operator A:  $X \rightarrow X^*$  satisfies condition (5) if and only if  $Y = \overline{AX}$  is total for X and

$$(44) (Ax, x) \ge 0, x \in X.$$

*Proof.* Condition (5) clearly implies (44), and if  $x \in X$  and (y, x) = 0 for every  $y \in Y$ , then in particular (Ax, x) = 0, and hence x = 0, so that Y is indeed total for X. Conversely, suppose that (44) holds and that Y is total for X. Then if  $x \in X$ ,  $x \neq 0$ , there exists  $y \in Y$  such that  $(y, x) \neq 0$ . Since AX is dense in Y it follows that there exists  $x' \in X$  such that  $(Ax', x) \neq 0$ . But from the symmetry of A, (44) and the Schwarz inequality there follows,

$$0 < (Ax', x)^2 \leq (Ax', x') (Ax, x)$$
,

and thus we conclude that (5) holds.

Now let  $\Omega$  be a bounded region in Euclidean *n*-space, let K(t, s) be a symmetric kernel defined for  $(t, s) \in \Omega \times \Omega$ , and let f(t, y), defined for  $(t, y) \in \Omega \times R$  satisfy the Carathéodory conditions, [8]. Assume moreover that f(t, y) satisfies the conditions set down in §1, namely

(45) 
$$f(t, y) = -f(t, -y) , \qquad t \in \Omega, y \in R ,$$

(46) 
$$yf(t, y) > 0$$
,  $t \in \Omega, y \in R, y \neq 0$ ,

and

(47) 
$$f(t, y_2) \ge f(t, y_1)$$
  $t \in \Omega, y_1, y_2 \in R, y_2 \ge y_1$ .

With X and Y yet to be specified, we take A to be the integral operator

$$Au = \int_{a} K(\cdot, s) u(s) ds$$
,

and  $\Phi$  to be the Nemytsky operator

$$arPsi(y)=f(\ ullet$$
 ,  $y(\ ullet$  )) ,

finally we take

(48) 
$$\gamma(y) = \int_{\mathscr{Q}} \int_{0}^{y(t)} f(t, u) \, du \, dt \, .$$

We consider first the case where K(t, s) is continuous on  $\overline{\Omega} \times \overline{\Omega}$ and f(t, y) is continuous on  $\overline{\Omega} \times R$ . In this case we take  $X = L^1(\Omega)$ , and Y is then determined and will be a subspace of

$$C(\overline{\varOmega}) \subseteq L^{\infty}(\overline{\varOmega}) = (L^1(\varOmega)^*$$
.

The complete continuity of A:  $L^{1}(\Omega) \rightarrow C(\overline{\Omega})$  and the continuity of  $\Phi: C(\overline{\Omega}) \to L^1(\Omega)$  are easily verified; weaker conditions on f(t, y) suffice for the latter continuity. The symmetry of A follows from the sym-The operator  $\Phi$ , which can be regarded as an metry of K(t, s). operator from  $C(\Omega)$  into  $(C(\overline{\Omega}))^*$ , since the latter contains a subspace naturally isomorphic to  $L^{1}(\Omega)$ , is the gradient of  $\gamma$ , given by (48). The properties (6), (7) and (8) for  $\Phi$  follow from (45), (46) and (47). If K(t, s) is nonnegative definite in the ordinary sense, *i.e.* if  $A \mid L^2(\Omega)$ is nonnegative definite, then A satisfies (44), thus the applicability of Theorem 1 hinges on the totality of Y for  $L^{\iota}(\Omega)$ . This depends on more special properties of the kernel K(t, s). However, if  $\Omega$  is a bounded interval and if K(t, s) is the Green's function for a regular self-adjoint two-point boundary value problem on  $\Omega$ , then, provided K(t, s) in positive definite, Y will contain all continuous functions which vanish identically near the endpoints of  $\Omega$  and consequently will be total for  $L^{1}(\Omega)$ . Thus Theorem 1 implies the following.

THEOREM 2. Let

be a formally self-adjoint regular differential operator of order 2m on [a, b], let

(50) 
$$M_k(y) = \sum_{j=0}^{2m-1} (\alpha_{kj} y^{(j)}(a) + \beta_{kj} y^{(j)}(b)) = 0$$
,  $k = 1, \dots, 2m$ ,

be self-adjoint boundary conditions for  $\checkmark$ , and suppose that if  $y \in C^{2m}$ ([a, b]),  $y \neq 0$  in [a, b], and y satisfies (50) then

$$\int_a^b y(t) [\swarrow y](t) dt > 0$$
 .

Let f(t, y) by continuous on  $[a, b] \times R$  and satisfy (45), (46), and (47). Then if c > 0, the problem

(51) 
$$\qquad \qquad \swarrow(y) = \mu f(t, y) , \qquad M_k(y) = 0, \ k = 1, \cdots, 2m ,$$

has infinitely many eigenfunctions y satisfying

$$\gamma(y) = \int_a^b \int_0^{y(t)} f(t, u) du dt = c .$$

The principle of linearization derived in §8 applies to the problem (51), provided there exists a positive function q(t) on [a, b] such that

$$\lim_{y \to 0 \ y \neq 0} y^{-1} f(t, y) = q(t)$$
, uniformly with respect to  $t$ .

Substantially weaker conditions actually suffice for the Fréchet differentiability at zero of the Nemytsky operator from C to  $L^1$ .

We next consider the case where the kernel K(t, s) is singular. We are primarily interested in the particular case where K(t, s) is the Green's function associated with an elliptic boundary value problem, thus, of the various conditions on K implying complete continuity of A we shall consider only the one which is satisfied by such a Green's function, namely,

(52) 
$$\sup_{t\in a}\int_{a}|K(t,s)|^{a} ds < \infty ,$$

for some a > 1. Concerning f(t, y) we then assume

(53) 
$$|f(t, y)| \leq c |y|^b + d$$
,

for some b such that

(54) 
$$1 < b < 2a - 1$$
,

and we take  $X = L^{q}(\Omega)$ , where,

(55) 
$$q > 2a/(2a-1)$$
.

If 1/p + 1/q = 1, we have by [7, Th. 95.6, p. 658], that  $A: L^q(\Omega) \to L^p(\Omega)$ is completely continuous, and by [18, Th. 19.1, p. 154] that  $\Phi: L^p(\Omega) \to L^q(\Omega)$  is continuous. Also,  $\Phi$  is the gradient of the functional  $\gamma$  on  $L^p(\Omega)$ ; see [18, § 6.3, p. 63]. Assume that K(t, s) is positive definite in the ordinary sense, then since, (because of the reflexivity of  $L^p(\Omega)$ ), no proper subspace of  $L^p(\Omega)$  is total for  $L^q(\Omega)$ , the applicability of our main result hinges in this case on the density of the range of A in  $L^p(\Omega)$ . We note however that in any case, in the presence of conditions (52), (53), (54) and (55) and when 1/p + 1/q = 1, any  $L_p$ -eigen-

function of (2) actually belongs to  $L^{\infty}(\Omega)$ ; see the proof of Theorem 3, [5].

In order for the Nemytsky operator from  $L^{p}(\Omega)$  to

$$L^{q}(\Omega)(p > 2, \frac{1}{p} + \frac{1}{q} = 1)$$

to be Fréchet differentiable at 0 it suffices that

$$f(t, y) = y(P(t) + q(t, y))$$

where  $P \in L^{r}(\Omega)$ ,  $r = p \setminus (p-2)$ , q(t, y) satisfies the Carathéodory conditions,  $q(t, 0) \equiv 0$ , and

(56) 
$$|q(t, y)| \leq m_1 |y|^{\delta} + m_2$$
,

where  $m_1, m_2 > 0$ ,  $\delta = (p-2)$ . If p(t) is positive almost everywhere on  $\Omega$ , then the principle of linearization is applicable in this case. In the presence of (52), (53), (54), the small  $L^p$ -solutions of (2), (p > 2a), can be shown to be shall in the  $L^{\infty}$  norm also, and thus the growth condition (56) on g(t, y) can be dropped when these conditions hold.

We will not state a general theorem concerning the equation (2) with a singular kernel but rather we state the following result, which is a principal application of such a theorem.

THEOREM 3. Let  $\Omega$  be a bounded region of class  $C^{2m}$  in  $\mathbb{R}^n$ , and let

$$au = \sum_{|\alpha|, |\beta| \leq m} (-1)^{|\alpha|} D^{lpha} a_{lpha eta}(t) D^{eta} ,$$

where  $D^{\alpha} = \partial^{|\alpha|} / \partial t_1^{\alpha_1} \cdots \partial t_n^{\alpha_n}$ ,  $|\alpha| = \alpha_1 + \cdots + \alpha_n$ , and for some  $\mu$ :  $0 < \mu < 1$ ,

$$a_{\alpha\beta} \in C^{\max(|\alpha|, |\beta|), \mu}$$
, all  $\alpha, \beta$ .

Moreover assume that  $\tau$  is formally self-adjoint and that there exists  $k_0 > 0$ , such that

$$\sum_{|\alpha|, |\beta| \leq m} \int_{\mathscr{Q}} a_{\alpha\beta}(t) D^{\alpha} y(t) D^{\beta} y(t) dt \geq k_0 \sum_{|\alpha| \leq m} \int_{\mathscr{Q}} |D^{\alpha} y|^2 dt ,$$

for all  $y \in C_0^m(\Omega)$ .

Let f(t, y) be uniformly Hölder continuous on  $\Omega \times R$  and satisfy (45), (46), (47) and (53) with

(57) 
$$1 \leq b$$
,  $b(n-2m) < n + 2m$ .

Then for any c > o the eigenvalue problem

(58) 
$$\tau y = \mu f(t, y) , \qquad D_y^{\alpha}|_{\delta \Omega} = 0 , \qquad |\alpha| \leq m-1 ,$$

has infinitely many eigenfunctions  $y \in C^{2m}(\Omega) \cap c^{m-1}(\overline{\Omega})$  and satisfying

$$\gamma(y) = \int_{a} \int_{0}^{y(t)} f(t, u) du dt = c .$$

*Proof.* It follows from results of [6] that (58) is equivalent to an integral equation of the form (2), and where (52) is satisfied for a(n-2m) < n. The theorem then follows from the remarks above, preceding its statement, and with the use of the arguments employed in the proof of the main theorem in [6]. The sharpness of the condition (53), (56) is shown by an example in [15].

For the special case where  $\tau = -\Delta$ , the hypothesis can be weakened slightly, compare Theorems 4 and 5 in [5].

We remark finally that X and Y in Theorem 1 can also be taken to be Orlicz spaces other than  $L^p$ . Continuity conditions for the integral operator A and the Nemytsky operator  $\Phi$ , when X and Y are Orlicz spaces, are given in [9]. In particular, by taking X and Y to be Orlicz spaces one can replace the growth condition (53), (57) in Theorem 3 by an exponential growth condition when  $\tau = -\Delta$  and n = 2; see for example the hypothesis of the main theorem of [10].

### References

1. F. E. Browder, Infinite dimensional manifolds and nonliner elliptic eigenvalue problems, Ann. of Math. 82 (1965), 459-477.

2. \_\_\_\_\_, Nonlinear eigenvalue problems and Galerkin approximations, Bull. Amer. Math. Soc. **74** (1968), 651-656.

3. E. S. Citlanadze, Existence theorems for minimaximal points in Banach spaces and their applications, Trud. Mosk. Mat. Obshch. 2 (1952), 235-274 (Russian).

4. C. V. Coffman, A minimum-maximum principle for a class of non-linear integral equations, J. d'Analyse Mathematique 22 (1969), 391-419.

5. \_\_\_\_\_, An existence theorem for a class of non-linear integral equations with applications to a nonlinear elliptic boundary value problem, J. Math. Mech. **18** (1968), 411-420.

6. \_\_\_\_\_, On a class of non-linear elliptic boundary value problems, J. Math. Mech. **19** (1969), 351-356.

7. R. E. Edwards, *Functional Analysis*, Holt Rinehart and Winston, Inc., New York, 1965.

8. M. A. Krasnosel'skii, Topological Methods in the Theory of Nonlinear Integral Equations, MacMillan, New York, 1964.

9. M. A. Krasnosel'skii and Ya. B. Rutickii, Convex Functions and Orlicz Spaces, Noordhoff, Groninger, 1961.

10. N. Levinson, Positive eigenfunctions for  $\Delta u + \lambda f(u) = 0$ , Arch. Rat. Mech. Anal. **11** (1962), 258-272.

11. R. A. Moore and Z. Nehari, Nonoscillation theorems for a class of nonlinear differential equations, Trans. Amer. Math. Soc. 93 (1959), 30-52.

12. Z. Nehari, On a class of nonlinear second order differential equations, Trans. Amer. Math. Soc. **95** (1960), 101-123.

13. Z. Nehari, On a class of nonlinear integral equations. Math. Zeit. 72 (1959), 175-183.

14. R. S. Palais, Lusternik-Schnirelman theory on Banach maniforlds, Topology 5 (1966), 115-132.

15. S. I. Pohozaev, Eigenfunctions of the equation  $\Delta u + \lambda f(u) = 0$ , Dokl. Akad. Nauk. SSSR 165 (1965), 36-39; Soviet Math. 5 (1965), 1408-1411.

16. J. T. Schwartz, Generalizing the Lusternik-Schnirelman theory of critical points, Comm. Pure App. Math. 17 (1964), 382-396.

17. V. I. Sobolev, On a nonlinear integral equation, Dokl. Akad. Nauk. SSSR 71 (1950), 831-834. (Russian).

18. M. M. Vainberg, Variational Methods for the Study of Nonlinear Operators, Holden-Day, Inc., San Francisco, 1964.

19. M. M. Vainberg, R. I. Kacurovskii, On the variational theory of nonlinear operators and equations, Dokl. Akad. Nauk. SSSR **129** (1959), 1199-1202. (Russian)

Received February 10, 1970. Research performed at the U.S.A.F. Aerospace Research Laboratories while in the capacity of an Ohio State University Research Foundation Visiting Research Associate under Contract F33615 67 C 1758.

CARNEGIE-MELLON UNIVERSITY PITTSBURGH, PENNSYLVANIA

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON Stanford University Stanford, California 94305

J. DUGUNDJI Department of Mathematics University of Southern California Los Angeles, California 90007

RICHARD ARENS

University of California Los Angeles, California 90024

# ASSOCIATE EDITORS

# SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA CALIFORNIA INSTITUTE OF TECHNOLOGY UNIVERSITY OF CALIFORNIA MONTANA STATE UNIVERSITY UNIVERSITY OF NEVADA NEW MEXICO STATE UNIVERSITY OREGON STATE UNIVERSITY UNIVERSITY OF OREGON OSAKA UNIVERSITY UNIVERSITY OF SOUTHERN CALIFORNIA STANFORD UNIVERSITY UNIVERSITY OF TOKYO UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF WASHINGTON \* \* \* \* AMERICAN MATHEMATICAL SOCIES

AMERICAN MATHEMATICAL SOCIETY CHEVRON RESEARCH CORPORATION NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The Pacific Journal of Mathematics is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

R. R. PHELPS University of Washington Seattle, Washington 98105

# Pacific Journal of Mathematics Vol. 36, No. 2 December, 1971

George E. Andrews, On a partition problem of H. L. Alder	279
Thomas Craig Brown, An interesting combinatorial method in the theory of	
locally finite semigroups	285
Yuen-Kwok Chan, A constructive proof of Sard's theorem	291
Charles Vernon Coffman, Spectral theory of monotone Hammerstein	
operators	303
Edward Dewey Davis, Regular sequences and minimal bases	323
Israel (Yitzchak) Nathan Herstein and Lance W. Small, <i>Regular elements in</i>	
P.Irings	327
Marcel Herzog, Intersections of nilpotent Hall subgroups	331
W. N. Hudson, Volterra transformations of the Wiener measure on the space	
of continuous functions of two variables	335
J. H. V. Hunt, An n-arc theorem for Peano spaces	351
Arnold Joseph Insel, A decomposition theorem for topological group	
extensions	357
Caulton Lee Irwin, Inverting operators for singular boundary value	
problems	379
Abraham A. Klein, <i>Matrix rings of finite degree of nilpotency</i>	387
Wei-Eihn Kuan, On the hyperplane section through a rational point of an	
algebraic variety	393
John Hathway Lindsey, II, On a six-dimensional projective representation of	
PSU <sub>4</sub> (3)	407
Jorge Martinez, Approximation by archimedean lattice cones	427
J. F. McClendon, On stable fiber space obstructions	439
Mitsuru Nakai and Leo Sario, <i>Behavior of Green lines at the Kuramochi</i>	
boundary of a Riemann surface	447
Donald Steven Passman, Linear identities in group rings.	457
Donald Steven Passman, Linear identities in group rings. II	485
David S. Promislow, The Kakutani theorem for tensor products of	
W*-algebras	507
Richard Lewis Roth, On the conjugating representation of a finite group	515
Bert Alan Taylor, On weighted polynomial approximation of entire	
functions	523
William Charles Waterhouse, <i>Divisor classes in pseudo Galois</i>	
extensions	541
Chi Song Wong, Subadditive functions	549
Ta-Sun Wu, A note on the minimality of certain bitransformation	
groups	553