

Pacific Journal of Mathematics

**INVERTING OPERATORS FOR SINGULAR BOUNDARY VALUE
PROBLEMS**

CAULTON LEE IRWIN

INVERTING OPERATORS FOR SINGULAR BOUNDARY VALUE PROBLEMS

C. L. IRWIN

Let S denote a Banach Space, B the bounded linear transformations on S , and let Q and A denote functions from $[0, \infty)$ into B with Q continuous. The objective here is to derive a Green's function K_A and hence an integral inverting operator R_A for the singular boundary value problem

$$(1) \quad \begin{cases} Y' - QY = H \\ A(0)Y(0) + \lim_{n \rightarrow \infty} A(c_n)Y(c_n) = 0, \end{cases}$$

where $\{c_n\}_{n=1}^{\infty}$ is a positive, increasing, unbounded number sequence and H is a continuous function from $[0, \infty)$ into S .

The method here provides Green's functions for singular boundary value problems associated with nonself-adjoint, as well as self-adjoint, linear differential expressions. The asymptotic boundary conditions in (1) permit one to extend some of the regular two-point boundary value problem techniques suggested by [3] and [4] to the singular case without being restricted to the Hilbert Space $L_2[0, \infty)$. Similar, but different, asymptotic boundary conditions are used by Codrington and Levinson in [2, Chapter 10], and by Benzinger [1].

As noted in § 3 of [3] there exists a unique continuous function M from $[0, \infty) \times [0, \infty)$ to B so that if each of x , t , and u is in $[0, \infty)$,

(i) $M_1(x, t) = Q(x)M(x, t)$ and $M(t, t) = I$

(ii) $M(x, t)M(t, u) = M(x, u)$

(iii) if H is a continuous function from $[0, \infty)$ to S and α is in S , then the only function Y such that $Y' - QY = H$ and $Y(0) = \alpha$ is given by

$$Y(x) = M(x, 0)\alpha + \int_0^x dt M(x, t)H(t)$$

for all x in $[0, \infty)$.

DEFINITION. A is a determinate boundary condition function for Q on c_1, c_2, \dots means that if H is a continuous function on $[0, \infty)$ and Y is a solution of the boundary value problem (1) for the non-homogeneous term H , then Y is unique.

NOTATION. If A is a boundary condition function for Q on c_1, c_2, \dots and n is a positive integer, let T_n denote the transformation

$$[A(0) + A(c_n)M(c_n, 0)].$$

THEOREM 1. *A is a determinate boundary condition function for Q on c_1, c_2, \dots if and only if the convergence of $\{T_n\alpha\}_{n=1}^\infty$ to zero implies that α is the zero of S.*

Proof. The proof follows from property (iii) of the M function and the linearity of the problem.

NOTATION. Let D_0 denote the continuous functions with compact support on $[0, \infty)$.

THEOREM 2. *Suppose A is a determinate boundary condition function for Q on c_1, c_2, \dots ; the following two statements are equivalent.*

(i) *There is an integral inverting operator R_A with kernel K_A of the form*

$$K_A(x, t) = \begin{cases} M(x, 0)K_A(0, 0)M(0, t) & \text{if } 0 \leq t \leq x \\ M(x, 0)[K_A(0, 0) - I]M(0, t) & \text{if } 0 \leq x < t \end{cases}$$

for boundary value problem (1) so that D_0 is a subset of the domain of R_A .

(ii) *There is a transformation π in B such that if α is in S, then $\{T_n(\pi\alpha)\}_{n=1}^\infty$ converges to $A(0)\alpha$.*

Proof. Assume (i) holds; if H is in D_0 and $U = R_A H$, then U is a solution of boundary value problem (1) and so

$$\lim_{n \rightarrow \infty} [A(0)U(0) + A(c_n)U(c_n)] = 0.$$

Let b denote a positive number so that if $x > b$, $H(x) = 0$; then, if $c_n > b$,

$$\begin{aligned} A(0)U(0) + A(c_n)U(c_n) &= A(0) \int_0^b dt [K_A(0, 0) - I]M(0, t)H(t) \\ &\quad + A(c_n) \int_0^b dt M(c_n, 0)K_A(0, 0)M(0, t)H(t) \\ &= T_n \left(K_A(0, 0) \int_0^b dt M(0, t)H(t) \right) \\ &\quad - A(0) \int_0^b dt M(0, t)H(t) \end{aligned}$$

and so

$$\lim_{n \rightarrow \infty} T_n \left(K_A(0, 0) \int_0^b dt M(0, t)H(t) \right) = A(0) \int_0^b dt M(0, t)H(t) .$$

Now, if α is in S , define H as

$$H(t) = \begin{cases} M(t, 0)(2 - 2t)\alpha & \text{if } 0 \leq t \leq 1 \\ 0 & \text{if } t > 1 . \end{cases}$$

H belongs to D_0 and $\int_0^b dt M(0, t)H(t) = \alpha$, so (ii) holds with $\pi = K_A(0, 0)$.

Now, assume (ii) holds; since A is a determinate boundary condition function for Q on c_1, c_2, \dots , π must be unique. Define K_A on $[0, \infty) \times [0, \infty)$ as

$$(2) \quad K_A(x, t) = \begin{cases} M(x, 0)\pi M(0, t) & \text{if } 0 \leq t \leq x \\ M(x, 0)[\pi - I]M(0, t) & \text{if } 0 \leq x < t \end{cases}$$

and let R_A denote the integral operator with kernel K_A . Let H be in D_0 and b denote a positive number so that if $x > b$, $H(x) = 0$. Define U on $[0, \infty)$ as

$$U(x) = \begin{cases} \int_0^x dt M(x, 0)\pi M(0, t)H(t) \\ \quad + \int_x^b dt M(x, 0)[\pi - I]M(0, t)H(t) & \text{if } 0 \leq x \leq b \\ \int_0^b dt M(x, 0)\pi M(0, t)H(t) & \text{if } x > b . \end{cases}$$

Differentiation yields that $U'(x) - Q(x)U(x) = H(x)$ for each x in $[0, \infty)$ and if $c_n > b$,

$$\begin{aligned} & A(0)U(0) + A(c_n)U(c_n) \\ &= T_n \left(\pi \int_0^b dt M(0, t)H(t) \right) - A(0) \int_0^b dt M(0, t)H(t) . \end{aligned}$$

By the definition of π , $\lim_{n \rightarrow \infty} [A(0)U(0) + A(c_n)U(c_n)] = 0$ and so (i) holds.

For the remainder of the paper suppose that A is a determinate boundary condition function for Q on c_1, c_2, \dots , condition (ii) in Theorem 2 holds and K_A is defined on $[0, \infty) \times [0, \infty)$ by (2). (Condition (ii) is implied, for example, in case the sequence $\{T_n\}_{n=1}^\infty$ converges in norm to a regular element of B .) Let D denote the set of continuous functions H on $[0, \infty)$ such that $\int_0^\infty dt K_A(x, t)H(t)$ exists for each x and furthermore, if U is defined as

$$U(x) = \int_0^\infty dt K_A(x, t)H(t) \quad \text{for } x \text{ in } [0, \infty) ,$$

then U is a solution of boundary value problem (1) for the nonhomo-

geneous term H . Let R_A denote the integral operator with kernel K_A and domain D ; i.e., if H belongs to D

$$(R_A H)(x) = \int_0^\infty dt K_A(x, t)H(t) .$$

Two aspects of the present development which differ from other treatments of Green's functions for singular boundary value problems are: (1) the Green's functions here are not necessarily square integrable in either place and (2) the domains of the associated integral inverting operators are not restricted to functions which are square integrable on $[0, \infty)$. However, the domain of R_A does depend upon the problem, i.e., upon the particular Q and A involved. This dependence is the subject of the following two theorems.

Two sets of continuous function on $[0, \infty)$ which are relevant to the description of D are defined as follows. Let D_1 denote the collection of continuous functions H on $[0, \infty)$ for which there exists a solution of (1) for the nonhomogeneous term H . Let D_2 denote the collection of continuous functions H on $[0, \infty)$ such that $\int_0^\infty dt(\pi - I)M(0, t)H(t)$ exists.

It is clear that D is a subset of the intersection of D_1 and D_2 ; not so obvious is the extent to which $D_1 \cap D_2$ is contained in D .

LEMMA. Suppose H belongs to $D_1 \cap D_2$; let Y denote the solution of (1) for H and let $X(x) = \int_0^\infty dt K_A(x, t)H(t)$ for all x in $[0, \infty)$, then

$$\begin{aligned} T_n [Y(0) - X(0)] &= [A(0)Y(0) + A(c_n)Y(c_n)] \\ &\quad + [A(0) - T_n \pi] \int_0^{c_n} dt M(0, t)H(t) \\ &\quad - T_n \int_{c_n}^\infty dt(\pi - I)M(0, t)H(t) \end{aligned}$$

for each positive integer n .

Proof. Let n denote a positive integer; property (iii) of the M function provides that

$$Y(c_n) = M(c_n, 0)Y(0) + M(c_n, 0) \int_0^{c_n} dt M(0, t)H(t) ,$$

so

$$\begin{aligned} T_n Y(0) &= A(0)Y(0) + A(c_n)M(c_n, 0)Y(0) \\ &= A(0)Y(0) + A(c_n)Y(c_n) - A(c_n)M(c_n, 0) \int_0^{c_n} dt M(0, t)H(t) . \end{aligned}$$

Also,

$$T_n X(0) = [A(0) + A(c_n)M(c_n, 0)] \int_0^\infty dt(\pi - I)M(0, t)H(t) .$$

A straightforward computation provides the result of the lemma.

The domain D of the inverting operator R_A may be studied for the following three cases.

Case 1. There is an increasing sequence of positive integers n_1, n_2, \dots such that $T_{n_i}^{-1}$ exists for all i and the transformation sequence $\{T_{n_i}^{-1}\}_{i=1}^\infty$ is uniformly norm bounded.

Case 2. There is an increasing sequence of positive integers n_1, n_2, \dots such that $T_{n_i}^{-1}$ exists for all i , but no subsequence of inverses is uniformly norm bounded.

Case 3. There is a positive integer N such that if $n > N$, then T_n^{-1} does not exist.

Note. Case 1 above is a sufficient condition for a function A from $[0, \infty)$ into B to be a determinate boundary condition function for Q on c_1, c_2, \dots .

THEOREM 3. *Suppose Case 1 above holds; if H is in $D_1 \cap D_2$ then H is in D if*

$$\lim_{i \rightarrow \infty} [A(0) - T_{n_i} \pi] \int_0^{c_{n_i}} dt M(0, t)H(t) = 0 .$$

Proof. By the lemma and existence of $T_{n_i}^{-1}$ for all i , we obtain in the notation of the lemma that

$$\begin{aligned} Y(0) - X(0) &= T_{n_i}^{-1}[A(0)Y(0) + A(c_{n_i})Y(c_{n_i})] \\ &\quad + T_{n_i}^{-1}[A(0) - T_{n_i} \pi] \int_0^{c_{n_i}} dt M(0, t)H(t) \\ &\quad - \int_{c_{n_i}}^\infty dt(\pi - I)M(0, t)H(t) \qquad \text{for each } i . \end{aligned}$$

H in D_2 provides that $\lim_{i \rightarrow \infty} \int_{c_{n_i}}^\infty dt(\pi - I)M(0, t)H(t) = 0$ and Y satisfies the asymptotic boundary condition so

$$Y(0) - X(0) = \lim_{i \rightarrow \infty} T_{n_i}^{-1}[A(0) - T_{n_i} \pi] \int_0^{c_{n_i}} dt M(0, t)H(t) .$$

Now, if $\lim_{i \rightarrow \infty} [A(0) - T_{n_i} \pi] \int_0^{c_{n_i}} dt M(0, t)H(t) = 0$, then $Y(0) - X(0) = 0$

and so $Y = X$, i.e., X is the unique solution of (1) for H and so H is in the domain of R_A .

A subcase of Cases 1, 2, and 3 above is that the transformation sequence $\{T_n\}_{n=1}^\infty$ be uniformly norm bounded, which occurs, for example, with $S = E_2$ and $T_n = \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$ n odd, $\begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}$ if n is even.

THEOREM 4. *Suppose $\{T_n\}_{n=1}^\infty$ is uniformly norm bounded; if H is in $D_1 \cap D_2$, then H is in D if and only if*

$$\lim_{n \rightarrow \infty} [A(0) - T_n \pi] \int_0^{c_n} dt M(0, t) H(t) = 0.$$

Proof. Let H denote a function in $D_1 \cap D_2$; by the lemma

$$\begin{aligned} T_n [Y(0) - X(0)] &= [A(0)Y(0) + A(c_n)Y(c_n)] \\ &\quad + [A(0) - T_n \pi] \int_0^c dt M(0, t) H(t) \\ &\quad - T_n \int_{c_n}^\infty dt (\pi - I) M(0, t) H(t) \end{aligned}$$

for each positive integer n . Where Y denotes the solution of (1) for H and X is defined on $(0, \infty)$ by

$$X(x) = \int_0^\infty dt K_A(x, t) H(t) \quad x \text{ in } [0, \infty).$$

Y satisfies the asymptotic boundary condition so

$$A(0)Y(0) + \lim_{n \rightarrow \infty} A(c_n)Y(c_n) = 0.$$

The transformation sequence $\{T_n\}_{n=1}^\infty$ is uniformly norm bounded and H is in D_2 so

$$\lim_{n \rightarrow \infty} T_n \int_{c_n}^\infty dt (\pi - I) M(0, t) H(t) = 0.$$

So

$$\lim_{n \rightarrow \infty} T_n [Y(0) - X(0)] = \lim_{n \rightarrow \infty} [A(0) - T_n \pi] \int_0^{c_n} dt M(0, t) H(t).$$

The result of the theorem follows from A being a determinate boundary condition function.

The following example illustrates the subcase for a Case 1 problem and shows that the domain of R_A may be a proper subset of $D_1 \cap D_2$.

EXAMPLE. Let c_1, c_2, \dots denote a positive, increasing, unbounded number sequence. Consider the singular boundary value problem associated with the differential expression $Ly = y''$ and the boundary condition function A defined as

$$A(x) = \begin{cases} \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} & \text{if } x = 0, \\ \begin{bmatrix} -1/[1 + \log(1 + x)] & x/[1 + \log(1 + x)] \\ 0 & -1 \end{bmatrix} & \text{if } x > 0, \end{cases}$$

We have $Q(x) = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ for $x \geq 0$, $M(x, t) = \begin{bmatrix} 1 & x - t \\ 0 & 1 \end{bmatrix}$ for all numbers x and t and if n is a positive integer,

$$T_n = \begin{bmatrix} \log(1 + c_n)/[1 + \log(1 + c_n)] & 0 \\ 0 & 1 \end{bmatrix}.$$

So, $\lim_{n \rightarrow \infty} T_n = I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ and $\pi = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$. A is a determinate boundary condition function for Q on c_1, c_2, \dots and K_A is calculated by equation (2).

Let $H(x) = \begin{bmatrix} 0 \\ 1/(1 + x^2) \end{bmatrix}$ $x \geq 0$. H is in D_1 since the function Y defined by

$$Y(x) = \begin{bmatrix} x \arctan x - \log + (1x^2)^{1/2} + (\pi/2)x - 1 \\ \arctan x + \pi/2 \end{bmatrix}$$

for $x \geq 0$ is a solution of the singular boundary value problem with nonhomogeneous term H . Also, $\int_0^\infty dt(\pi - I)M(0, t)H(t)$ exists so H is in D_2 and $\int_0^\infty dt K_A(x, t)H(t)$ exists for each $x \geq 0$. The function X defined by $X(x) = \int_0^\infty dt K_A(x, t)H(t)$ for $x \geq 0$ does not satisfy the asymptotic boundary condition and so H is in $D_1 \cap D_2$ but not in the domain of R_A .

It remains to more completely describe how the domain of R_A depends upon the problem and to investigate the complex numbers λ for which one obtains an inverting operator $R(A, \theta, \lambda)$ for the singular boundary value problem

$$\begin{cases} Y' - (Q + \lambda\theta)Y = H \\ A(0)Y(0) + \lim_{n \rightarrow \infty} A(c_n)Y(c_n) = 0 \end{cases}$$

where θ denotes a function from $[0, \infty)$ into B .

REFERENCES

1. H. E. Benzinger, *Green's functions on large intervals*. Preliminary Report, Notices Amer. Math. Soc. **16** (1969), 114.
2. E. A. Coddington and N. Levinson, *Theory of Ordinary Differential Equations*, McGraw Hill Book Co., Inc., 1955.
3. J. W. Neuberger, *Concerning boundary value problems*, Pacific J. Math. **10** (1960), 1385-1392.
4. A. J. Zettl, *Quasi-Differential Operators in Hilbert Space*, M. R. C. Technical Summary Report No. 845, 1968.

Received February 11, 1970. This paper is part of the author's doctoral dissertation which was directed by Professor John W. Neuberger.

WEST VIRGINIA UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R. PHELPS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLE

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics

Vol. 36, No. 2

December, 1971

George E. Andrews, <i>On a partition problem of H. L. Alder</i>	279
Thomas Craig Brown, <i>An interesting combinatorial method in the theory of locally finite semigroups</i>	285
Yuen-Kwok Chan, <i>A constructive proof of Sard's theorem</i>	291
Charles Vernon Coffman, <i>Spectral theory of monotone Hammerstein operators</i>	303
Edward Dewey Davis, <i>Regular sequences and minimal bases</i>	323
Israel (Yitzchak) Nathan Herstein and Lance W. Small, <i>Regular elements in P.I.-rings</i>	327
Marcel Herzog, <i>Intersections of nilpotent Hall subgroups</i>	331
W. N. Hudson, <i>Volterra transformations of the Wiener measure on the space of continuous functions of two variables</i>	335
J. H. V. Hunt, <i>An n-arc theorem for Peano spaces</i>	351
Arnold Joseph Insel, <i>A decomposition theorem for topological group extensions</i>	357
Caulton Lee Irwin, <i>Inverting operators for singular boundary value problems</i>	379
Abraham A. Klein, <i>Matrix rings of finite degree of nilpotency</i>	387
Wei-Eihn Kuan, <i>On the hyperplane section through a rational point of an algebraic variety</i>	393
John Hathway Lindsey, II, <i>On a six-dimensional projective representation of $PSU_4(3)$</i>	407
Jorge Martinez, <i>Approximation by archimedean lattice cones</i>	427
J. F. McClendon, <i>On stable fiber space obstructions</i>	439
Mitsuru Nakai and Leo Sario, <i>Behavior of Green lines at the Kuramochi boundary of a Riemann surface</i>	447
Donald Steven Passman, <i>Linear identities in group rings. I</i>	457
Donald Steven Passman, <i>Linear identities in group rings. II</i>	485
David S. Promislow, <i>The Kakutani theorem for tensor products of W^*-algebras</i>	507
Richard Lewis Roth, <i>On the conjugating representation of a finite group</i> . . .	515
Bert Alan Taylor, <i>On weighted polynomial approximation of entire functions</i>	523
William Charles Waterhouse, <i>Divisor classes in pseudo Galois extensions</i>	541
Chi Song Wong, <i>Subadditive functions</i>	549
Ta-Sun Wu, <i>A note on the minimality of certain bitransformation groups</i>	553
Keith Yale, <i>Invariant subspaces and projective representations</i>	557