

Pacific Journal of Mathematics

TRIANGULAR MATRICES WITH THE ISOCLINAL PROPERTY

LEROY JOHN DERR

TRIANGULAR MATRICES WITH THE ISOCLINAL PROPERTY

LEROY J. DERR

Consider the system V_n of $n \times n$, lower triangular matrices over the real numbers with the usual operations of addition, multiplication and scalar multiplication and with the additional property that $a_{i+1,j+1} = a_{i,j}$ (isoclinal). It is shown that V_n is a commutative vector algebra. The principal theorem (§ 3) establishes the existence of an algebraic mapping of V_n into a ring of rational functions. This mapping associates a special set of basis elements in V_n with the classically known Eulerian Polynomials.

Some properties of the space V_n are outlined in § 2. Section 4 gives an application of the main theorem to a problem which motivated this study, namely, the inversion of certain matrices in V_n for arbitrary dimension n . The matrices with first columns $[1^m, 2^m, \dots, n^m]$, $m = 0, 1, 2, \dots$, are considered in particular.

2. Properties.

2.1. Nomenclature. A matrix $A = \{a_{i,j}\}$ is called isoclinal if $a_{i+1,j+1} = a_{i,j}$ for all values of the indices permitted. Further we designate by V_n the class of $n \times n$ lower-triangular, isoclinal (L.T.I.) matrices (over the reals).

REMARK. The isoclinal property has appeared in studies of commutativity, under other names; for example see [4].

THEOREM 2.2. *The class V_n is a commutative sub-ring of matrices. Further, if $A \in V_n$ is nonsingular then $A^{-1} \in V_n$.*

Proof. A simple computation using the L.T.I. property will show multiplicative closure. Now, for $A, B \in V_n$ let $\{a_i\}, \{b_i\}$ be the elements of their first columns; these clearly define the matrices. The first column of AB is given by the Cauchy Product formula $\sum_{j=1}^k a_j b_{k-j+1}$ for $k = 1, 2, \dots, n$, which is commutative. Finally, if $A \in V_n$ is nonsingular then its diagonal element $a_1 \neq 0$ and the system $a_1 x_1 = 1, \sum_{j=1}^k a_j x_{k-j+1} = 0$ is solvable. Hence $X \in V_n$ and $X = A^{-1}$.

The algebra of V_n is closely allied to that of the polynomials over the reals, $P(Y)$. Let $A \in V_n$ be given by its first column $\{a_i\}$. Define $\phi_n: V_n \rightarrow P(Y)$ as the injection, $\phi_n(A) = \sum_{j=1}^n a_j Y^{j-1}$ and let $\pi_n: P(Y) \rightarrow V_n$ be the projection. We then have:

COROLLARY 2.3.

(i) π_n is a ring homomorphism onto, with kernel the principal ideal generated by Y^n .

(ii) $\pi_n \phi_n$ is the identity and $\pi_n \{\phi_n(A)\phi_n(B)\} = AB$.

Finally we note the useful operating rule for L.T.I. matrices that the product Ax , where x is a vector, is equivalent to AX where X is the L.T.I. matrix with first column x .

3. A Mapping of V_n by means of Eulerian Polynomials.

3.1. *Definitions and Nomenclature.* (i). The Eulerian Polynomials $A_m(\lambda)$ may be defined recursively, with $A_0(\lambda) = 1$, by:

$$A_{m+1}(\lambda) = (1 + m\lambda)A_m(\lambda) + \lambda(1 - \lambda)A'_m(\lambda).$$

(ii) Let $M_{m,n} \in V_n$ be defined, (giving the matrices' first columns), by:

$$M_{m,n} = (1^m, 2^m, \dots, n^m) \quad \text{for } m = 0, 1, 2, \dots.$$

(iii) Let $M_m(\lambda) = \sum_{p=1}^{\infty} p^m \cdot \lambda^{p-1}$, for $|\lambda| < 1$ and $m = 0, 1, \dots$.

(iv) Let $R = \{P(\lambda)/Q(\lambda)\}$ be the sub-ring of rational functions such that $Q(0) \neq 0$.

3.2. *Assertion.* (i) The matrices $M_{m,n}$ constitute a basis for V_n , $m = 0, 1, \dots, n - 1$.

(ii) $M_m(\lambda) = A_m(\lambda)/(1 - \lambda)^{m+1} \in R$.

The second part of the assertion may be easily proved by noting the recursion $M_{m+1}(\lambda) = d\{\lambda M_m(\lambda)\}/d\lambda$. The Eulerian Polynomials and rational functions closely related to the $M_m(\lambda)$ were used by Frobenius [2] in studies of Bernoulli numbers; a further exposition of their properties has been given by Carlitz [1] and they have been used by Riordan [3] in combinatorial analysis. The inversion of the matrices $M_{m,n}$ was the author's original problem and will be discussed in the next section. Now, using the above notations and definitions, we give the following algebraic mapping theorem.

THEOREM 3.3. *In the following diagram:*

$$V_n \xrightarrow{f_n} R \xrightarrow{h_n} R/\langle \lambda^n \rangle \xrightarrow{j_n} V_n$$

f_n is defined by identifying the basis elements of V_n , $f_n(M_{m,n}) = M_m(\lambda) \in R$. h_n is the natural homomorphism with kernel, $K(h_n)$, the principal ideal generated by λ^n . Then, there exists a ring isomorphism

j_n such that $j_n h_n(M_m(\lambda)) = M_{m,n}$.

Proof. We first note that an element γ of the ring $R/\langle \lambda^n \rangle$ has a unique antecedent in R of the form $\sum_{p=1}^n a_p \lambda^{p-1}$. This enables us immediately to define j_n as an additive isomorphism onto by $j_n(\gamma) = (a_1, a_2, \dots, a_n) \in V_n$. The product of two elements in $R/\langle \lambda^n \rangle$ can be expressed as $\sum_{p=1}^n c_p \lambda^{p-1} + K(h_n)$ where the c_p are formed by Cauchy Products of the unique antecedents. This gives a ring isomorphism since the multiplication in V_n is also Cauchy Product, truncated to n components.

The conclusion $j_n h_n(M_m(\lambda)) = j_n h_n\{\sum_{p=1}^n p^m \lambda^{p-1}\} = M_{m,n}$ follows at once by noting $M_m(\lambda) = \sum_{p=1}^n p^m \lambda^{p-1} + \sum_{p=n+1}^{\infty} p^m \lambda^{p-1}$. Other immediate consequences are:

- COROLLARY 3.4. (i) f_n is one-to-one and $j_n h_n f_n$ is the identity.
 (ii) $j_n h_n\{f_n(A) \cdot f_n(B)\} = AB$.

4. Application. By making use of the previous theorem:

$$M_{m,n}^{-1} = j_n h_n\{(1 - \lambda)^{m+1}\} \cdot j_n h_n\{1/A_m(\lambda)\} = BC^{-1}.$$

The matrix B is given by its first column (b_1, \dots, b_n) where $b_i = (-1)^{i-1} \frac{m+1}{(i-1)}$ if $i \leq m+2$ and $b_i = 0$ if $i > m+2$. The nonzero components for $C \in V_n$ are also finite in extent, being the coefficients of the Eulerian Polynomial $A_m(\lambda)$. These are known explicitly: $A(m, k) = \sum_{j=0}^k (-1)^{k-j} (j+1)^m \frac{m+1}{(k-j)}$, $k = 0, 1, \dots, m-1$. The problem is then reduced to finding C^{-1} which may be expressed in terms of a recursion on the $A(m, k)$. For $m = 0, 1, 2$ the solutions are trivial. For $m = 3$ the n^{th} component, c_n , of C^{-1} is $c_n = U_n(-2)$ (Chebyshev polynomials of the second kind). These are readily given in explicit form.

REFERENCES

1. L. Carlitz, *Eulerian numbers and polynomials*, Math. Mag. **32** (1959), 247-260.
2. G. Frobenius, *Über die Bernoullischen und die Eulerschen Polynome*, Sitzungsberichte der Preussische Akademie der Wissenschaften (1910), 809-847.
3. J. Riordan, *An introduction to combinatorial analysis*, Wiley, New York, 1958.
4. D. Suprunenko and R. Tyshevich, *Commutative Matrices*, Academic Press, New York 1968.

Received July 22, 1970

LOUISIANA STATE UNIVERSITY IN NEW ORLEANS

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 37, No. 1

January, 1971

Gregory Frank Bachelis and Haskell Paul Rosenthal, <i>On unconditionally converging series and biorthogonal systems in a Banach space</i>	1
Richard William Beals, <i>On spectral theory and scattering for elliptic operators with singular potentials</i>	7
J. Lennart (John) Berggren, <i>Solvable and supersolvable groups in which every element is conjugate to its inverse</i>	21
Lindsay Nathan Childs, <i>On covering spaces and Galois extensions</i>	29
William Jay Davis, David William Dean and Ivan Singer, <i>Multipliers and unconditional convergence of biorthogonal expansions</i>	35
Leroy John Derr, <i>Triangular matrices with the isoclinal property</i>	41
Paul Erdős, Robert James McEliece and Herbert Taylor, <i>Ramsey bounds for graph products</i>	45
Edward Graham Evans, Jr., <i>On epimorphisms to finitely generated modules</i>	47
Hector O. Fattorini, <i>The abstract Goursat problem</i>	51
Robert Dutton Fray and David Paul Roselle, <i>Weighted lattice paths</i>	85
Thomas L. Goulding and Augusto H. Ortiz, <i>Structure of semiprime (p, q) radicals</i>	97
E. W. Johnson and J. P. Lediaev, <i>Structure of Noether lattices with join-principal maximal elements</i>	101
David Samuel Kinderlehrer, <i>The regularity of minimal surfaces defined over slit domains</i>	109
Alistair H. Lachlan, <i>The transcendental rank of a theory</i>	119
Frank David Lesley, <i>Differentiability of minimal surfaces at the boundary</i>	123
Wolfgang Liebert, <i>Characterization of the endomorphism rings of divisible torsion modules and reduced complete torsion-free modules over complete discrete valuation rings</i>	141
Lawrence Carlton Moore, <i>Strictly increasing Riesz norms</i>	171
Raymond Moos Redheffer, <i>An inequality for the Hilbert transform</i>	181
James Ted Rogers Jr., <i>Mapping solenoids onto strongly self-entwined, circle-like continua</i>	213
Sherman K. Stein, <i>B-sets and planar maps</i>	217
Darrell R. Turnidge, <i>Torsion theories and rings of quotients of Morita equivalent rings</i>	225
Fred Ustina, <i>The Hausdorff means of double Fourier series and the principle of localization</i>	235
Stanley Joseph Wertheimer, <i>Quasi-compactness and decompositions for arbitrary relations</i>	253
Howard Henry Wicke and John Mays Worrell Jr., <i>On the open continuous images of paracompact Čech complete spaces</i>	265