Pacific Journal of Mathematics

LIE ALGEBRAS OF GENUS ONE AND GENUS TWO

JAMES W. BOND

Vol. 37, No. 3

March 1971

LIE ALGEBRAS OF GENUS ONE AND GENUS TWO

JAMES BOND

The genus of a finite dimensional algebra is the difference between its dimension and the number of elements in a minimal generating set for the algebra. In this paper the classification of finite dimensional Lie algebras of genus one and genus two is accomplished in four steps. First, it is shown that every such algebra is either solvable or contains a simple subalgebra. Second, the algebras containing a simple subalgebra are determined. Third, nonminimal genus one and two Lie algebras are shown to have a one, two dimensional ideal, J, respectively, with genus zero quotient. Fourth, the different possible L module structures for J are analyzed and completely determined except for the genus two index two nilpotent algebras.

The purpose of this paper is to classify and solve the isomorphism problem for finite dimensional Lie algebras of genus 1 and genus 2 over arbitrary fields. The classification problem was attacked for Lie algebras with genus less than three in Bond [1]. The classification obtained was somewhat unsatisfactory for genus 1 and genus 2 algebras because the results were presented by specifying different possible multiplication tables in terms of minimal generating sets for the algebras and therefore provided no ready access to the isomorphism problems. Nevertheless, the classification of genus 1 algebras was fairly short and had interesting consequences and therefore was made the basis of a short paper, which because of the perceptive comments of the referee grew into the present paper.

The concept of genus seems to have originated with Knebelman [5]. He obtained among other results a valid classification of the structure constants for genus 0 and 1 Lie algebras. His paper contains errors, which have been documented in Patterson [8] and Marshall [7]. Wallace [9] has obtained a classification of Lie algebras of dimension four over the complex numbers. It should be remarked that not unlike Wallace, who was unaware of Knebelman's work, the author was unaware of both Wallace's and Knebelman's work until quite recently.

Patterson [8] establishes the most far reaching results. His main theorem for algebras implies an interesting theorem about Lie algebras, (Theorem 2, 1) which seems to have gone unnoticed.

The author first became aware of the term "genus of a Lie algebra" through [7]. The results obtained in [7] are indirectly related to ours.

JAMES BOND

Throughout this paper use is made of the results in [2]. It should be pointed out that a minimal set of generators of a finite dimensional algebra is always a weak minimal generating set (a concept particular to [2]) for the algebra. The converse is false in general. The results needed from [2] are proved under the hypothesis of the existence of weak minimal generating sets with specific properties and hence hold under the stronger hypothesis of the existence of a minimal generating set with the same properties.

The following notation will be in force throughout the paper. L will denote a finite dimensional Lie algebra over an arbitrary field denoted by F. The elements of L will be denoted by small Roman letters and the elements of F by small Greek letters. J(a, b, c) = 0will stand for the Jacobi identity (ab)c+(bc)a+(ca)b=0. $\langle a, b, c, \cdots \rangle$ will denote the subalgebra of an algebra A generated by the elements a, b, c, \cdots of A. $\langle S, T, \cdots, a, b, \cdots \rangle$ will denote the subalgebra generated by the subsets of elements S, T, \cdots of A and the elements a, b, \cdots of A.

1. The genus of an algebra. A subset S of elements of a finite dimensional algebra A is said to minimally generate A if S generates A as an algebra and no subset of A having fewer elements than S generates A. A finite dimensional algebra A has genus k if A has a minimal generating set with m elements and dim L=k+m.

PROPOSITION 1. If B is a subalgebra of an algebra A then genus $B \leq genus A$.

Proof. Let M be a minimal generating set of B consisting of m elements. Let $A = B \dotplus C$, C a subspace of A. Then for any basis N of $C, M \cup N$ generates A. Therefore

genus $A \ge \dim A - (\dim C + m)$ = $(\dim A - \dim C) - m$ = $\dim B - m$ = genus B.

COROLLARY 1. If A has genus k then the dimension of a subalgebra generated by t elements of A is at most t + k.

PROPOSITION 2. Let I be an ideal of an algebra A. Then either A = B + I, B a subalgebra of A or genus A/I < genus A.

Proof. Let $\varphi: A \to A/I$ be a homomorphism with ker $\varphi = I$. Let M be a set of elements of A with $\varphi(M)$ minimally generating A/I. Let B be the subalgebra of A generated by M. Then genus $A \ge A$

genus $B \ge$ genus A/I. If genus B = genus A/I then dim B =dim A/I and hence $\varphi \mid B$ is an isomorphism.

THEOREM 1. Let L be a finite dimensional solvable genus k Lie algebra over any field. Then L is at most (k + 2)-step solvable.

Proof. If k = 0 then A is metabelian by Theorem 1 of [2].

Suppose the theorem has been proved for all finite dimensional Lie algebras of genus < k, k a positive integer.

Let L be of minimal dimension among the Lie algebras of genus k. Then genus $L^2 < \text{genus } L$. Hence L^2 is at most (k + 1)-step solvable, i.e., L is at most (k + 2)-step solvable.

Suppose the theorem has been proved for all genus k algebras of dimension less than n. Let dim L = n. Let I be a minimal non-zero ideal of L. Then I is abelian since I^2 is also an ideal of L.

If genus L/I < genus L then L/I is at most (k + 1)-step solvable. Hence L is at most (k + 2)-step solvable.

If genus L/I = genus L then the last proposition implies L = S + I, S a subalgebra. Any minimal generating set of S extended by any basis of I is a minimal generating set for L since genus S =genus L. It follows that Proposition 1, [2] applies. Hence Theorem 2, [2] applies with S = B and I = C. Consider K of Theorem 2, [2]. If K = S then $KI \subset K \cap I = 0$. Hence $L^2 = S^2$ and L is at most (k + 2)-step solvable by the induction hypothesis applied to S. If Kis not S then K + Fb = S, F the ground field, with $KC \subset B$ and bs - s in B. But, $KC \subset I$ and bs - s in I, so that KC = 0 and bs = s. Therefore from L = K + Fb + C it follows that $L^2 = (K + Fb + C)^2 \subset K^2 \oplus C$. Hence $(L^2)^2 = (K^2)^2$ and the theorem follows from the induction hypothesis applied to K.

Note that no inequality bounding the genus k by the length of the derived series is possible. Indeed, a metabelian Lie algebra can have any genus greater than one. Let A be a cyclic F[x] module with respect to a linear transformation T of A. Let L = A + Ft and define $t \cdot a = T(a), t^2 = 0, ab = 0$, for all a and b in A. Then L is 2-generated and hence has genus dim A - 1.

Recall that a Lie algebra L is said to be perfect if $L^2 = L$. The remaining results of this section relate assumptions on the genus of a Lie algebra to simplicity of the algebra.

PROPOSITION 3. Let P be a proper perfect subalgebra of a Lie algebra L over an infinite field F and x an element of L not in P. Then genus $\langle P, x \rangle >$ genus P.

JAMES BOND

Proof. Suppose $\langle P, x \rangle = P + Fx$. If M minimally generates P while $M \cup \{x\}$ does not minimally generate $\langle P, x \rangle$ then genus $\langle P, x \rangle >$ genus P. If $M \cup \{x\}$ minimally generates $\langle P, x \rangle$ apply Theorem 2, [2] to $\langle P, x \rangle$ with B = P and C = Fx to conclude $B = B^{2} \subset K$. Hence P is an ideal in $\langle P, x \rangle$. Apply Theorem 4, [2] and derive a contradiction to P perfect.

COROLLARY 2. Let S be a simple subalgebra of a Lie algebra L over an infinite field F. If genus S = genus L then S = L.

PROPOSITION 4. Let B be a subalgebra of a simple finite dimensional Lie algebra S over any field F. If genus B = genus S then B = S.

Proof. Suppose $B \neq S$. Let $S = B \dotplus C$, C a subspace of S. If dimension C is at least two by Theorem 2, [2] S has an ideal. If dim C = 1 then $B = K \dotplus Fb$, otherwise K is an ideal, from which it follows that $S^2 = (K \dotplus Fb \dotplus Fc)^2 \subset K + Fbc \neq S$, a contradiction.

COROLLARY 3. A simple Lie algebra of genus 1 over any field has dimension 3.

Proof. There exist a, b in S generating a 3 dimensional subalgebra.

We state without proof Theorem 1, [8] by E. M. Patterson:

Let A be an algebra of genus k and dimension n over an arbitrary field F. If $n \ge 3k + 3$ there exist elements a_1, a_2, \dots, a_{k+s} of A, with $s \le 2k$ such that the multiplication in A is given by $xy = \phi(y)x + \psi(x)y + \sum_{\lambda=1}^{k+s} \theta^{\lambda}(x, y)a_{\lambda}$, where ϕ and ψ are linear functions, θ^{λ} are bilinear functions, and a_1, \dots, a_k are generated by a_{k+1}, \dots, a_{k+s} . If F is not the field GF(2) the theorem is also true for n = 3k + 2.

If A is assumed to be a Lie algebra this result implies:

THEOREM 2. Let L be a Lie algebra of genus k and dimension n over an arbitrary field F. If $n \ge 3k + 4$ there exists an ideal J with dim $J \le 2k$ such that genus L/J = 0. If F is not the field GF(2), the theorem is true also for n = 3k + 3.

Proof. The multiplication in L is given by $xy = \phi(y)x + \psi(x)y + \sum_{\lambda=1}^{k+s} \theta^{\lambda}(x, y)a_{\lambda}$, $s \leq 2k$, ϕ and ψ linear functions, the θ^{λ} bilinear functions, a_1, \dots, a_{k+s} fixed linearly independent elements of L. Suppose further that the a^{λ} have been chosen so that the number of $\theta^{\lambda} \neq 0$

is minimal among all such multiplication rules for L.

If x, y and the a_{λ} are linearly independent xy = -yx implies $\phi(x) = -\psi(x)$. Then from $\phi(x + a) = -\psi(x + a)$ and $\phi(x) = -\psi(x)$ conclude by linearity $\phi(a) = -\psi(a)$. Hence $\phi = -\psi$. Take x, y, z and the a_{λ} linearly independent elements of L. Then the identity J(x, y, z) = 0 implies $\sum_{\lambda=1}^{k+s} \theta^{\lambda}(x, y) \ \phi(a_{\lambda}) = 0$ (the coefficient of z). Since this sum is zero for any x, y linearly independent of the a's by bilinearity it is zero for all x and y in L.

Suppose for some μ that $\phi(a_{\mu}) \neq 0$. Define

$$b_{\lambda} = egin{cases} a_{\lambda} - \phi(a_{\mu})^{-1} \phi(a_{\lambda}) a_{\mu} \ ext{ for } \lambda
eq \mu \ a_{\lambda} \ ext{ for } \lambda = \mu \ . \end{cases}$$

Then $xy = \phi(y)x - \phi(x)y = \sum_{\substack{\lambda=1\\\lambda\neq\mu}}^{k+s} \theta^{\lambda}(x, y)b_{\lambda}$ for all x and y in L. This contradicts the choice of the a_{λ} . Hence for all $\lambda, \phi(a_{\lambda}) = 0$, i.e. the a_{λ} span an ideal J. Obviously, genus L/J = 0.

COROLLARY 4. If L is a genus 1 Lie algebra over any field F and dim $L \ge 7$ then L has a 1-dimensional ideal with genus 0 quotient.

Proof. By Patterson, Theorem 2, [8], the s occurring in the last theorem can be taken equal to zero when k = 1.

2. The structure of finite dimensional Lie algebras of genus 1 and 2.

PROPOSITION 5. There are no perfect 4-dimensional Lie algebras over any field F.

Proof. If genus L = 0 then L is not perfect by Theorem 1, [2]. If genus L = 1 let L = B + Fc, B a 2-generated subalgebra of L. Apply Theorem 2, [2]. Either B = K and c is not in L^2 or B = K + Fb and $L^2 \subset K + Fbc$.

It remains to show that a 2-generated 4-dimensional Lie algebra cannot be perfect. Let a, b, ab, and a(ab) be a basis for L and adjust b so that b(ab) is in Fa + Fb + Fab. Suppose a(a(ab)) = $\alpha a + \beta b + \gamma ab + \delta a(ab)$ and $b(ab) = \alpha' a + \beta' b + \gamma' ab$. Then b(a(ab) = $\beta' ab + \gamma' a(ab)$ and $(ab)(a(ab)) = a(b(a(ab))) - b(a(a(ab))) = (\alpha \gamma' - \gamma \alpha')a +$ $(\beta \gamma' - \gamma \beta')b + (\alpha - \delta \beta')ab + \beta' a(ab)$ so that the whole multiplication table is determined. Observe L is a Lie algebra if and only if the parameters are so chosen that J(a, ab, a(ab)) = 0 and J(b, ab, a(ab)) = 0. The first Jacobi identity is equivalent to (1) $2\alpha - \delta \beta' = 0$, (2) $2\beta \gamma' - \alpha \delta = 0$, (3) $2\beta \beta' - \delta(\beta \gamma' - \gamma \beta') = 0$, and (4) $\beta' \alpha + \beta \alpha' - \delta(\alpha \gamma' - \gamma \alpha') = 0$. Use 1) to rewrite the coefficient of ab in ab(a(ab)) as $-\alpha$. Then find that the second Jabobi identity is equivalent to (5) $2\beta'\gamma + \alpha'\delta = 0$, (6) $\alpha\beta' + \alpha'\beta + 2\gamma'(\gamma'\beta - \beta'\gamma) = 0$, and (7) $2\alpha\alpha' + 2\gamma'(\gamma'\alpha - \alpha'\gamma) = 0$.

First suppose $\alpha' \neq 0$. Replace a by $\alpha' = \alpha' a + \beta' b$ and obtain with change of notation $\beta' = 0$. Then the equation (5) implies $\delta = 0$ and (4) that $\beta = 0$ and hence b is not in L^2 .

Second suppose $\alpha' = 0$. First, we show if char $F \neq 2$ then $\alpha = 0$. It would then follow that α is not in L^2 . By (7) $\alpha \gamma' = 0$ so that (4) implies $\alpha \beta' = 0$, which implies by multiplication of (1) by α that $\alpha = 0$. Finally suppose $\alpha \neq 0$. Then char F = 2 and (2) implies $\delta = 0$ and (4) $\beta' = 0$. Here $\alpha \alpha + \beta b$ is the only linear combination of α and b to occur in L^2 .

THEOREM 3. Let L be a finite dimensional Lie algebra over any field F. Suppose genus $L \leq 2$. Then L is either solvable or contains a simple subalgebra.

Proof. This theorem is proved by induction on the dimension of L. Every Lie algebra of dimension less than 3 is solvable. Suppose the theorem proved for all Lie algebras of genus k of dimensions $\leq n, k$ fixed.

Let L be an (n + 1)-dimensional genus k Lie algebra containing no simple subalgebras. A minimal ideal I of L is solvable and hence abelian. If L/I is solvable then L is solvable. Otherwise L/I contains a simple subalgebra T. Let S be the subalgebra generated by the preimage of a minimal generating set of T under the natural map of L onto L/I. S cannot be solvable for it has a nonsolvable homomorphic image. If $S \neq L, S$, and so L, would have a simple subalgebra. Hence S = L and so T = L/I.

Genus L/I < genus L since the preimage of a minimal generating set of L/I is a generating set of L. Since all genus 0 Lie algebras are solvable the theorem has been proved for k = 0, 1. If k = 2, dim L/I = 3 by Proposition 2 and Corollary 3. Then dim L = 4 since L is generated by two elements. Thus, by Proposition 5, $L^2 \neq L$ so that L^2 (and hence L) is solvable.

The very last argument fails for k=3. For there is a 2-generated 5-dimensional perfect Lie algebra with a 3-dimensional simple quotient algebra. For example: L = S + A, with S the 2 by 2 matrices of trace zero operating on the 2-dimensional underlying vector space of the abelian ideal A.

THEOREM 4. If a finite dimensional genus 1 Lie algebra over any field F contains a simple subalgebra S then S = L.

596

Proof. Suppose $0 \neq x$ in L - S. Then dim $\langle S, x \rangle = 4$. Therefore $\langle S, x \rangle^2 \neq \langle S, x \rangle$ by Proposition 5 and hence equals S.

Consider D_x on L^2 . Let m(t) and c(t) denote its minimal and characteristic polynomials, respectively. Then deg $m(t) \leq 2$, otherwise L^2 would have a 2-generated subspace of dim > 2, contradicting genus L = 1. The classical decomposition of L^2 into cyclic subspaces for D_a then contains a 1-dimension subspace. Therefore, a linear factor, $t - \lambda$, which divides m(t). Hence $c(t) = (t - \lambda)^3$. It follows that L^2 has two eigenvectors for λ , their product cannot be zero since L^2 is perfect and 3-dimensional, hence corresponds to eigenvalue 2λ . Thus $\lambda = 0$, x is central. The argument applies to x + a for any a in L^2 , a contradiction.

LEMMA 1. A genus 2 Lie algebra L over any field with dim L > 5 contains a proper genus 2 subalgebra.

Proof. If a genus 2 Lie algebra L has more than six generators then there exist a, b, c, d with dim $\langle a, b, c, d \rangle = 6$.

Suppose a, b, c, d, ab, cd is a basis for A. Consider

 $ac = \alpha a + \beta b + \gamma c + \delta d + \mu ab + \kappa cd$.

If $\mu \neq 0$, a, b, and c generate a genus 2 subalgebra. If $\kappa \neq 0$, a, d, c generate a genus 2 subalgebra. Hence $ac = \alpha a + \gamma c$, or done. Similarly, $bc = \beta_1 b + \gamma_1 c$, $ad = \alpha_2 a + \delta_2 d$, $bd = \beta_3 b + \delta_3 d$, or done. Then a + c, b, and d generate a genus 2 subalgebra.

PROPOSITION 6. A 5-dimensional simple Lie algebra over any field is 2-generated.

Proof. By Corollary 3 genus $L \neq 1$ and by earlier work genus $L \neq 0$. There remains the possibility that L has genus 2, i.e., L is minimally 3-generated. So suppose a, b, and c minimally generate L.

If B is a 4-dimensional subalgebra of L then B has genus less than 2 by Proposition 4.

Suppose a, b, ab are linearly independent. If ac and bc are both in Fa + Fb + Fc + Fab then so also is (ab)c = a(bc) - b(ac); for a(ab)and b(ab) are in Fa + Fb + Fab. Therefore with a change of notation, if necessary, L has basis a, b, c, ab, cb.

If $ac = \alpha a + \beta b + \gamma c + \delta ab + \mu cb$ replace a by $a + \mu b$ and c by $c - \delta b$ and find to avoid L being 2-generated that $ac = \alpha' a + \gamma' c$. By symmetry between a and c it can be supposed that $ac = \varepsilon c$, $\varepsilon = 0$ or 1. Let

$$egin{aligned} a(ab) &= lpha a + eta b + \gamma ab & c(cb) &= \mu c + \delta b + \kappa cb \ b(ab) &= lpha' a + eta' b &= \gamma' ab & b(cb) &= \mu' c + \delta' b + \kappa' cb \ c(ab) &= lpha'' a + eta'' b + \gamma'' ab + \mu'' c + \kappa'' cb \ . \end{aligned}$$

Suppose $\varepsilon = 0$. To avoid c in $\langle a, b + c \rangle$ from a(a(b + c)) conclude $\beta = 0$ and from (b + c)(a(b + c)) conclude $\kappa'' = 0$. To avoid a in $\langle c, b + a \rangle$ from c(c(b + a)) conclude $\delta = 0$ and from (b + a)(c(b + a)) conclude $\gamma' = 0$. Then J(c, a, ab) = 0 implies $\beta'' = 0$. Either $\beta' \neq 0$ or $\delta' \neq 0$ because b is in L^2 , say $\beta' \neq 0$ without loss of generality. Replace c by $c - \beta'^{-1}\delta'a$ and find that it can be supposed that $\delta' = 0$, hence $\mu = 0$. J(a, b, ab) = 0 implies $\alpha = \beta', \gamma = 0$, and $\gamma' = 0$. From b((a + c)b) conclude $\eta' = 0$. Note c(ab) = a(cb). Calculate (cb)(ab) using J(c, b, ab) = 0 and using J(cb, a, b) = 0 find that $\beta' = -2\mu''$, so that char $F \neq 2$ and $2\alpha'' = 0$; hence $\alpha'' = 0$. Then the subalgebra Fc + Fbc is an ideal of L because each generator carries it into itself.

Suppose $\varepsilon = 1$. From (b + c)(a(b + c)) conclude $\eta' = 1$. Note a(cb) = (ac)b + c(ab). Calculate the coefficient of cb in J(c, a, ab) = 0 and find that $\beta + \gamma = 1$. Then c is in $\langle a, b + c \rangle$ follows from a(a(b + c)).

LEMMA 2. Let L be a 5-dimensional Lie algebra over any field F. Let B be a proper ideal of L containing a proper simple subalgebra S. Then genus L = 3.

Proof. S must be genus 1 and 3-dimensional and hence B must be 4-dimensional and have genus 2.

Let a, b, ab, and a(ab) be a basis for B. Some linear combination of a and b is an element of B^2 . Suppose $b + \lambda a$ is an element of B^2 for some λ in F. If $ac = \alpha''a + \beta''b + \gamma''ab + \delta''a(ab)$ set $c' = c - \delta''ab - \gamma''b$ and find that $ac' = \alpha a + \beta b$. If $\beta \neq 0$ a and c' generate L. If $\beta = 0$ let b' = b + c'. Claim: a and b' generate L. Since the multiplication table of B is determined by the products b(ab) and a(a(ab)) b occurs in one of these products. The algebra generated by a and b' contains ab and a(ab) because $ab' = ab + \alpha a$ and a(ab') = a(ab).

Suppose $b + \lambda a$ is not an element of B^{2} for any λ in F. Suppose a, b, chosen so that $b(ab) = \alpha' a + \gamma' ab$ and $(*)a(a(ab)) = \alpha a + \gamma ab + \delta a(ab)$. Then S = Fa + Fab + Fa(ab).

Observe if s, t, and st are a basis of a simple Lie algebra L with $s(st) = \alpha s + \beta t + \gamma st$ and $t(st) = \alpha' s + \beta' t + \gamma' st$ then from J(s, t, st) = 0 it follows that $\gamma(\alpha' s + \beta' t) + \gamma'(\alpha s + \beta t) = 0$. This is a linear relation

598

between the contributions of s(st) and t(st) to Fs + Ft and therefore $\gamma = 0$ and $\gamma' = 0$, otherwise S would not be perfect.

Therefore since a and ab generate S from (*) it follows that $\delta = 0$ and from $(ab)(a(ab)) = \gamma'\alpha a + \gamma'\gamma ab + (\gamma'\delta - \gamma')a(ab)$ that $\gamma'\delta - \gamma' = 0$, which implies $\gamma'=0$. Hence (ab)(a(ab))=0, contradicting the simplicity of S.

THEOREM 5. Let L be a finite dimensional genus 2 Lie algebra over any field F. Suppose L contains a simple subalgebra. Then dim L = 4 and L^2 is simple 3-dimensional, Furthermore, if char $F \neq 2$ then $L = L^2 + Fz$, Fz the center of L.

Proof. Lemma 1 and Proposition 4 imply a simple genus 2 Lie algebra is at most 5-dimensional. Proposition 5 implies there are no 4-dimensional simple Lie algebras and Proposition 6 that there are no genus 2 5-dimensional simple algebras. Therefore L must contain a simple genus 1 Lie algebra S.

Suppose B is a 5-dimensional subalgebra of L containing S. Then genus $B \leq \text{genus } L$ and hence B is not simple. Let I be a proper ideal of B of maximal dimension. By Lemma 2 and Proposition 1, I cannot contain S so it follows that B = S + I. S has dimension 3 so that if $B = S \oplus I$ then S would be an ideal of larger dimension than I. This means that I cannot be trivial as an S module and hence I must be abelian since the other 2-dimensional Lie algebra has a solvable derivation algebra. Since $S^2 = S$ the elements of S must operate on I as transformations of trace 0. Considering that S has dimension 3 while I has dimension 2 we see that B is isomorphic to the semi-direct sum of a 2-dimensional abelian Lie algebra V and the Lie algebra of transformation of trace 0 on V.

This being the case the characteristic of F cannot be 2 since then the 2×2 transformations of trace 0 are solvable.

Let V = Fe + Ff and $a = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, $b = \begin{pmatrix} 00 \\ 10 \end{pmatrix}$, and $c = \begin{pmatrix} 01 \\ 00 \end{pmatrix}$ be a basis for S. Then a + b and c + 2e + f generate B. To see this observe (a + b)(c + 2e + f) - (c + 2e + f) = c - a. Therefore genus $B \ge 3$, contradicting Proposition 1.

If char $F \neq 2$ then S has a basis so that $a(ab) = \alpha b$ and $b(ab) = \beta a$ (Jacobson, page 14, [4]). Let L = S + Fx. Then $xa = \alpha'a + \beta'b + \gamma'ab$ and $xb = \alpha''a + \beta''b + \gamma''ab$. Set $z = x + \gamma'b - \gamma''a + \alpha^{-1}\beta'ab$, then $za = \alpha'a$ and $zb = \alpha''a + \beta''b$. Thus $z(ab) = (\alpha' + \beta'')ab$. From J(z, a, ab) = 0 conclude $\alpha' = 0$ and $\alpha'' = 0$ and from J(z, b, ab) = 0 that $\beta'' = 0$.

3. A characterization of nonminimal genus 1 and 2 Lie algebras. A Lie algebra L will be called a minimal genus k Lie

algebra, if genus B = genus L, B a subalgebra of L, implies B = L.

Suppose L is a finite dimensional Lie algebra with a k-dimensional ideal J with genus L/J = 0. Then genus $L \leq k$ because if dim L - k elements were to generate L their cosets would generate L/J. It is surprising that conversely nonminimal finite dimensional genus 1 and 2 Lie algebras always have ideals J of dimension 1 and 2, respectively, with genus 0 quotients, except possibly for F = GF(2).

There do exist solvable minimal genus 1 and 2 Lie algebras over some fields without such ideals. Indeed, the construction of metabelian Lie algebras in section one of arbitrarily high genus, provide such an example whenever A can be also taken as an irreducible F[x]module. This is equivalent to requiring that the characteristic polynomial of T is irreducible.

It has been shown that a genus 1 Lie algebra L, dim $L \ge 7$, had a 1-dimensional ideal J with genus L/J = 0 (Corollary 4). The next theorem shows this to be the case for dim $L \ge 4$, which is clearly the best possible result, since there are 3-dimensional genus 1 simple Lie algebras over every field.

THEOREM 6. Let L be a nonminimal finite dimensional genus 1 Lie algebra over any field F. Then L has a 1-dimensional ideal J with genus L/J = 0.

Proof. Let $L = B \dotplus C$, C a subspace and B a 3-dimensional subalgebra. Apply Theorem 2, [2] to conclude $B = \langle a, b \rangle$ with (1) ac - c, bc in B for all c in C, $B^{2} \subset Fb \dotplus Fab$, or (2) $BC \subset B$. (1) If $ac = c + \alpha'a + \beta'b + \gamma'ab$, c in C, then $ac' = c' + (\beta' + \gamma')b$, where $c' = c - \gamma'b + \alpha'a$, so that ac' = c', otherwise genus $\langle a, c' \rangle > 1$. Therefore C can be taken so that ac = c for all c in C. Suppose $bc = \alpha a + \beta b + \gamma ab$. Then $b(c + \gamma a) = \alpha a + \beta b$ and therefore $\alpha = 0$, otherwise genus $\langle c + \gamma a, b \rangle > 1$. Next $(a + b)(c + \gamma a) = (c + \gamma a) - \gamma a + \beta b$ so that $\beta = -\gamma$, i.e. $bc = \beta(b - ab)$. If $a(ab) = \gamma b + \delta ab$ then set $c' = c + ab - (\delta - 1)b$ and conclude from $ac' = c' + (\gamma + \delta - 1)b$ that $\delta = 1 - \gamma$.

Set J = F(b - ab). Then $a(b - ab) = -\gamma(b - ab)$ so that $aJ \subset J$. Next, $(ab)c = -\beta(1 + \gamma)(b - ab)$ so that $cJ \subset J$. J(a, b, ab) = 0 implies $b(ab) = \delta'ab$ because a(ab) and b(ab) are in Fb + Fab. Then

$$(b + c)(a(b + c)) = b(ab) + (b - ab)c$$

and must be in F(b+c) + Fa + F(ab+c) to avoid genus $\langle b+c, a \rangle > 1$. Hence $\delta' = 0$. Therefore $bJ \subset J$ and J is an ideal of L.

For c, d in C, cd is either an eigenvector for D_a for the eigenvalue 2 or cd = 0. Note a is a candidate if char F = 2. If cd = b', b'

in B and not in Fb + Fab, then every term of the derived series of L would contain C + Fa modulo Fb + Fab, contradicting L solvable by Theorem 4. If $\gamma \neq -1$, $C + Fa + F(ab + \gamma b) + F(b - ab)$ is an eigenspace decomposition of L for D_a and cd must be in J since $a(ab + \gamma b) = ab + \gamma b$. If $\gamma = -1$ then Fb + F(ab - b) is an indecomposable eigenspace for $D_a (ab = b + (ab - b))$. Again cd must be in J. Thus genus L/J = 0. (2) Clearly C can be taken so that ac and bc are in Fa + Fb for all c in C. Hence to avoid a genus 2 subalgebra, $ac = \alpha a$ and $bc = \alpha b$. If $\alpha \neq 0$ replace $\langle a, b \rangle$ by $\langle a + c, b \rangle$ reducing this case to (1). To avoid a genus 2 subalgebra from (a + c)((a + c)b) and a(a(b + c)) conclude a(ab) is in Fab and from b((a + c)b) and (b + c)(a(b + c)) that b(ab) is in Fab. Thus $L^2 = Fab$. Hence if dim C = 1 the proof is complete.

Suppose c, d are linearly independent elements of C with cd = e + x, e in C, x in $\langle a, b \rangle$. Note from (a + c)d one may suppose e is in Fc, and from c(a + d) that e is in Fd, otherwise this case is reduced to (1). Hence it may be supposed that e = 0. If $x = \alpha a + \beta b + \gamma ab$ then $(c - \gamma a)(d + b) = \alpha a + \beta b$ so that $\alpha \neq 0$ would imply genus $\langle c - \gamma a, d + b, b \rangle > 1$ while $\beta \neq 0$ implies genus $\langle c - \gamma a, d + b, a \rangle > 1$. Thus $J = L^2 = Fab$.

The proof of Theorem 6 provides an excellent model for the proof of the next theorem. It should be remarked that a simpler proof would be possible for the next theorem if fields of char 2 and GF(3) were excluded. We were unable to handle the case when F = GF(2).

THEOREM 7. Let L be a nonminimal finite dimensional genus 2 Lie algebra over any field F, except GF(2), the field of two elements. Then L has a 2-dimensional ideal J with genus L/J = 0.

Proof. I. Suppose there exists a subalgebra B of L with dim B = 4 and genus B = 2. Let C be a subspace complement of B in L. Theorem 2, [2] can be applied. There are three cases to be considered. There exists a basis a, b, ab, a(ab) of B such that for every c in C:

(1) ac and bc-c are in B, which $b(ab) = \delta_0 a + \delta_1 ab$ and $a(a(ab)) = \gamma_0 a + \gamma_1 ab + \gamma_2 a(ab)$.

(2) ac - c and bc are in B, $B^2 \subset Fb \dotplus Fab \dotplus Fa(ab)$.

(3) ac and bc are in B.

(1) Let $bc = c + \alpha' a + \gamma' a(ab)$ and $ac = \alpha a + \beta ab + \gamma a(ab)$, where c has been taken so that bc does not involve b or ab, while if ac involved b then genus $\langle a, c - \beta b - \gamma ab \rangle > 2$.

Set $c_{\lambda} = c - (\gamma + \lambda \gamma')ab - (\beta + \lambda(\gamma + \lambda \gamma')(1 - \delta_1))b$. Then $(a + \lambda b)c_{\lambda}$

can be expressed as a linear combination of c_{λ} , $a + \lambda b$, and b. The coefficient of b must be zero, otherwise genus $\langle a + \lambda b, c_{\lambda} \rangle > 2$. Therefore (*) $\lambda^{2}\gamma'(1 + \delta_{0} - \delta_{1}) + \lambda(\alpha' + \gamma(1 + \delta_{0} - \delta_{1})) + (\beta - \alpha) = 0$ for all nonzero λ in F.

The candidate for J is F(a + ab) + Fa(ab). To show that J is an ideal with genus L/J = 0 it suffices to show (1) $\delta_1 = 1 + \delta_0$ and (2) $\gamma_0 = \gamma_1$. This is because (1) implies $b(a + ab) = \delta_0(a + ab)$ and always $b(a(ab)) = \delta_1 a(ab)$ by J(a, b, ab) = 0, while (2) implies a(a(ab))is in J. Hence J is an ideal in B. Also, (1) implies $\alpha' = 0$ and $\alpha = \beta$ from (*) because $F \neq GF(2)$. Then bc - c and ac are in J. The Jacobi identities J(a, b, c) = 0, and J(a, ab, c) = 0 then imply (ab)cand (a(ab))c are in J. Therefore J is an ideal in L. If $cd = e + \lambda a + \mu b + J$, c, d, e in C, j in J, then conclude from J(b, c, d) = 0 that $cd = j + \mu b$ (calculate modulo J). Similarly J(a, c, d) = 0 implies $\mu = 0$. Thus genus L/J = 0.

 $J(a, b, a(ab)) = 0 \text{ implies } (ab)(a(ab)) = (\gamma_0\delta_1 - \gamma_1\delta_0)a + \gamma_0ab. \text{ From}$ the coefficient of a(ab) in J(a, ab, a(ab)) = 0 conclude $2\gamma_0 = 0$. From J(b, ab, a(ab)) = 0 conclude $\gamma_2\delta_0 = 0$ and $2\gamma_1\delta_0\delta_1 = 0$. Note, also J(a, b, c) = 0 implies $(ab)c = (\alpha - \beta\delta_0 + \gamma_0\gamma')a + (\alpha + \beta - \beta\delta_1 + \gamma_1\gamma')ab + (\gamma - \gamma\delta_1 + \gamma_2\gamma')a(ab).$

Suppose $bc = c + \alpha' a + \gamma' a(ab)$, $\gamma' \neq 0$, and c cannot be chosen so that $\gamma' = 0$. Then $\delta_1 = 1$, otherwise take $c' = c + (1 - \delta_1)^{-1} \gamma' a(ab)$. Note (1) holds unless F = GF(3) because of (*) and because F = GF(2)has been excluded by hypothesis. Suppose for the moment that $\delta_0 \neq 0$. Then $\gamma_0 = 0$, $\gamma_1 = 0$, and $\gamma_2 = 0$. From the coefficient of abin J(a, ab, c) = 0 conclude $\alpha = 0$, and from the coefficient of a that $\beta = 0$, contradicting (*). To establish (2) calculate J(b, ab, c) = 0 and find that $2\alpha + \gamma_1 \gamma' = 0$ and $2\alpha + \gamma_0 \gamma' = 0$. Multiplication of the last relation by 2 implies $2\alpha = 0$ since $2\gamma_0 = 0$. Therefore $\gamma_0 = \gamma_1 = 0$.

Suppose bc = c. The (*) implies $\alpha = \beta$. To avoid c in $\langle b, a+c \rangle$ conclude from b((a+c)b) that (1) holds. Then $(ab)c = (\alpha - \alpha\delta_0)(a+ab) - \gamma\delta_0a(ab)$. J(b, ab, c) = 0 implies $2\alpha - \alpha\delta_0 = 0$. Use $\gamma_2\delta_0 = 0$, $\alpha\delta_0 = 2\alpha$, and $2\gamma_0 = 0$ to find because of J(a, ab, c) = 0 that $a(ab)c = -\gamma(\gamma_0 - \gamma_1\delta_0)a - \gamma(\gamma_0 + \gamma_1\delta_0)ab - \alpha a(ab)$. Set $a' = a + \lambda c$. Then $a'b = ab - \lambda c$, $a'(a'b) = \lambda'(a(ab)$, where $\lambda' = 1 - \lambda(\gamma - \gamma\delta_0)$, and $a'(a'(a'b) = \lambda'(\gamma_0 + \lambda\gamma(\gamma_0 - \gamma_1\delta_0))a + \lambda'(\gamma_1 + \lambda\gamma(\gamma_0 + \gamma_1\delta_0))ab + \lambda'(\gamma_2 + \lambda\alpha)a(ab)$. Finally, the discriminant of a', b, a'b, a'(a'b), a'(a'(a'b)) with respect to a, b, ab, a(ab), c is $\lambda'^2\lambda^2[(\gamma_1 - \gamma_2) + 2\lambda\gamma_1\delta_0]$, which must be zero for all λ in F to avoid genus $\langle a', b \rangle = 3$. This establishes (2) with the possible exception of F = GF(3). Here $\gamma_0 = 0$ and $\gamma_1\delta_0\delta_1 = 0$. Done unless $\delta_1 = 0$, in which case $\delta_0 = -1$. Then $\lambda' = 1 - 2\alpha$ and $\gamma_1 - \gamma_0 + 2\lambda\gamma_1\delta_0 = \gamma_1(1-2\lambda)$ so that there exists a choice of λ so that $\gamma_0 = \gamma_1$.

The remaining cases are relatively easy.

(2) Take C so that ac = c and suppose $bc = \alpha_c a + \beta_c b + \gamma_c ab + \delta_c a(ab)$ for all c in C. Note if $b(ab) = \delta_0 b + \delta_1 ab + \delta_2 a(ab)$, with $\delta_2 \neq 0$, setting $b' = b - \delta_2 a$ reduces this case to (1). Hence suppose $\delta_2 = 0$. Let $a(a(ab)) = \gamma_0 b + \gamma_1 ab + \gamma_2 a(ab)$.

Set $c' = c + a(ab) - (\gamma_2 - 1)ab - (\gamma_1 + \gamma_2 - 1)b$. Then $ac' = c' + (\gamma_0 + \gamma_1 + \gamma_2 - 1)b$. Therefore (1) $\gamma_2 = 1 - \gamma_0 - \gamma_1$.

Set $c_{\lambda} = c - \lambda \delta_{c} a b - \lambda (\delta_{c} + \gamma_{c} - \lambda \delta_{c} \delta_{1}) b$. Conclude from $(a + \lambda b) c_{\lambda}$ that (2) $\beta_{c} + \gamma_{c} + \delta_{c} = 0$ and (3) $\alpha_{c} + \delta_{c} (\delta_{0} + \delta_{1}) = 0$.

Consider $(b + \lambda c)(a(b + \lambda c))$. Note to avoid c in $\langle a, b + \lambda c \rangle$ the sum of the coefficients of b, ab, and a(ab) in the linear expression of the product under consideration must be zero. This implies (4) $\delta_0 + \delta_1 = 0$. Hence (5) $\alpha_c = 0$ follows from (3).

Set J = F(b - ab) + F(ab - a(ab)). Then (1) implies $aJ \subset J$ and (4) implies b(b - ab) is in J, from which it follows b(ab - a(ab)) = b(ab) - a(b(ab)) is in J. Next, bc is in J because of (2) and (5). The Jacobi identities J(a, b, c) = 0 and J(a, ab, c) = 0 then imply $cJ \subset J$. Thus J is an ideal in L.

Suppose $cd = e + \gamma a + \mu b + j$, c, d, e in C, j in J. The J(a, c, d) = 0 implies e = 0 and $\mu = 0$, while J(b, c, d) = 0 implies $\gamma = 0$. Therefore genus L/J = 0.

(3) Suppose $b(ab) = \delta_0 a + \delta_1 b + \delta_2 ab$ and $a(a(ab)) = \gamma_0 a + \gamma_1 b + \gamma_2 ab + \gamma_3 a(ab)$. Take C any complement of B so that $ac = \alpha_c a$ and $bc = \beta_c a + \gamma_c b + \delta_c(ab)$ for all c in C.

There exists a nonzero λ in F so that $a' = a + \lambda c$, b, a'b, and a'(a'b) are linearly independent because $F \neq GF(2)$. Then $\alpha_c = 0$ because $a'c = \alpha_c a' - \lambda \alpha_c d$, otherwise this case would be reduced to (1) or (2). Repeating this argument for $c' = c + a(ab) - \gamma_3 ab - \gamma_2 b$ from ac' conclude $\gamma_0 = 0$ and $\gamma_1 = 0$.

If $\gamma_c = 0$ replace b by b + c to reduce this case to either (1) or (2). From (b + c)(a(b + c)) it follows that $\delta_1 = 0$. From the coefficient of a(ab) in J(b, ab, c) = 0 conclude $\beta_c = 0$.

Note Fa'b + Fa'(a'b) = Fab + Fa(ab), a' chosen as above. Then from b(a'b) conclude $\delta_0 = 0$.

Set J = Fab + Fa(ab). If dim C = 1 the proof of this case is complete. If not, suppose c and d are linearly independent elements of C, with cd = e + b', e in C, b' in B. Note dim $\langle b + c, a \rangle$ and dim $\langle b + d, a \rangle = 4$. Then in order to avoid a reduction to a previous case conclude from (b + c)d that e is in Fc and from (b + d)c that eis in Fd. Hence e is zero. If $bc = \gamma_c a(ab)$ then $(a(ab))c = \gamma_c a(a(a(ab)))$. Therefore $b(cd) = (bc)d - c(bd) = \gamma_c(a(ab))d - \gamma_d(a(ab))c = 0$. Clearly a(cd) = 0. Thus cd is in the center of B.

Suppose x is a nonzero element of the center of B. Then x =

 $\gamma a + \mu b + \kappa a b + \eta a(ab), \ \eta \neq 0$, otherwise ax = 0 and bx = 0 imply x = 0. From bx = 0 conclude $\delta_2 = 0$ and hence $\gamma = 0$. If cd = x, $\mu \neq 0$ then $(b + c)d = \mu c + b''$, b'' in $\langle b + c, a \rangle$, contradicting genus L = 2. Therefore $L^2 = J$.

II. Let L = B + D, D a subspace, B a subalgebra with genus B=2 and dim B=5. Suppose L has no genus 2 4-dimensional subalgebras. Apply Theorem 2, [2] to conclude (1) for some generator of B, say, b, bd - d is in B for all d in D, and for the remaining generators a and c of B ad and cd are always in B or (2) $BD \subset D$. In (1) a further normalization is desirable Either a, b, and ab or b, c, and cb are linearly independent, say a, b and ab are linearly independent. If cb depends on a, b, ab, c replace b by b + c and obtain with a change in notation that a, b, c, ab, cb span B. Note $B^2 \subset Fa + Fc + Fab + Fcb$. (1) D can be taken so that bd = d for all d in D. Suppose $b(ab) = \gamma_0 a + \gamma_1 ab$. Conclude from $b(d + ab - ab) = \gamma_0 a + \gamma_1 ab$. $(1 - \gamma_1)a)$ that $\gamma_1 = 1 + \gamma_0$. An exactly analogous argument leads to $b(cb) = \gamma'_0 c + (1 + \gamma'_0) cb$. Then $\gamma_0 = \gamma'_0 = \gamma$ because b((a + c)b) must be a linear combination of a + c, b, and ab + cb. Next $a(ab) = \delta_1 ab$ because a(ab) and b(ab) are in Fa + Fab by J(a, b, ab) = 0. Note $(b + \lambda a)(a(b + \lambda a)) = b(ab) + \lambda a(ab) = \gamma a + (1 + \gamma + \lambda \delta_1)ab$. Therefore $\delta_1 = 0$ since there exists λ in F such that a, $b + \lambda a$, c, ab, $c(b + \lambda a)$ are linearly independent. Analogously, c(cb) = 0.

Suppose $ac = \beta a + \gamma c + \delta ab + \mu cb$. For any nonzero λ in $F(a + \lambda d)c$ must be a linear combination of $a + \lambda d$, $ab - \lambda d$, b, c, and bc. Therefore $\alpha = \delta$. Analogously, $\gamma = \mu$. The same argument applies to $c((a + \lambda d)b)$. This implies $c(ab) = \alpha'(a + ab) + \gamma'c + \mu'cb$. Analogously, $a(cb) = \alpha''a + \beta''ab + \gamma''(c + cb)$. Set J = F(a + ab) + F(c + cb). Then $bJ \subset J$. Then J(a, b, c) = 0 implies $\gamma' = \mu'$ and $\alpha'' = \beta''$, i.e. c(ab) and a(cb) are in J. It is immediate that J is an ideal in B.

Suppose $ad = \alpha_1 a + \beta_1 b + \gamma_1 c + \delta_1 ab + \mu_1 cb$ and $cd = \alpha_2 a + \beta_2 b + \gamma_2 c + \delta_2 ab + \mu_2 cb$. For any nonzero λ in F such that $1 - \lambda \gamma \neq 0$ set $d_{\lambda} = d + (\lambda/(1 - \lambda \gamma))\mu_1 c + (\lambda(\delta_1 + \lambda/(1 - \lambda \gamma))\mu_1 \alpha)a$. Then $(b + \lambda a)d_{\lambda} = d_{\lambda} + \lambda\beta_1 b + \lambda(\alpha_1 - \delta_1)a + \lambda(\gamma_1 - \mu_1)c$. Therefore $\gamma_1 = \mu_1$ and $\lambda\beta_1 = \alpha_1 - \delta_1$. Analogously, $\alpha_2 = \delta_2$ and $\lambda\beta_2 = \gamma_2 - \mu_2$. Hence ad and cd are in J with the possible exception F = GF(3), F = GF(2) being excluded by hypothesis. To handle this special case note first (ab)d = ad + (ad)b by J(a, b, d) = 0. Then J(a, ab, d) = 0 can be rewritten as the relation $2(1 + \gamma)ad + (2 + \gamma)(ad)b = b((ab)d)$. Here a b term can occur only in $2(1 + \gamma)ad$. Done, unless $2(1 + \gamma) = 0$. The above relation now simplifies to (ad)b = b((ab)d). Note b(ab) = -a, b(cb) = -c. Calculate both sides of this last relation and find $\alpha_1 = \gamma_1 = \delta_1 = \mu_1 = 0$. Then $\beta_1 = 0$, otherwise genus $\langle a, d, c \rangle > 2$. A similar

argument implies cd is always in J. It follows that $dJ \subset J$. Hence J is an ideal in L.

If $de = f + \alpha a + \beta b + \gamma c + j$, d, e, f in D, j in J, then J(b, d, e) = 0 implies $de = \beta b + j$. If $\beta \neq 0$ then L/J would not be solvable for every term of its derived series would contain (D + Fb)/J. Hence de is in J. Therefore genus L/J = 0.

Suppose a and c chosen so that ac is in Fa + Fc. Then (2)take D so that $bd = \lambda_d b$. If $\lambda_d \neq 0$ this case reduces to last one since there exists λ in F so that genus $\langle b + \lambda d, a, c \rangle = 2$. Suppose bd = 0 for all d in D. If $b(ab) = \delta_0 a + \delta_1 b + \delta_2 ab$ then repeat this argument for b and $d + ab + \delta_2 a$. Conclude $b(ab) = \delta_2 ab$. Likewise b(cb) is in Fcb. Suppose $ad = \alpha a + \beta b + \gamma c + \delta ab + \mu cb$. Repeat the argument for $b + \lambda a$ and $d + \lambda \delta a + \lambda \mu c$ and conclude ad is in J = Fab + Fcb. Similarly, cd is in J. It now follows immediately that ac = 0 from (a + d)c a linear combination of a + d, b, ab, c, cb and a(c + d) a linear combination of a, b, ab, c + d, cb. Thus (ab)c =a(bc) by J(a, b, c) = 0. Clearly genus $\langle a, b, c+d \rangle > 2$ if a(bc) involved c, while genus $\langle a, b, c+d \rangle > 2$ if a(bc) involved c, while genus $\langle a+d, d \rangle < 2$ b, c > 2 if (ab)c involved a. Therefore a(bc) and (ab)c are both in J. J(a, b, ab) = 0 implies $a(ab) = \gamma_1 b + \gamma_2 ab$. Consider a(a(b + c)) to conclude $\gamma_1 = 0$. Likewise J(c, b, cb) = 0 and c(c(b + a)) forces c(cb)to be in J. Suppose $de = f + \alpha a + \beta b + \gamma c + j, d, e, f$ in D, j in J. Consider $(a + \lambda d)e$. To avoid a previous case or genus L > 2 conclude f=0 and $\alpha=0$. It has been shown that ad is in B^2 , hence $(a+\lambda d)e$ is in $\langle a + \lambda d, b, c \rangle^2 = J$. Therefore $\beta = 0$ and $\gamma = 0$. Thus $J = L^2$.

4. The classification of genus 1 and 2 Lie algebras.

PROPOSITION 7. If L is nilpotent then genus $L = \dim L^2$.

Proof. The result follows from the fact that the span of any minimal set of generators intersects the derived algebra in zero.

PROPOSITION 8. Let L be nilpotent with $L^2 = Fz$. Let μ be the alternating bilinear form defined by $xy = \mu(x, y)z$ for all x and y in L. Then the isomorphism class of L is determined by the integers dim L and rank μ .

Proof. Suppose ψ is an isomorphism of L onto $\psi(L)$. Then if $\psi(\mu)$ is defined by $\psi(x)\psi(y) = \psi(\mu)(\psi(x), \psi(y))\psi(z)$ from $\psi(xy) = \psi(\mu(x, y)z) = \mu(x, y)\psi(z)$ it follows that μ and $\psi(\mu)$ are equivalent.

Conversely, suppose that V = W + Fz is a vector space over Fand that μ and $\hat{\mu}$ are two equivalent bilinear forms on $V \times V$, i.e.

JAMES BOND

 $\mu(x, y) = \hat{\mu}(T(x), T(y))$ for a linear transformation T of V. Turn V into an algebra (V, \cdot) over F by $xy = \mu(x, y)z$ and an algebra (V, *) by $T(x)*T(y) = \hat{\mu}(T(x), T(y))T(z)$. Then T is an isomorphism of (V, \cdot) onto (V, *) because $T(xy) = T(\mu(x, y)z) = \mu(x, y)T(z) = \hat{\mu}(T(x), T(y))T(z) = T(x)*T(y)$.

PROPOSITION 9. A nonnilpotent Lie algebra with dim $L^2 = 1$ is the direct sum of an abelian ideal and a 2-dimensional non-abelian ideal.

Proof. Let $L^2 = Fz$. If Lz = 0 then L is nilpotent. Take x so that xz = z. If $xy = \gamma z$ then $x(y - \gamma z) = 0$. Thus the inner derivation D_x decomposes L into 1-dimensional eigenspaces all with eigenvalue 0 except for Fz corresponding to eigenvalue 1. The proposition follows from the fact that the product of two eigenvectors is again an eigenvector.

PROPOSITION 10. Let L be a genus 1 Lie algebra with a 1dimensional ideal J having nonabelian genus 0 quotient. Then

(1) L^2 is abelian and either

(a) $L = L^2 + Fx$, $L^2 = B + Fa + Fz$ with xz = z, xa = a + z, xb = b for all b in B. or

(b) L = A + Fz + Fx with $xz = \lambda z$, $\lambda \neq 1$, az = 0 and xa = a for all a in A. or

(2) L^2 is nilpotent genus one, $L^2 = A + Fz$, $L = L^2 + Fx$, with xz = 2z and xa = a for all a in A.

In (b) x is characterized so that λ determines the isomorphism class of L, while in (2) L^2 determines the isomorphism class of L. The algebras described in (a) and (b) are never isomorphic because D_x has an indecomposable 2-dimensional eigenspace in (a) and not in (b).

Proof. Let L = A + Fx + Fz, where $xz = \lambda z$, $az = \beta_a z$, $xa = a + \alpha_a z$, and $ab = \mu_{a\ b} z$ for all a, b in A. For any a in $A, \beta_a = 0$, because a is in L^2 modulo Fz and therefore the trace of D_a on Fz is zero. From J(x, a, b) = 0 conclude $(2 - \lambda)\mu_{a,b} = 0$

If $\lambda = 1$ some $\alpha_a \neq 0$ otherwise *L* has genus 0. Take *a* so that $\alpha_a = 1$. Then Fa + F(xa - a) is an indecomposable eigenspace for D_x . If $\lambda \neq 1$ replace *a* by $a + (1 - \lambda)^{-1}\alpha_a z$ and find that xa = a.

The algebra (1) (a) is obtained when $\lambda = 1$; (1) (b) when $\lambda \neq 1$ and $\mu_{a,b} = 0$ for all a and b in A, and (2) when $\lambda = 2$ and some $\mu_{a,b} \neq 0$.

The nonminimal genus 1 Lie algebras have been classified. A

606

minimal genus 1 Lie algebra has dimension 3 and the classification of all such algebras is discussed in [4], pages 11 through 14.

The next two propositions furnish the key to the classification of the genus 2 Lie algebras.

PROPOSITION 11. Let M be a commutative associative algebra of 2 by 2 matrices over a field F. Then dim $M \leq 2$. If dim M = 2 then M has an identity.

Proof. Let E_{ij} denote the 2 by 2 matrix with a 1 in the *i*th *j*th spot and zeros elsewhere. The only two distinct E_{ij} to commute are E_{11} and E_{22} . If dim M > 2 no spot can always be zero. Hence M contains $A = \begin{pmatrix} 1\beta \\ 0\gamma \end{pmatrix}$, $B = \begin{pmatrix} 0\beta' \\ 1\gamma' \end{pmatrix}$, and $C = \begin{pmatrix} 0\beta'' \\ 0\gamma'' \end{pmatrix}$. If $\gamma'' \neq 0$ take $\gamma = 0$ and $\gamma' = 0$ and it follows $AB \neq BA$, while if $\lambda'' = 0$, $\beta'' \neq 0$, it follows by taking $\beta = 0$ and $\beta' = 0$ that $BC \neq CB$.

If dim M = 2 then dim $\langle M, I \rangle = 2$, since $\langle M, I \rangle$ is commutative. Hence I is in M.

PROPOSITION 12. Let M be a 2-dimensional commutative associative algebra of 2 by 2 matrices over any field F.

If char $F \neq 2$ then M has a basis I and T with $T^2 = \beta I$, β in F. If char F = 2 then if M has a basis I and T with $T^2 = \delta T + \beta I$, $\delta \neq 0$, then M contains no S with $S^2 = \beta' I$, and furthermore T may be taken so that $\delta = 1$.

If M_i is such an algebra with basis I and T_i , $T_i^2 = \beta_i I$, i = 1, 2. Then M_1 and M_2 are isomorphic if and only if $\beta_1 = \lambda^2 \beta_2$, λ a nonzero element of F. If char F = 2 and $T_i^2 = T_i + \beta_i I$, i = 1, 2, then M_1 and M_2 are isomorphic if and only if there exists μ in F with $\mu^2 + \mu + \beta_1 + \beta_2 = 0$.

Proof. If $\varphi: M_1 \to M_2$ is an onto isomorphism then I and T a basis of M_1 go into I and $\varphi(T)$ a basis for M_2 . Therefore, $T^2 = \delta T + \beta I$ implies $\varphi(T)^2 = \delta \varphi(T) + \beta I$. If also $\varphi(T)^2 = \delta_1 \varphi(T) + \beta_1 I$. Then $(\delta - \delta_1)\varphi(T) = (\beta_1 - \beta)I$ implies $\delta = \delta_1$ and $\beta = \beta_1$. Hence to prove the proposition it is only necessary to find a canonical choice for T within M.

Suppose $T^2 = \delta T + \beta I$. Set $S = \lambda T + \mu I$, $\lambda \neq 0$. Then $S^2 = (\lambda \delta + 2\mu)S + (\beta \lambda^2 - \mu \lambda \delta - \mu^2)I$.

If char F = 2 and $\delta = 0$ then β can be adjusted by squares as in the char $F \neq 2$ case. If $\delta \neq 0$ no allowable choice of λ and μ gives a zero coefficient of S; hence choose $\lambda = \delta^{-1}$. Then the coefficient of I becomes $\mu^2 + \mu + \beta \delta^{-2}$. Here if I and T_1 and I and T_2 were

JAMES BOND

basis for M with δ_1 and δ_2 nonzero then there would exist μ_1 and μ_2 such that $\mu_1^2 + \mu_1 + \beta_1 = \mu_2^2 + \mu_2 + \beta_2$, i.e. $\mu^2 + \mu + \beta_1 + \beta_2 = 0$ for $\mu = \mu_1 + \mu_2$.

A nilpotent algebra A is said to have index n if $A^n \neq 0$ while $A^{n+1} = 0$.

A genus 2 index 2 nilpotent Lie algebra can only be decomposed into two nonzero genus 1 Lie algebras. The only decomposition of a genus 1 Lie algebra is the decomposition into its center and an indecomposable genus 1 nilpotent algebra.

The classification and isomorphism of problems for genus 1 nilpotent Lie algebras are both solved by considering the bilinear form introduced in Proposition 8. A generalization of this approach to the genus 2 index 2 nilpotent Lie algebras leads to a solution of the isomorphism problem over some fields, but never to a solution of the classification problem. This point is clarified by introducing the definition: $L = L_1^* + L_2$, L_1 and L_2 subalgebras of L, $L_1 \bigcap L_2 \subset L^2$ and $[L_1, L_2] = 0$. If $L = L^* + L_2$ implies $L \leq L^2$ or $L_2 \leq L^2$ then L is said to be (*) indecomposable. The only (*) indecomposable genus 1 nilpotent Lie algebras are Fx with x in $L - L^2$ and a 3-dimensional genus 1 nilpotent Lie algebra (L = Fx + Fy + Fz with [x, y] = z). The number of 3-dimensional summands in L is counted by one-half the rank of μ (as introduced in Proposition 8). The situation for the genus 2 index 2 case appears to be quite different. We are unable to describe the (*) indecomposable Lie algebras in this case.

Let $\mu_1 + L^2, \dots, \mu_n + L^2$ be a basis for L/L^2 . If $L^3 = 0$ then the multiplication in L can be specified by an n by n skew symmetric matrix $M = (u_{ij}), u_{ij}$ in L^2 and defined by $[\mu_i, \mu_j] = u_{ij}$. A change in basis for L/L^2 leads to the replacing of M by DMD^t , D a nonsingular matrix with entries in F.

Ordinary decomposition of M is equivalent to (*) decomposition of L. The classification problem for the genus 2 index 2 nilpotent Lie algebras reduces to finding a canonical form for the matrix M. An



FIGURE 1

infinite number of reasonable candidates for (*) indecomposable algebras are described by matrices of Figure 1 with $Fu + Fv = L^2$, $k \ge 1$, $r \ge 1$.

We have not found a calculable property of M invariant under $M \rightarrow DMD^{t}$, D a nonsingular matrix with elements in F, which would distinguish indecomposable from decomposable M.

The classical theory of regular pencils of skew symmetric matrices leads to a partial solution of the isomorphism problem for genus 2 index 2 nilpotent Lie algebras. We associate to M the regular pencil of symmetric matrices $A = A + \lambda B$, where $M = v_1 A + v_2 B$ and the notation and terminology is that of Gantmacher [3]. Theorem 6 and its corollary, p. 91 and p. 92, vol. II [3], when applicable to F, give a criterion in terms of the elementary divisors of Λ . A change in the choice of a basis v_1 , v_2 of L^2 is equivalent to substituting $\alpha x + \beta y$ for x and $\gamma x + \delta y$ for y, with $\alpha \delta - \beta \gamma \neq 0$, $\alpha, \beta, \gamma, \delta$ in F. Once the elementary divisors associated to L by writing M = $v_1A + v_2B$ and the elementary divisors associated to L' by writing $M' = v'_1A + v'_2B$ have been calculated, the isomorphism question for L reduces to calculating whether there exists a substitution replacing x by x + y and y by x + y carrying the elementary divisors associated with L into those associated with L'.

PROPOSITION 13. Let L be a nilpotent genus 2 Lie algebra over any field F. Then either the index of L is 2 or $L = C + Fa + Fb + L^2$, with $L^2 = Fu + Fv$, C + Fu a genus 1 nilpotent subalgebra, $C(Fa + Fb + L^2) = 0$, au = 0, av = u, ba = v, bu = 0, and bv = 0. The isomorphism class of L is determined by dim L and that of C + Fu.

Proof. Let $L^2 = Fu + Fv$, which is possible by Proposition 7. Let $M\left\{\begin{pmatrix} \alpha\beta\\\gamma\delta \end{pmatrix}, xu = \alpha u + \gamma v, xv = \beta u + \delta v$ for some x in $L\right\}$. Then M is a commutative associative algebra which must consist of nilpotent transformations and hence dim $M \leq 1$ (Proposition 12). Therefore M = (0) and index L = 2 or $M = F\begin{pmatrix} 01\\00 \end{pmatrix}$ for u, v a suitably chosen basis of L^2 .

If index L > 2 there is an element a of L determined up to L^2 by requiring au = 0 and av = u. Note if $bL^2 = 0$ and $cL^2 = 0$ then J(a, b, c) = 0 implies bc is in Fu. Therefore there must exist b' in L such that $b'L^2 = 0$ and $b'a = v + \lambda u$, otherwise v would not be in L^2 . Set b = b'' + v and find ba = v. Let C be a complement of $L^2 + Fa + Fb$ in L such that $C(L^2 + Fa) = 0$. Note b is also determined up to C and C is determined up to L^2 , and hence the isomorphism class of L is determined by rank μ , where μ is defined as in the statement of Proposition 8.

The following situation arises a number of times in the propositions to follow. Therefore it seems best to determine the suitable invariants once and for all.

Let A be an n-dimensional subspace of a Lie algebra L over a field F such that $A^2 \subset Fz$ for some z in L. Suppose A = B + Fa. Let $\mu: A \times A \to F$ denote the alternating bilinear from defined by $cd = \mu(c, d)z$ for all c and d in A. Choose any basis of B and any complementary vector of B in A and associate to μ a matrix C in the usual way. It is easy to see that a canonical form for C is:



FIGURE 2

where $K = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. Let is (B, A) denote the triplet of integers (r, s, n). Note the integer r is the rank of μ restricted to B, while s is dim $\{d \text{ in the radical of } \mu \text{ restricted to } B \mid ad \neq 0\}$.

If $A = C \dotplus Fa \dotplus Fb$ and the allowable basis for A must be of the form a basis of C and elements of the form a + c, b + d, c and d in C then the canonical form of μ becomes:



FIGURE 3

In this case let is (C, a, b) = (r, s, t, n).

610

PROPOSITION 14. Let L be a nonnilpotent genus 2 Lie algebra with a 2-dimensional abelian derived algebra. Let ρ denote the adjoint representation of L on L². Then either (1) $\rho(L)$ contains a nonsingular transformation and the kernel of ρ is the center of L, with the isomorphism class of L determined by the isomorphism class of the associative algebra $\rho(L)$, or (2) $\rho(L) \cong F\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$. Here L =B + Fa + Fu + Fv, au = u, av = 0, bu = 0, bv = 0 for all b in B, and B + Fa + Fv is nilpotent genus 1. Here is(B, a) determines the isomorphism class of L.

Proof. (1) Let $\rho(a)$ be a nonsingular. Let $B = \{b - \rho(a)^{-1}ab \mid b \text{ in } L\}$. Then $\rho(B) = 0$. From J(a, b, c) = 0, if either $\rho(b) = 0$ or $\rho(c) = 0$ it follows that bc = 0.

(2) Let au = u, av = 0. Note if ab' = u + v then ab = v for b = b' - u. Hence B can be taken so that L = B + Fa + Fu + Fv, with B(Fu + Fv) = 0 and $aB \subset Fv$. From J(a, b, c) = 0 conclude bc in Fv for b and c in B. Thus B + Fa + Fv is a nilpotent genus 1 algebra. It cannot be abelian for then v would not be in L^2 . Note regardless of the choice of C with B = C + D as in the statement of the proposition that a can be taken so that ac = 0, while dim $\{d \text{ in } D \mid ad \neq 0\}$ is independent of the choice of a.

PROPOSITION 15. Let L be a genus 2 Lie algebra with a 2dimensional nonabelian ideal J with genus 0 quotient. Then there exists an ideal A, which is non-abelian genus 0, such that $L = A \oplus J$.

Proof. Let L = B + J, B a subspace of L. Suppose uv = v, u and v a basis for J. Since the derived algebra of an ideal is an ideal $bu = \alpha u + \gamma v$ and $bv = \beta v$, b in B. Then J(b, u, v) = 0 implies $\alpha = 0$. Set $b' = b - \gamma v + \beta u$ and find that b'u = 0 and b'v = 0. Let A be spanned by the b', b in B. Suppose ab = c + k, c in A, k in J. Then J(u, a, b) = 0 and J(v, a, b) = 0 imply ab = c. Hence $L = A \oplus J$, A genus 0. If A were abelian then genus L = 0.

LEMMA 3. Let L be a genus 2 Lie algebra over a field F with a 2-dimensional abelian ideal J having a non-abelian genus 0 quotient. Let ρ denote the adjoint representation of L on J. If dim $\rho(L) > 1$ then J has a basis so that

(i)
$$\rho(a) = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$
 and $\rho(x) = \begin{pmatrix} \alpha & 0 \\ 0 & \alpha + 1 \end{pmatrix}$, or
(ii) Char $F = 2$, (a) $\rho(a) = \begin{pmatrix} 0 & \lambda \\ 1 & 0 \end{pmatrix}$, $\lambda \neq 0$, $\rho(x) = \begin{pmatrix} \alpha & 0 \\ 0 & \alpha + 1 \end{pmatrix}$
(b) $\rho(a) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $\rho(x) = \begin{pmatrix} 0 & \alpha \\ 1 & 1 \end{pmatrix}$.

Proof. Let L = A + Fx + J, with xa - a and ab in J for all a and b in A. Then (*): $\rho(x)\rho(a) - \rho(a)\rho(x) = \rho(a)$, for all a in A. dim $\rho(A) = 2$ implies some $\rho(a) = I$, contradicting (*).

Suppose $\rho(a) \neq 0$ and its eigenvalues are not in F. Then there exists a basis of J such that $\rho(a) = \begin{pmatrix} 0 & \lambda \\ 1 & 0 \end{pmatrix}$, since trace $\rho(a) = 0$ because of (*). If $\rho(a)$ has zero as its only eigenvalue then there exists a basis of J such that $\rho(a) = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$. In either case it may be supposed that $\rho(x) = \begin{pmatrix} \alpha & \beta \\ 0 & \delta \end{pmatrix}$. Then (*) implies (i) or (ii) a.

If $\rho(a)$ has nonzero eigenvalues in F, a may be taken so that 1 and -1 are its eigenvalues. Choose a basis for J so that $\rho(a) = \begin{pmatrix} 1 & \varepsilon \\ 0 & -1 \end{pmatrix}$, $\varepsilon = 0$ or 1, $\varepsilon = 1$ only if char F = 2. Take $\rho(x) = \begin{pmatrix} 0 & \beta \\ \alpha & \delta \end{pmatrix}$. Then (*) implies ii (b).

PROPOSITION 16. A non-minimal genus 2 Lie algebra over a field F, with char $F \neq 2$, has a nilpotent derived algebra.

Proof. If J and L/J are abelian then the desired result is clear. If J is abelian and L/J is non-abelian genus 0 the result follows immediately from Lemma 3. If J is non-abelian and L/J is genus 0 the result is clear from Proposition 15.

The solution of the problem of extending an abelian 2-dimensional algebra J by a non-abelian genus 0 algebra is quite complicated. In the next proposition, accordingly, the description of the different algebras which arise is less explicit than those given in Proposition 10. Nevertheless, it is not difficult to derive a similarly explicit list of Lie algebras from the information given in the next proposition.

PROPOSITION 17. Let L be a genus 2 Lie algebra over a field F having a 2-dimensional abelian ideal J with non-abelian genus 0 quotient. Suppose L^2 is nilpotent. Then

(1) $L=J \dotplus A \dotplus Fx$, with xa=a for all a in A. Here $(A \dotplus J)^{\circ} \subset K \subset J$, where $(\rho(x) - 2I)K = 0$. If AJ = 0, to avoid genus L < 2 it is necessary that $\rho(x) \neq \begin{pmatrix} 1 & 0 \\ 0 & \lambda \end{pmatrix}$, λ in F, for any basis of J. The isomorphism class of L is determined when AJ = 0 by the equivalence class of $\rho(x)$ and the isomorphism class of the nilpotent subalgebra $A \dotplus J$, which as of now is undetermined, and when $\rho(a) \neq 0$ by is (ker ρ, a).

(2) L = J + B + Fa + Fx, with xb = b for all b in B, xa - aa nonzero element of J. If (B + Fa)J = 0 then either $\rho(x) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and J + B + Fa is abelian or $\rho(x) = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}$ and J + B + Fa is nilpotent: with isomorphism class determined by is (B, a). If $(B + Fa)J \neq 0$ then J = Fu + Fv, $xu = \lambda u$, xv = v, $\lambda \neq 1$, xa = a + v, with either $B = \ker \rho$, λ and is(B, a) determining the isomorphism class of L, or $\ker \rho = C + Fa$ with bu = v, bv = 0, and is(C, a, b) determining the isomorphism class, or bu = 0, bv = u, $\lambda \neq 2$ $ab = 1/(2 - \lambda)u$, with aC = 0 and bC = 0.

(3) L = J + C + Fa + Fb + Fx, with xc = c for all c in C, $xa - a \ xb - b$ linearly independent elements of J. Here $\rho(x) = I$, AJ = 0, and C + Fa + Fb is abelian.

Proof. (1) Suppose x and A chosen so that L = J + A + Fx with xa = a for all a in A. From J(x, a, b) = 0 it follows that x(ab) = 2ab. Hence $(A + J)^2 \subset K \subset J$ with $(\rho(x) - 2J)K = 0$.

If $\rho(x) = I$ then genus L = 0 and if $\rho(x) = \begin{pmatrix} 1 & 0 \\ 0 & \lambda \end{pmatrix}$, $\lambda \neq 1$ with respect to basis u, v of J then Fv has genus 0 quotient and hence genus L = 1.

If dim $\rho(c) = 1$ then choose basis $\{b_i\}$ for A and basis u, v for J and write $b_i b_j = \alpha_{ij} u + \beta_{ij} v$ with is $((\alpha_{ij}), (\beta_{ij}))$ determining multiplication in A + J.

If dim $\rho(c) = 2$, the hypothesis of L^2 nilpotent implies the structure of case (3) Lemma 3. Thus au = v, av = 0 and $xu = \alpha u$. $xv = (\alpha + 1)v$. Thus dim $K \leq 1$ and the invariant $is(\ker \rho, a)$ determines the multiplication in A + J.

(2) Suppose x and A chosen so that L = J + B + Fa + Fx with xb = b for all b in B and xa - a a nonzero element of J, and x and B + Fa cannot be taken so that xa = a, xb = b for all b in B.

If $D_x - I$ were non-singular and xa = a + k', k' in J, set a' = a + k, where $(D_x - I)k = k'$, and find xa' = a'. It follows that $D_x - I$ is nonzero and singular. Thus $\rho(x) - I = \begin{pmatrix} \lambda & 0 \\ 0 & 0 \end{pmatrix}$, $\lambda \neq 0$, or $\rho(x) - I = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$.

If (B + Fa)J = 0 then J(x, a, b) = 0 implies $(B + Fa + J)^2 \subset K \subset J$, K as in (1).

Suppose $(B + Fa)J \neq 0$ then by Lemma 3, (i) $\rho(x)$ has two distinct eigenvalues, hence also $\rho(x) - I$, and therefore $\rho(x) = \begin{pmatrix} \lambda & 0 \\ 0 & 1 \end{pmatrix}$.

If $B = \ker \rho$ then au = v, av = 0 or au = 0, av = u. In either case J(x, a, b) = 0 implies ab in K, K as in (1).

Suppose $B \neq \ker \rho$ then since a can be modified by multiples of b and dim (B + Fa)J = 1, by Lemma 3 a can be taken so that aJ = 0and then normalized so that $xa = a + \beta'v$, $\beta' \neq 0$. Let $bJ \neq 0$ and $C + Fa = \ker \rho$. Then J(x, a, b) = 0 implies $x(ab) = 2ab + \beta'bv$. Either bu = v, bv = 0 and is(C, a, b) determines the multiplication of B + Fa + J or bu = 0 and bv = u. In the latter case let $ab = \alpha u + \beta v$ and conclude $\beta = 0$ and $\alpha(\lambda - 2) + \beta' = 0$. Hence $\lambda \neq 2$ and it follows that aC = 0 and bC = 0 from J(x, a, c) = 0 and J(x, b, c) = 0 for c in C.

(3) Suppose L = J + C + Fa + Fb + Fx, with xc = c for all cin C, xa - a, xb - b linearly independent elements of J, and it is possible to choose A and x so that xa = a for all elements of A. Then $\rho(x) = I$, for $\rho(x) - I$ nonsingular was ruled out in the last case, while if $\rho(x) - I$ has rank 1, then the argument there employed can be used to show that every a can be chosen so that xa - a is in Fv, v any fixed element of J not in the range of $\rho(x)-I$. Lemma 3, (i) gives (C + Fa + Fb)J = 0 and J(x, c, d) = 0, for any c, d in C + Fa + Fb, then implies cd = 0.

PROPOSITION 18. Let L be a genus 2 Lie algebra over a field F having a 2-dimensional abelian ideal J with non-abelian genus 0 quotient. Suppose L^2 is not nilpotent. Then char F = 2, $L = L^2 + K + Fx$, dim $\rho(L) = 2$ and either $L^2 = J + A$ with xa = a for all a in A or $L^2 = J + \ker \rho + Fa$, xb = b for all b in ker ρ , xa = a a nonzero element of J. In every case $(\ker \rho + Fa)^2$ is contained in a 1-dimensional subspace of J so that the isomorphism class of $\rho(L)$ and is $(\ker \rho, a)$ determine the isomorphism class of L.

Proof. If L^2 is not nilpotent then dim $\rho(L) = 2$. Then char F = 2 and $L^2 \subset J$ by Lemma 3. But, A modulo J is in L^2 , hence $L = L^2 + Fx$.

Suppose $L = J \dotplus A \dotplus Fx$, xa = a for all a in A. Here x(ab) = 0and $\rho(x)$ as in Lemma 3, (ii) cannot have two identical eigenvalues. Therefore it must have rank 1, which excludes (ii) b of Lemma 3. If ker $\rho = B$ then $(B \dotplus Fa \dotplus J)^2$ is contained in a 1-dimensional subspace of J and is (B, a) is the appropriate invariant. If ker $\rho \neq B$ then we can take a in ker ρ , since $\rho(x) - I$ has rank 1, $\alpha = 0, 1$ in (ii) a of Lemma 3. Here J(x, a, b) = 0 leads to a contradiction.

PROPOSITION 19. Let L be a solvable minimal genus 2 Lie algebra without a 2-dimensional ideal with genus 0 quotient. Then L belongs to the same isomorphism class as one of the following algebras:

(1) $L = L^2 + Fx$, L^2 a 3-dimensional abelian ideal without a 2dimensional invariant subspace for D_x . The equivalence class of D_x determines the isomorphism class of L.

(2) $L = L^2 \dotplus Fx$, L^2 3-dimensional nilpotent with $(L^2)^2 = Fx \neq 0$ and D_x irreducible on L^2/Fz . Here the equivalence class of D_x determines the isomorphism class of L.

(3) $L = L^2 \dotplus Fx$, L^2 4-dimensional abelian, with the characteristic polynomial of D_x on L of the form $p(y) = y(q(y))^2$, q(y) irreducible.

Two such algebras are isomorphic if and only if the corresponding p(y) and p'(y) are projectively equivalent, i.e. there exist nonzero element λ of F such that $p(\lambda y) = p'(y)$.

Proof. (1, 2) dim L = 4 and $L = L^2 + Fx$. If L^2 is abelian then (1). If not let $L^2 = Fa + Fb + Fab$. If $(L^2)^2$ were 2-dimensional then its quotient would be genus 0. Hence $(L^2)^2 = Fab$ is a 1-dimensional ideal and therefore annihilated by L^2 . The multiplication table of Lis determined by D_x acting on L^2/Fab since x is specified only up to the addition of elements of L^2 . This gives (2).

(3) dim L = 5. L does not have a 1-dimensional ideal J. Otherwise, genus L/J = 2 implies J complemented by a genus 2 subalgebra by Proposition 2; genus L/J = 0 implies genus L = 1; genus L/J = 1 implies L/J has a 1-dimensional ideal with genus 0 quotient (hence L has a 2-dimensional ideal with genus 0 quotient).

Suppose L has a maximal abelian ideal J of dimension 2. Then genus L/J = 1. Let x and y be preimages of two generators of L/Junder the natural map L onto L/J. Then x and y generate a 3dimensional subalgebra S. If dim $S^2 = 1$ take x and y so that x(xy) =0; if dim $S^2 = 2$ with x in S^2 then J(x, y, xy) = 0 implies x(xy) = 0. It follows (x(x(y + a)) = x(xa) = 0 for all a in I to avoid $\langle x, y + a \rangle$ containing a nonzero element of I and thus having dimension ≥ 4 . If $D_x = 0$ then $[D_x, D_y] = 0$ and $I + S^2$ is an abelian ideal. Therefore there exist a and b spanning I with xa = b and xb = 0. Take y so that $ya = \gamma a$. Then $(xy)a = x(ya) + (xa)y = \gamma b + by$ and therefore (x + a)((x + a)y) = -by = 0 to avoid dim $\langle x + a, y \rangle \geq 4$. Therefore Fb is an ideal in L, a contradiction.

Suppose L has a maximal abelian ideal J of dimension 3. Note dim $L^2 \ge 3$, for genus L = 2 implies dim $L^2 \ge 2$ and if dim $L^2 = 2$ then genus $L/L^2 = 0$. If $J \ne L^2$ then dim $(J \cap L^2) = 2$ to avoid $J \cap L^2$ a 1-dimensional ideal. Write $L = J \cap L^2 \dotplus Fx \dotplus Fy \dashv Fz$, with x in L^2 , y in J, and conclude from $xz = \alpha x + a$ and $yz = \beta y + b$ with a and b in $J \cap L^2$ that genus $L/(J \cap L^2) = 0$. If $J = L^2$ let $L = L^2 \dotplus$ $Fx \dotplus Fy$. Then $xy \ne 0$ implies Fxy is a 1-dimensional ideal because x(xy) and y(xy) are in $L^2 \cap (Fx \dotplus Fy \dotplus Fxy) = Fxy$. Therefore xy =0 for all x and y in L^2 , i.e. (x + a)y = 0 and x(y + a) = 0 for all a in L^2 , a contradiction. If $J \subsetneq L^2$ let $L = J \dotplus Fx \dotplus Fy$ with x in L^2 and y chosen so that xy = x + a, a in J, which is possible because x is in L^2 . If $a \ne 0$ then from x(xy) = xa and y(xy) = x + a + yain L^2 and $Fx \dotplus Fy \dotplus Fxy$ conclude xa and ya are in Fa so that Fa is an ideal. It follows from x(y + b) and (x + b)y that xb = 0 and yb = 0 for all b in J, a contradiction.

Suppose L has a maximal abelian ideal J of dimension 4. Then

JAMES BOND

there exist and b in J such that x, xa, xb, a, and b span L. To avoid a 2-generated 4-dimensional subalgebra $x(xa) = \alpha a + \beta xa$ and $x(xb) = \alpha'b + \beta'xb$ with $\alpha = \alpha'$ and $\beta = \beta'$ necessary for x(x(a + b))to be in F(a + b) + Fx(a + b). The characteristic polynomial of D_x on L^2 is $(y^2 - \beta y + \alpha)^2$. If $y^2 - \beta y + \alpha$ were not irreducible L would contain a 2-dimensional ideal with genus 0 quotient.

References

1. James Bond, The structure of Lie algebras with large minimal generating sets, Math. Ph. D. thesis, University of Notre Dame, Notre Dame, Indiana, June 1964.

2. _____, Weak minimal generating set reduction theorems for associative and Lie algebras, Illinois J. Math, 10, No. 4, (1966), 579-591.

3. F. R. Gantmacher, The Theory of Matrices, Vol. 2, Chelsea Publishing Company, 1959.

4. N. Jacobson, Lie algebras, Interscience Publishers, 1962.

5. M. S. Knebelman, Classification of Lie algebras, Ann. of Math., 36 (1935), 46-56.

6. G. Leger, A particular class of Lie algebras, Proc. Amer. Math. Soc., 16 (1965), 293-296.

7. E. I. Marshall, The genus of a perfect Lie algebra, J. London Math. Soc., 40 (1965), 276-282.

8. E. M. Patterson, Genertors of Linear algebras, Proc. London Math. Soc., (3) 7 (1957), 467-480.

9. E. W., Wallace, Complex four dimensional Lie algebras, Proc. Royal Soc. Edinburg, Sec A, 65 (1958), 72-83.

Received January 23, 1967, and in revised form October 19, 1970,

THE PENNSYLVANIA STATE UNIVERSITY AND PRESENTLY AT NUC, SAN DIEGO

616

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON Stanford University Stanford, California 94305

C. R. HOBBY

University of Washington Seattle, Washington 98105 J. DUGUNDJI Department of Mathematics University of Southern California Los Angeles, California 90007

RICHARD ARENS University of California Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH B. H. NEUMANN F. WOLF K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA CALIFORNIA INSTITUTE OF TECHNOLOGY UNIVERSITY OF CALIFORNIA MONTANA STATE UNIVERSITY UNIVERSITY OF NEVADA NEW MEXICO STATE UNIVERSITY OREGON STATE UNIVERSITY UNIVERSITY OF OREGON OSAKA UNIVERSITY UNIVERSITY OF SOUTHERN CALIFORNIA STANFORD UNIVERSITY UNIVERSITY OF TOKYO UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF WASHINGTON

AMERICAN MATHEMATICAL SOCIETY CHEVRON RESEARCH CORPORATION NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The Pacific Journal of Mathematics is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics Vol. 37, No. 3 March, 1971

Mohammad Shafqat Ali and Marvin David Marcus, <i>On the degree of the minimal polynomial of a commutator operator</i>	561
Howard Anton and William J. Pervin, <i>Integration on topological</i> semifields	567
Martin Bartelt, Multipliers and operator algebras on bounded analytic	
functions	575
Donald Earl Bennett, Aposyndetic properties of unicoherent continua	585
James W. Bond, Lie algebras of genus one and genus two	591
Mario Borelli, The cohomology of divisorial varieties	617
Carlos R. Borges, <i>How to recognize homeomorphisms and isometries</i>	625
J. C. Breckenridge, Burkill-Cesari integrals of quasi additive interval functions	635
J. Csima, A class of counterexamples on permanents	655
Carl Hanson Fitzgerald, <i>Conformal mappings onto</i> ω -swirly domains	657
Newcomb Greenleaf. Analytic sheaves on Klein surfaces	671
G. Goss and Giovanni Viglino. <i>C-compact and functionally compact</i>	
<i>spaces</i>	677
Charles Lemuel Hagopian, Arcwise connectivity of semi-aposyndetic plane	
continua	683
John Harris and Olga Higgins, <i>Prime generators with parabolic limits</i>	687
David Michael Henry, Stratifiable spaces, semi-stratifiable spaces, and their	
relation through mappings	697
Raymond D. Holmes, <i>On contractive semigroups of mappings</i>	701
Joseph Edmund Kist and P. H. Maserick, <i>BV-functions on semilattices</i>	711
Shûichirô Maeda, On point-free parallelism and Wilcox lattices	725
Gary L. Musser, <i>Linear semiprime</i> (p; q) radicals	749
William Charles Nemitz and Thomas Paul Whaley, <i>Varieties of implicative</i>	750
	739
Jaroslav Nesetrii, A congruence theorem for asymmetric trees	771
Robert Anthony Nowlan, A study of H-spaces via left translations	779
Gert Kjærgaard Pedersen, Atomic and diffuse functionals on a C*-algebra	795
Tilak Raj Prabhakar, On the other set of the biorthogonal polynomials suggested by the Laguerre polynomials	801
Leland Edward Rogers, <i>Mutually aposyndetic products of chainable</i> continua	805
Frederick Stern, An estimate for Wiener integrals connected with squared	
error in a Fourier series approximation	813
Leonard Paul Sternbach, On k-shrinking and k-boundedly complete basic	
sequences and quasi-reflexive spaces	817
Pak-Ken Wong, <i>Modular annihilator A*-algebras</i>	825