

Pacific Journal of Mathematics

**ARCWISE CONNECTIVITY OF SEMI-APOSYNDETTIC PLANE
CONTINUA**

CHARLES LEMUEL HAGOPIAN

ARCWISE CONNECTIVITY OF SEMI-APOSYNDETTIC PLANE CONTINUA

CHARLES L. HAGOPIAN

Suppose M is a bounded semi-aposyndetic plane continuum and for any positive real number ε there are at most a finite number of complementary domains of M of diameter greater than ε . In this paper it is proved that M is arcwise connected.

Let M be a continuum (a closed connected point set) and let x and y be distinct points of M . If M contains a continuum H and an open set G such that $x \in G \subset H \subset M - \{y\}$, then M is said to be *aposyndetic* at x with respect to y [4]. M is said to be *semi-aposyndetic* if for each pair of distinct points x and y of M , M is aposyndetic either at x with respect to y or at y with respect to x . In [3] it is proved that every bounded semi-aposyndetic plane continuum which does not have infinitely many complementary domains is arcwise connected. For other results concerning semi-aposyndetic plane continua see [1] and [2].

Let x and y be distinct points of a metric space S . A finite collection $\{A_1, A_2, \dots, A_m\}$ of sets in S is a *chain* in S from x to y provided A_1 contains x , A_m contains y , and for i and j belonging to $\{1, 2, \dots, m\}$, $A_i \cap A_j \neq \emptyset$ if and only if $|i - j| \leq 1$. If each element of a chain \mathcal{A} has diameter less than r (a positive real number) then \mathcal{A} is said to be an *r -chain*. Suppose $\mathcal{A} = \{A_1, A_2, \dots, A_m\}$ and $\mathcal{B} = \{B_1, B_2, \dots, B_n\}$ are chains in S from x to y . The chain \mathcal{B} is said to *run straight through* \mathcal{A} provided the closure of each element of \mathcal{B} is contained in an element of \mathcal{A} and if B_i and B_k ($1 \leq i \leq k \leq n$) both lie in an element A_s of \mathcal{A} , then for each integer j ($i < j < k$), B_j is contained in an element of \mathcal{A} whose intersection with A_s is nonvoid.

If M is a bounded plane continuum and for any positive real number ε there are at most a finite number of complementary domains of M of diameter greater than ε , then M is said to be an *E -continuum* [6, p. 112].

The boundary of a set A is denoted by $\text{Bd } A$.

THEOREM 1. *Suppose M is a semi-aposyndetic E -continuum in S (a 2-sphere with metric φ), U is a disk in S , x and y are distinct points which belong to the same component of $M \cap U$, and V is an open disk in S containing U . Then for any positive real number r less than both $\varphi(x, y)/5$ and $\varphi(\text{Bd } U, \text{Bd } V)/5$ there exists an r -chain $\{H_1, H_2, \dots, H_n\}$ ($n > 3$) in S from x to y such that for each positive*

integer i less than or equal n , H_i is a continuum in $M \cap V$ and $\varphi(H_i, \text{Bd } V)$ is greater than $4r$.

Proof. Let G be the union of all components of $S - M$ which have diameter less than $r/3$. Since M is a semi-aposyndetic E -continuum, $M \cup G$ is a semi-aposyndetic continuum which does not have infinitely many complementary domains [5, Th. 2 (proof)]. Let F be the x -component of $U \cap (M \cup G)$. F is a semi-aposyndetic continuum in S which does not have infinitely many complementary domains [3, Th. 1] (D and M in [3] are $S - U$ and $M \cup G$ respectively). Hence F is arcwise connected [3, Th. 2]. Let A be an arc in F from x to y . There exists a finite point set B in $A - \{x, y\}$ such that each component of $A - B$ has diameter less than $r/3$. For each component C of $A - B$, let $G(C)$ be C union all components of G which intersect C and let $Z(C)$ be the boundary (relative to S) of $G(C)$. For each component C of $A - B$, since the boundary of each component of G is a continuum [6, Th. 2.1, p. 105] and each point of C that is not in G belongs to $Z(C)$, $Z(C)$ is a continuum of diameter less than r in M . Let \mathcal{H} be the finite coherent collection of continua $\{Z(C) \mid C \text{ is a component of } A - B\}$. The points x and y each belong to an element of \mathcal{H} and each element of \mathcal{H} intersects U . It follows that any chain from x to y whose elements are members of \mathcal{H} has the specified conditions.

THEOREM 2. *If M is a semi-aposyndetic E -continuum, then M is arcwise connected.*

Proof. Let S be a 2-sphere which contains M and let φ be a distance function on S . Let p and q be distinct points of M . Define r_1 to be a positive real number less than both $1/8$ and $\varphi(p, q)/5$ and let $s_1 = 4r_1$. According to Theorem 1, there exists an r_1 -chain $\{H_1^1, H_2^1, \dots, H_{n_1}^1\}$ ($n_1 > 3$) in S from p to q such that for each positive integer i less than or equal n_1 , H_i^1 is a continuum in M . Let m_1 be the smallest integer greater than or equal to $(n_1 - 1)/2$. There exist a set of disks $\{U_1^1, U_2^1, \dots, U_{m_1}^1\}$ and a set of open disks $\{V_1^1, V_2^1, \dots, V_{m_1}^1\}$ such that $\{V_1^1, V_2^1, \dots, V_{m_1}^1\}$ is an s_1 -chain in S from p to q and for each positive i less than or equal m_1 , $H_{2i-1}^1 \cup H_{2i}^1 \cup H_{2i+1}^1 \subset U_i^1 \subset V_i^1$ (if n_1 is even, let $H_{n_1+1}^1 = \phi$).

Let $\{p_1^1, p_2^1, \dots, p_{m_1+1}^1\}$ be a point set such that $p_1^1 = p$, $p_{m_1+1}^1 = q$, and for each positive integer i less than or equal m_1 , p_i^1 belongs to H_{2i-1}^1 . Let t_1 be the smallest number in the set $\{\varphi(\text{Bd } U_i^1, \text{Bd } V_i^1) \mid i \leq m_1\} \cup \{\varphi(p_i^1, p_{i+1}^1) \mid i \leq m_1\}$. Let r_2 be a positive real number less than both $t_1/5$ and $1/16$. Define s^2 to be $4r_2$. For each positive in-

teger i less than or equal m_1 , there exists an r_2 -chain \mathcal{C}_i in S from p_1^i to p_{i+1}^i such that each element of \mathcal{C}_i is a continuum in $M \cap V_i^i$ and at a distance greater than $4r_2$ from $\text{Bd } V_i^i$ (Theorem 1). There exists an r_2 -chain $\{H_1^2, H_2^2, \dots, H_{n_2}^2\}$ in S from p to q whose elements belong to $\bigcup_{i=1}^{m_1} \mathcal{C}_i$ such that for each positive integer i less than or equal m_1 , $\mathcal{C}_i \cap \{H_1^2, H_2^2, \dots, H_{n_2}^2\}$ is a coherent collection. Let m_2 be the smallest integer greater than or equal to $(n_2 - 1)/2$. There exist a set of disks $\{U_1^2, U_2^2, \dots, U_{m_2}^2\}$ and a set of open disks $\{V_1^2, V_2^2, \dots, V_{m_2}^2\}$ such that $\{V_1^2, V_2^2, \dots, V_{m_2}^2\}$ is an s_2 -chain in S from p to q and for each positive integer i less than or equal m_2 , $H_{2i-1}^2 \cup H_{2i}^2 \cup H_{2i+1}^2 \subset U_i^2 \subset V_i^2$ (if n_2 is even, let $H_{n_2+1}^2 = \emptyset$). Note that $\{V_1^2, V_2^2, \dots, V_{m_2}^2\}$ runs straight through $\{V_1^1, V_2^1, \dots, V_{m_1}^1\}$.

Continue this process. For $i = 3, 4, 5, \dots$, there exists a chain $\{H_1^i, H_2^i, \dots, H_{n_i}^i\}$ in S from p to q whose elements are continua in M , and there exists an s_i -chain $\{V_1^i, V_2^i, \dots, V_{m_i}^i\}$ ($s_i < 1/2^i$) in S from p to q whose elements are open disks in S such that $\bigcup_{j=1}^{m_i} V_j^i$ contains $\bigcup_{j=1}^{n_i} H_j^i$ and $\{V_1^i, V_2^i, \dots, V_{m_i}^i\}$ runs straight through $\{V_1^{i-1}, V_2^{i-1}, \dots, V_{m_{i-1}}^{i-1}\}$. For each positive integer i , let L_i be the continuum $\bigcup_{j=1}^{m_i} H_j^i$. The limiting set L of the sequence L_1, L_2, L_3, \dots is a continuum in M containing p and q . Note that for each positive integer i , L is contained in $\bigcup_{j=1}^{m_i} V_j^i$.

Let x be a point of $L - \{p, q\}$. For each positive integer i , let $V_{j_i}^i$ be an element of $\{V_1^i, V_2^i, \dots, V_{m_i}^i\}$ which contains x . Assume without loss of generality that $4 < j_1 < m_1 - 4$. For each positive integer i , let P_i be $\{V_1^i, V_2^i, \dots, V_{j_i-4}^i\}$ and let F_i be $\{V_{j_i+4}^i, V_{j_i+5}^i, \dots, V_{m_i}^i\}$. Let $P = \bigcup_{i=1}^{\infty} (P_i \cap L)$ and $F = \bigcup_{i=1}^{\infty} (F_i \cap L)$. P and F are nonempty disjoint relatively open subsets of L and $P \cup F = L - \{x\}$. Hence x is a separating point of L . It follows that L has only two nonseparating points. Therefore L is an arc [6, Th. 6.2, p. 54]. Hence M is arcwise connected.

REMARK. Using [3, Th. 1] and Theorem 2 one can easily prove that if M is a semi-aposyndetic E -continuum, then M has Jones's cyclic property (that is, if p and q are distinct points of M and no point cuts p from q in M , then there exists a simple closed curve lying in M which contains p and q).

REFERENCES

1. C.L. Hagopian, *Arc-wise connectedness of semi-aposyndetic plane continua*, to appear in Trans. Amer. Math. Soc., **158** (1971).
2. ———, *An arc theorem for plane continua*, soon to appear in Ill. J. of Math.
3. ———, *A class of arcwise connected continua*, soon to appear in Proc. Amer. Math. Soc.

4. F. B. Jones, *Aposyndetic continua and certain boundary problems*, Amer. J. Math., **63** (1941), 545-553.
5. ———, *A characterization of a semi-locally-connected plane continuum*, Bull. Amer. Math. Soc., **53** (1947), 170-175.
6. G. T. Whyburn, *Analytic topology*, Amer. Math. Soc. Colloquium Publications, vol. 28, Rhode Island, 1963 (Revised edition).

Received December 4, 1970 and in revised form April 28, 1971.

SACRAMENTO STATE COLLEGE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics

Vol. 37, No. 3

March, 1971

Mohammad Shafqat Ali and Marvin David Marcus, <i>On the degree of the minimal polynomial of a commutator operator</i>	561
Howard Anton and William J. Pervin, <i>Integration on topological semifields</i>	567
Martin Bartelt, <i>Multipliers and operator algebras on bounded analytic functions</i>	575
Donald Earl Bennett, <i>Aposyndetic properties of unicoherent continua</i>	585
James W. Bond, <i>Lie algebras of genus one and genus two</i>	591
Mario Borelli, <i>The cohomology of divisorial varieties</i>	617
Carlos R. Borges, <i>How to recognize homeomorphisms and isometries</i>	625
J. C. Breckenridge, <i>Burkill-Cesari integrals of quasi additive interval functions</i>	635
J. Csima, <i>A class of counterexamples on permanents</i>	655
Carl Hanson Fitzgerald, <i>Conformal mappings onto ω-swirly domains</i>	657
Newcomb Greenleaf, <i>Analytic sheaves on Klein surfaces</i>	671
G. Goss and Giovanni Viglino, <i>C-compact and functionally compact spaces</i>	677
Charles Lemuel Hagopian, <i>Arcwise connectivity of semi-aposyndetic plane continua</i>	683
John Harris and Olga Higgins, <i>Prime generators with parabolic limits</i>	687
David Michael Henry, <i>Stratifiable spaces, semi-stratifiable spaces, and their relation through mappings</i>	697
Raymond D. Holmes, <i>On contractive semigroups of mappings</i>	701
Joseph Edmund Kist and P. H. Maserick, <i>BV-functions on semilattices</i>	711
Shūichirō Maeda, <i>On point-free parallelism and Wilcox lattices</i>	725
Gary L. Musser, <i>Linear semiprime $(p; q)$ radicals</i>	749
William Charles Nemitz and Thomas Paul Whaley, <i>Varieties of implicative semilattices</i>	759
Jaroslav Nešetřil, <i>A congruence theorem for asymmetric trees</i>	771
Robert Anthony Nowlan, <i>A study of H-spaces via left translations</i>	779
Gert Kjærgaard Pedersen, <i>Atomic and diffuse functionals on a C^*-algebra</i>	795
Tilak Raj Prabhakar, <i>On the other set of the biorthogonal polynomials suggested by the Laguerre polynomials</i>	801
Leland Edward Rogers, <i>Mutually aposyndetic products of chainable continua</i>	805
Frederick Stern, <i>An estimate for Wiener integrals connected with squared error in a Fourier series approximation</i>	813
Leonard Paul Sternbach, <i>On k-shrinking and k-boundedly complete basic sequences and quasi-reflexive spaces</i>	817
Pak-Ken Wong, <i>Modular annihilator A^*-algebras</i>	825