

# Pacific Journal of Mathematics

**LINEAR SEMIPRIME  $(p; q)$  RADICALS**

GARY L. MUSSER

## LINEAR SEMIPRIME $(p; q)$ RADICALS

GARY L. MUSSER

**This paper introduces McKnight's  $(p; q)$ -regularity and  $(p; q)$  radicals, a collection of radicals which contains the Jacobson radical and the radicals of regularity and strong regularity among its members. The linear semiprime  $(p; q)$  radicals are classified canonically and, as a result of this classification, these radicals can be distinguished by the fields  $GF(p)$  and are shown to form a lattice. The semiprime  $(p; q)$  radicals are found to be hereditary and the linear semiprime  $(p; q)$  radical of a complete matrix ring of a ring  $R$  is determined to be the complete matrix ring over the  $(p; q)$  radical of  $R$ . More generally, the  $(p; q)$  radical of a complete matrix ring over  $R$  is contained in the matrix ring over the  $(p; q)$  radical of  $R$  for all  $(p; q)$  radicals.**

A function  $\rho$  which assigns to each ring  $R$  an ideal  $\rho R$  of the ring is called a *radical function* in the sense of Amitsur and Kurosh [1; 5] if it has the following properties:

R1: If  $\phi: R \rightarrow S$  is a ring epimorphism and  $\rho R = R$ , then  $\rho S = S$ .

R2:  $\rho(\rho R) = \rho R$  for all rings  $R$  and if  $\rho I = I$  for any ideal  $I$  of  $R$ , then  $I \subseteq \rho R$ .

R3:  $\rho(R/\rho R) = 0$  for all rings  $R$ .

If  $\rho$  is a radical function, then the ideal  $\rho R$  is called the *radical* of  $R$ . If  $\rho R = R$  for some ring  $R$ , then  $R$  is called a  $\rho$ -*radical ring* while if  $\rho R = 0$  we call  $R$  a  $\rho$ -*semisimple ring*. If  $I$  is an ideal (right ideal) of a ring  $R$ , then  $I$  is called a  $\rho$ -*radical ideal* (right ideal) if  $I$  is a  $\rho$ -radical ring.

Now let  $p(x)$  and  $q(x)$  be polynomials with integer coefficients. An element  $r$  of a ring  $R$  is called  $(p; q)$ -*regular* if  $r \in p(r)Rq(r)$ , that is,  $r = p(r)sq(r)$  for some  $s \in R$  where an integer multiple of a ring element has its usual meaning. If every element of an ideal  $I$  of  $R$  is  $(p; q)$ -regular, that is, if  $r \in p(r)Iq(r)$  for all  $r \in I$ , then  $I$  is said to be a  $(p; q)$ -*regular ideal*. Examples of  $(p; q)$ -regularity are quasi-regularity,  $(x + 1; 1)$  [4], von Neumann regularity,  $(x; x)$  [7] and strong regularity,  $(x^2; 1)$  [2].

LEMMA 1. *If  $I$  and  $R/I$  are  $(p; q)$ -regular, then  $R$  is  $(p; q)$ -regular.*

*Proof.* Let  $r \in R$ . Then  $r + I \in R/I$ , which implies

$$r + I = p(r + I)(s + I)q(r + I) = p(r)sq(r) + I$$

for some  $s + I \in R/I$ . Thus  $r - p(r)sq(r) \in I$  and, since  $I$  is  $(p; q)$ -

regular,  $r - p(r)sq(r) = p[r - p(r)sq(r)]tq[r - p(r)sq(r)]$  for some  $t \in I$ . Moreover there exist  $u, v \in R$  such that

$$\begin{aligned} r - p(r)sq(r) &= p[r - p(r)sq(r)]tq[r - p(r)sq(r)] \\ &= [p(r) - p(r)u]t[q(r) - vq(r)] \end{aligned}$$

or  $r = p(r)(s + t - ut - tv + utv)q(r)$ . Therefore  $R$  is  $(p; q)$ -regular.

**LEMMA 2.** *If  $I$  and  $J$  are  $(p; q)$ -regular ideals of  $R$ , then  $I + J$  is a  $(p; q)$ -regular ideal of  $R$ .*

*Proof.* Immediate from Lemma 1, since the homomorphic image of a  $(p; q)$ -regular ring is a  $(p; q)$ -regular ring.

**COROLLARY 1.** *The sum of all  $(p; q)$ -regular ideals of a ring  $R$  is a  $(p; q)$ -regular ideal of  $R$ .*

*Proof.* This follows from Lemma 2, since  $(p; q)$ -regularity is defined elementwise.

We shall let  $(p(x)Rq(x))$  denote the largest  $(p; q)$ -regular ideal of the ring  $R$ . Then we have

**THEOREM 1.** (J. D. McKnight, Jr.) *If a function  $\rho$  is defined by  $\rho R = (p(x)Rq(x))$  for all rings  $R$ , then  $\rho$  is a radical function.*

*Proof.* We only need to show R3 holds. Let  $I/\rho R$  be a  $(p; q)$ -regular ideal of  $\rho(R/\rho R)$ . Then by Lemma 1,  $I$  is a  $(p; q)$ -regular ideal of  $R$  and  $I \subseteq \rho R$ .

We shall call  $(p(x)Rq(x))$  the  $(p; q)$  radical of the ring  $R$ . Thus the Jacobson radical and the radicals of regularity and strong regularity of  $R$  are given by  $((x + 1)R)$ ,  $(xRx)$  and  $(x^2R)$  respectively.

1. A canonical representation for linear semiprime  $(p; q)$  radicals. A radical function  $\rho$  is called *semiprime* if  $\rho R$  is a semiprime ideal, equivalently, if  $\rho R$  contains the prime (Baer-lower) radical [6; 3]. Now we shall determine the form of the semiprime  $(p; q)$  radicals.

**LEMMA 3.**  *$\rho$  is a semiprime radical function if and only if  $\rho R = R$  for all zero rings  $R$ .*

*Proof.* The necessity is clear. Now if  $I^2 \subseteq \rho R$  for some ideal  $I$  of  $R$ , then  $\rho[(I + \rho R)/\rho R] = (I + \rho R)/\rho R$  since  $(I + \rho R)/\rho R$  is isomorphic to the zero ring  $I/(I \cap \rho R)$ . Also  $\rho(R/\rho R) = 0$  implies

$$\rho[(I + \rho R)/\rho R] = 0$$

and therefore  $I \subseteq \rho R$ .

**THEOREM 2.** (A. H. Ortiz)  $(p(x)Rq(x))$  is semiprime for all rings  $R$  if and only if the constant terms of  $p(x)$  and  $q(x)$  are 1 or  $-1$ .

*Proof.* Let  $p(x)$  and  $q(x)$  have constant terms 1 or  $-1$  and  $R$  be any zero ring. Then for  $r \in R$ , we have  $r = p(r)(\pm r)q(r)$  and  $R \subseteq (p(x)Rq(x))$ . Thus  $R = (p(x)Rq(x))$ . Conversely, if  $a_0$  and  $b_0$  are the constant terms of  $p(x)$  and  $q(x)$  respectively, then suppose  $a_0 \neq \pm 1$  or  $b_0 \neq \pm 1$ . Since we are assuming the  $(p; q)$  radical is semiprime, it follows from Lemma 3 that for the zero ring with additive group  $Z/(a_0b_0)$  we have  $(p(x)[Z/(a_0b_0)]q(x)) = Z/(a_0b_0)$  where  $Z$  denotes the ring of integers and  $(a_0b_0)$  the ideal generated by  $a_0b_0$ . However if  $r \in (p(x)[Z/(a_0b_0)]q(x))$ , then  $r \in p(r)[Z/(a_0b_0)]q(r)$  and  $r = 0$ . Hence  $Z/(a_0b_0) = 0$ , which is a contradiction.

Henceforth we shall be considering semiprime  $(p; q)$  radicals and, since  $(p(x)R) = (p(-x)R) = (-p(x)R)$ , we shall assume that the constant term of  $p(x)$ , similarly the constant term of  $q(x)$ , is 1.

**LEMMA 4.** If the constant term of  $p(x)$  is 1, then for all  $r \in R$  we have  $r \in p(r)R$  if and only if  $R = p(r)R$ .

*Proof.* The sufficiency is obvious. Now let  $r \in p(r)R$ . Since  $p(r) = rf(r) + 1$  for some integral polynomial  $f(x)$ , for any  $s \in R$  we have,  $p(r)s = rf(r)s + s$ . Since  $r \in p(r)R$  we have  $s \in p(r)R$  and  $R \subseteq p(r)R$ . Therefore,  $R = p(r)R$ .

**COROLLARY 2.** If the constant terms of  $p(x)$  and  $q(x)$  are both 1, then for all  $r \in R$  we have  $r \in p(r)Rq(r)$  if and only if  $R = p(r)Rq(r)$ .

**THEOREM 3.** If  $(p(x)Rq(x))$  and  $(p'(x)Rq'(x))$  are semiprime for all rings  $R$ , then  $(p(x)Rq(x)) \cap (p'(x)Rq'(x)) = (p(x)p'(x)Rq'(x)q(x))$ .

*Proof.* Clearly  $(p(x)p'(x)Rq'(x)q(x)) \subseteq (p(x)Rq(x)) \cap (p'(x)Rq'(x))$ . Now let  $r \in (p(x)Rq(x)) \cap (p'(x)Rq'(x))$ . Then  $r \in (p'(x)Rq'(x))$  implies  $r \in p'(r)Rq'(r)$  and, by Corollary 2,  $R = p'(r)Rq'(r)$ . Now  $r \in p(r)Rq(r)$  and  $R = p'(r)Rq'(r)$  implies  $r \in p(r)p'(r)Rq'(r)q(r)$ . The product polynomials  $p(x)p'(x)$  and  $q(x)q'(x)$  have constant terms 1, hence  $r = p(r)p'(r)sq'(r)q(r)$  implies that  $s \in (p(x)Rq(x)) \cap (p'(x)Rq'(x))$ . Therefore  $(p(x)Rq(x)) \cap (p'(x)Rq'(x))$  is  $(pp'; q'q)$ -regular and

$$(p(x)Rq(x)) \cap (p'(x)Rq'(x)) \subseteq (p(x)p'(x)Rq'(x)q(x)) .$$

In what follows we shall determine a canonical representation for all *linear semiprime*  $(p; q)$  radicals, that is,  $(p; q)$  radicals determined by integral polynomials  $p(x)$  and  $q(x)$  which are products of linear polynomials having constant term 1.

LEMMA 5.  $((ax + 1)(bx + 1)R) \subseteq ((a + b)x + 1]R)$  for all integers  $a, b$ .

*Proof.* Let  $r \in ((ax + 1)(bx + 1)R)$ . Then  $r = (ar + 1)s$  for  

$$s \in ((ax + 1)(bx + 1)R) .$$

Thus  $r = (ar + 1)(bs + 1)t = (br + ar + 1)t = ((a + b)r + 1)t$ , where  $t \in ((ax + 1)(bx + 1)R)$ , implies that

$$((ax + 1)(bx + 1)R) \subseteq ((a + b)x + 1]R .$$

COROLLARY 3.  $((ax + 1)R) \subseteq ((\max + 1)R)$  for all integers  $m$ .

*Proof.* By Theorem 3 we have  $((ax + 1)^m R) = ((ax + 1)R)$ .

COROLLARY 4.  $((ax + 1)R) \subseteq ((a^k x + 1)R)$  for  $k = 1, 2, 3, \dots$ .

LEMMA 6.  $((ax + 1)(bx + 1)R) \subseteq ((ma + nb)x + 1]R)$  for all integers  $m, n$ .

*Proof.* This is immediate from Corollary 3, Lemma 5 and Theorem 3.

Now Corollary 3, Lemma 6 and Theorem 3 yield

THEOREM 4.  $((ax + 1)(bx + 1)R) = ((a, b)x + 1]R)$  where  $(a, b)$  is the greatest common divisor of  $a$  and  $b$ .

We shall now show that the converse of Corollary 4 is true.

LEMMA 7.  $((a^k x + 1)R) \subseteq ((ax + 1)R)$  for  $k = 1, 2, 3, \dots$ .

*Proof.* We first show that  $((a^2 x + 1)R) \subseteq ((ax + 1)R)$ . For this inclusion it is sufficient to show that  $((a^2 x + 1)R) = 0$  whenever  $((ax + 1)R) = 0$  so suppose  $((ax + 1)S) = 0$  for some ring  $S$ . Then if  $r \in ((a^2 x + 1)S)$  we have  $r = (a^2 r + 1)s$  or  $ar = (a(ar) + 1)as$ . Thus  $a((a^2 x + 1)S) \subseteq ((ax + 1)S)$  and  $ar = 0$  for all  $r \in ((a^2 x + 1)S)$ . Therefore  $r = (a^2 r + 1)s = (ar + 1)s$  implies that  $((a^2 x + 1)S) \subseteq ((ax + 1)S) = 0$ . The result now follows by induction.

Combining Corollary 4 and Lemma 7 we have

**THEOREM 5.**  $((ax + 1)R) = ((a^kx + 1)R)$  for  $k = 1, 2, 3, \dots$ .

Our next lemma and Theorem 3 permit us to represent each linear semiprime  $(p; q)$  radical as a  $(pq; 1)$  radical.

**LEMMA 8.**  $((ax + 1)R) = (R(ax + 1))$ .

*Proof.* First, for  $r, s \in R$  define a circle product by  $r \circ s = r + s + ars$ . Then  $(r \circ s) \circ t = r \circ (s \circ t)$ . Now if  $r \in ((ax + 1)R)$ , then  $r \circ s = 0$  for some  $s \in ((ax + 1)R)$ . Since  $s \circ t = 0$  for some  $t \in ((ax + 1)R)$ , we have that  $r = t$  and  $s \circ r = 0$ . Therefore,  $((ax + 1)R) \subseteq (R(ax + 1))$ . A similar argument yields the opposite inclusion, hence equality.

We can now give a canonical representation for all linear semiprime  $(p; q)$  radicals.

**THEOREM 6.** *Every linear semiprime  $(p; q)$  radical can be uniquely represented by a radical of the form  $((ax + 1)R)$  where the nonnegative integer  $a$  is a finite product of distinct prime factors.*

*Proof.* Theorem 3 and Lemma 8 show that

$$(p(x)Rq(x)) = (p(x)q(x)R)$$

for the linear semiprime radical  $(p(x)Rq(x))$ . Then Theorems 3, 4 and 5 show that  $(p(x)q(x)R) = ((ax + 1)R)$  for some nonnegative integer  $a$  where  $a$  is a finite product of distinct prime factors.

To distinguish between the linear semiprime radicals observe that if  $a = \pi_{i=1}^n p_i$  for primes  $p_i$ , then  $((ax + 1)R) = R$  for  $R = GF(p_i)$ ,  $i = 1, 2, \dots, n$  and  $((ax + 1)R) = 0$  for  $R = GF(p)$  for all primes  $p \neq p_i, i = 1, 2, \dots, n$ .

**2. The lattice of linear semiprime  $(p; q)$  radicals.** Let  $(p; q)$  denote the radical function defined by  $(p; q)(R) = (p(x)Rq(x))$  for all rings  $R$ . We partially order the linear semiprime  $(p; q)$  radical functions by defining  $(ax + 1; 1) \leq (bx + 1; 1)$  if  $((ax + 1)R) \subseteq ((bx + 1)R)$  for all rings  $R$ . Then we have

**THEOREM 7.** *The collection of all linear semiprime  $(p; q)$  radicals form a lattice with respect to the partial order  $\leq$  where the infimum and supremum are given by the canonical representatives:*

$$(i) \quad (ax + 1; 1) \wedge (bx + 1; 1) = ((a, b)x + 1; 1)$$

$$(ii) \quad (ax + 1; 1) \vee (bx + 1; 1) = ([a, b]x + 1; 1)$$

where  $[a, b]$  denotes the least common multiple of  $a$  and  $b$ .

*Proof.* (i) By Corollary 3 we have  $((a, b)x + 1; 1) \leq (ax + 1; 1)$ ,  $(bx + 1; 1)$ . Now if  $(cx + 1; 1) \leq (ax + 1; 1)$ ,  $(bx + 1; 1)$ , then  $((cx + 1)R) \subseteq ((ax + 1)R) \cap ((bx + 1)R) = ((ax + 1)(bx + 1)R) = ([a, b]x + 1)R$ .

(ii) First let  $a$  and  $b$  be relatively prime. Since  $(abx + 1; 1)$  is clearly an upper bound of  $(ax + 1; 1)$  and  $(bx + 1; 1)$ , we show that for all rings  $R$ ,  $((abx + 1)R) \subseteq ((cx + 1)R)$  for any other upper bound  $(cx + 1; 1)$ . Again it is enough to show that this inclusion holds for any ring  $S$  for which  $((cx + 1)S) = 0$ . As in the proof of Lemma 7,  $a((abx + 1)S) \subseteq ((bx + 1)S) \subseteq ((cx + 1)S) = 0$  and similarly  $b((abx + 1)S) = 0$ . Therefore, since  $(a, b) = 1$ , for all  $r \in ((abx + 1)S)$  we have integers  $m, n$  such that  $r = m(ar) + n(br) = 0$ . Therefore  $((abx + 1)S) = 0$  and  $((abx + 1)R) \subseteq ((cx + 1)R)$ . Thus when  $(a, b) = 1$ , we have  $(ax + 1; 1) \vee (bx + 1; 1) = ([a, b]x + 1; 1)$ . Using this result it is easy to see that the statement is true for arbitrary integers  $a$  and  $b$ .

It is interesting to observe that  $((x + 1; 1)$ , the Jacobson radical, is the least element in this lattice.

**3. Hereditary  $(p; q)$  radicals.** A radical function  $\rho$  is called *hereditary* if every ideal of a  $\rho$ -radical ring is  $\rho$ -radical. Equivalently, if for any (associative) ring  $R$  and any ideal  $I$  of  $R$  we have the equation  $\rho I = I \cap \rho R$ , then  $\rho$  is hereditary [3, p. 125]. The linear semiprime  $(p; q)$  radical functions are hereditary. Moreover we have

**THEOREM 8.** *If  $(p; q)$  is semiprime, then it is hereditary.*

*Proof.* Let  $I$  be an ideal of  $R$ . For any radical function  $\rho$  we have  $\rho I \subseteq I \cap \rho R$  [3, p. 125]. Now  $r \in I \cap (p(x)Rq(x))$  implies  $r \in I$  and  $r = p(r)sq(r)$  for some  $s \in (p(x)Rq(x))$ . Since the constant terms of  $p(x)$  and  $q(x)$  are 1 we have  $s \in I$  and  $I \cap (p(x)Rq(x)) \subseteq (p(x)Iq(x))$ .

It is easy to see that if the polynomial  $p(x)q(x)$  has  $x^2$  as a factor, then  $(p; q)$  is also hereditary. Thus the radicals of von Neumann regularity and strong regularity are hereditary. The radical given by  $(xR)$  is not hereditary for if  $R$  is the ring of integers modulo 4 and  $I$  is the ideal  $\{0, 2\}$ , then  $(xI) = 0$  while  $I \cap (xR) = I \cap R = I$ .

**4.  $(p; q)$  radicals of matrix rings.** Let  $R_n$  denote the ring of all  $n \times n$  matrices whose elements are taken from the ring  $R$ . We shall show that for all  $(p; q)$  radicals and all rings  $R$  the inclusion

$(p(x)R_nq(x)) \subseteq (p(x)Rq(x))_n$  holds while for linear semiprime  $(p; q)$  radicals we have equality. D. M. Morris has shown

**LEMMA 9.** *If  $p(x) = \pm 1$  or  $p(x) = \pm x$ , then  $(p(x)\mathbf{Z}) = \mathbf{Z}$ ; otherwise  $(p(x)\mathbf{Z}) = 0$*

*Proof.* Clearly  $(p(x)\mathbf{Z}) = \mathbf{Z}$  when  $p(x) = \pm 1$  or  $p(x) = \pm x$ . Suppose that  $(p(x)\mathbf{Z}) \neq 0$  and that  $p(x) \neq \pm 1$ . Then  $(p(x)\mathbf{Z}) = (r)$  where  $(r)$  is the ideal generated by some positive integer  $r$ . Let  $m$  be any prime not dividing  $r$ . Since  $mr \in (p(x)\mathbf{Z})$  we have  $mr = p(mr)m'r$  for some  $m' \in \mathbf{Z}$ . Since  $p(x) \neq \pm 1$  we must have  $p(mr) = \pm m$  for infinitely many primes  $m$ . It follows that  $p(x) = \pm x$ .

**COROLLARY 5.**  $(1\mathbf{Z}1) = (x\mathbf{Z}) = (\mathbf{Z}x) = \mathbf{Z}$  and  $(p(x)\mathbf{Z}q(x)) = 0$  for all other choices of  $p(x)$  and  $q(x)$ .

*Proof.* Clearly  $(x\mathbf{Z}x) = 0$  and since  $(p(x)\mathbf{Z}q(x)) \subseteq (p(x)\mathbf{Z})$ , the corollary is established.

**LEMMA 10.** *Let  $\rho$  be any radical function such that  $\rho\mathbf{Z} = 0$ . Then any ring  $R$  can be embedded in a ring  $S$  with unity such that  $\rho R = \rho S$ .*

*Proof.* Let  $\phi$  be the usual embedding of a ring  $R$  into the ring  $S$  with unity and identify  $R$  with  $\phi R$ , [6, p. 8]. Then  $S/R \cong \mathbf{Z}$ , which implies that  $\rho(S/R) = 0$ . Therefore  $\rho S \subseteq R$  and  $\rho S \subseteq \rho R$ . But since  $R$  is an ideal of  $S$  we always have  $\rho R \subseteq \rho S$ , [3, p. 124]. Therefore  $\rho R = \rho S$ .

**LEMMA 11.** *Let  $\rho$  be any radical function satisfying (i)  $\rho\mathbf{Z} = 0$  and (ii) if  $S$  is a ring with unity, then  $\rho(S_n) \subseteq (\rho S)_n$ . Then  $\rho(R_n) \subseteq (\rho R)_n$  for all rings  $R$ .*

*Proof.* By Lemma 10 we can embed  $R$  as an ideal in a ring  $S$  with unity such that  $\rho R = \rho S$ . Therefore we have  $\rho(R_n) \subseteq \rho(S_n) \subseteq (\rho S)_n = (\rho R)_n$ .

**THEOREM 9.**  $(p(x)R_nq(x)) \subseteq (p(x)(Rq(x))_n)$  for all  $(p; q)$  radicals.

*Proof.* For the  $(1; 1)$  radical equality is obvious. Now consider all other  $(p; q)$  radicals except for the  $(x; 1)$  and  $(1; x)$  radicals. By Corollary 5,  $(p(x)\mathbf{Z}q(x)) = 0$ . If  $S$  has unity, then  $(p(x)S_nq(x)) = I_n$  for some ideal  $I$  of  $S$  [6]. If  $r \in I$ , then  $rE_{11} \in I_n$  and

$$rE_{11} = p(rE_{11})Mq(rE_{11}) = p(r)m_{11}q(r)E_{11}$$



where  $M \in I_n$ ,  $m_{11} \in M$  and  $E_{11}$  is the  $n \times n$  matrix  $|e_{ij}|$  where  $e_{11} = 1$ ,  $e_{ij} = 0$  otherwise. Therefore  $r = p(r)mq(r)$ , for  $m \in I$ , which implies that  $I \subseteq (p(x)Sq(x))$  and  $I_n = (p(x)S_nq(x)) \subseteq (p(x)Sq(x))_n$ . Now Lemma 11 yields  $(p(x)R_nq(x)) \subseteq (p(x)Rq(x))_n$  for all  $(p; q)$  radicals except the  $(x; 1)$  and  $(1; x)$  radicals. To show  $(xR_n) \subseteq (xR)_n$ , let  $A \in (xR_n)$ . Then there exists a  $B \in (xR_n)$  such that  $A = AB$ , where  $A = |a_{ij}|$  and  $B = |b_{ij}|$ . Let  $A_1$  denote the product matrix  $AC$  of  $(xR_n)$  where  $C = |c_{ij}|$ ,  $c_{11} = b_{11}$ ,  $c_{ij} = 0$  for  $j > 1$ , that is,  $A_1 = |a'_{ij}|$  where  $a'_{11} = a_{11}$  and  $a'_{ij} = 0$  for  $j > 1$ .  $A_1 \in (xR_n)$  implies that  $A_1 = A_1D$  or  $a'_{i1} = a'_{i1}d_{11}$  where  $d_{11} \in D$ ,  $D \in (xR_n)$ . Again, there is a matrix  $D_1 \in (xR_n)$ ,  $D_1 = |d'_{ij}|$ , where  $d'_{i1} = d_{i1}$  and  $d'_{ij} = 0$  for  $j > 1$ . Therefore  $D_1 = D_1F$  for  $F \in (xR_n)$  and  $d_{11} = d_{11}f_{11}$  where  $f_{11}$  is an element of  $F$ . Now for  $G = |g_{ij}|$  where  $g_{11} = d_{11}$  and  $g_{ij} = 0$  otherwise, we have  $G \in (xR_n)$  because  $G = GF$ . If we let  $J = \{r \in R \mid |r_{ij}| \in (xR_n), r_{11} = r, r_{ij} = 0 \text{ otherwise}\}$ , then  $J$  is an ideal of  $R$ . It follows that for all  $r \in J$  there exists an  $s \in J$  such that  $r = rs$ . Therefore  $J \subseteq (xR)$  and  $d_{11} \in (xR)$ . But  $d_{11} \in (xR)$  implies that  $a_{i1} \in (xR)$  for  $i = 1, 2, \dots, n$ . Similarly,  $a_{ij} \in (xR)$  for

$$i = 1, 2, \dots, n, j = 2, 3, \dots, n.$$

Thus  $A \in (xR)_n$  and  $(xR_n) \subseteq (xR)_n$ . Similarly  $(R_nx) \subseteq (Rx)_n$ .

R. L. Snider gave the following example to show that the inclusion  $\rho(R_n) \subseteq \rho(R)_n$  is not true for all radicals. Let  $\sigma R$  be the upper radical determined by declaring  $GF(2)$  to be semisimple (In [3, p. 6] let  $M = \{GF(2)\}$ ). Then since the ring of  $2 \times 2$  matrices over  $GF(2)$  cannot be mapped homomorphically onto  $GF(2)$ ,  $(GF(2))_2$  is not semisimple.

Finally we show that for all linear semiprime  $(p; q)$  radicals the opposite inclusion holds; hence we have equality.

LEMMA 12. *The sum of two  $(ax + 1; 1)$ -regular right ideals of the ring  $R$  is an  $(ax + 1; 1)$ -regular right ideal of  $R$ .*

*Proof.* Let  $I$  and  $J$  be  $(ax + 1; 1)$ -regular right ideals of  $R$  and  $r \in I, s \in J$ . Then there exists an  $r' \in I$  such that  $r = (ar + 1)r'$ . Now  $s - asr' \in J$ , which implies that there exists an  $s' \in J$  such that  $s - asr' = (a(s - asr') + 1)s'$ . It is easy to see that  $r + s = (a(r + s) + 1)(r' - ar's' + s')$ , hence  $I + J$  is an  $(ax + 1; 1)$ -regular right ideal.

COROLLARY 7. *The sum of all  $(ax + 1; 1)$ -regular right ideals of a ring  $R$  is an  $(ax + 1; 1)$ -regular right ideal of  $R$ .*

LEMMA 13. *The sum  $K$  of all  $(ax + 1; 1)$ -regular right ideals of a ring  $R$  is a two-sided ideal of  $R$ . Therefore  $K \subseteq ((ax + 1)R)$ .*

*Proof.* [cf. 3, p. 93] Let  $s \in K$  and  $r \in R$ . Then  $sr \in K$  implies that  $sr = (asr + 1)s'$  for some  $s' \in K$ . It is easy to see that  $rs = (ars + 1)(-ars's + rs)$ , hence  $rs$  is  $(ax + 1; 1)$ -regular. For  $m \in \mathbf{Z}$ ,  $t \in R$  we have  $sm + st \in K$ . Since from above  $r(sm + st)$  must be  $(ax + 1; 1)$ -regular,  $rs\mathbf{Z} + rsR$ , the right ideal generated by  $rs$ , is an  $(ax + 1; 1)$ -regular right ideal and we have  $rs\mathbf{Z} + rsR \subseteq K$ , therefore  $rs \in K$ . Since  $K$  is now an  $(ax + 1; 1)$ -regular ideal,  $K \subseteq ((ax + 1)R)$ .

**THEOREM 10.** *If  $(p; q)$  is a linear semiprime radical, then for all rings  $R$ ,  $(p(x)R_nq(x)) = (p(x)Rq(x))_n$ .*

*Proof.* [cf. 4, p. 11] We only need to show that  $((ax + 1)R)_n \subseteq ((ax + 1)R_n)$  for all positive integers  $a$ . Let  $k$  be a fixed positive integer,  $k \leq n$ , and  $|r_{ij}| \in ((ax + 1)R_n)$  where  $r_{ij} = 0$  for  $i \neq k$ . Then by Lemma 4,  $r_{kk} \in (ar_{kk} + 1)R$  implies that  $R = (ar_{kk} + 1)R$ . Therefore for each  $r_{kj}$  there exists an  $s_{kj}$  such that  $r_{kj} = (ar_{kk} + 1)s_{kj}$  for  $j = 1, 2, \dots, n$ . Thus  $|r_{ij}| = (a|r_{ij}| + 1)|s_{ij}|$  where  $s_{ij} = 0$  for  $i \neq k$ . Hence the right ideal  $P_k$  of  $n \times n$  matrices having elements of  $((ax + 1)R)$  in the  $k$ th row and zeros elsewhere is an  $(ax + 1; 1)$ -regular right ideal, thus  $P_k \subseteq ((ax + 1)R_n)$ . Since  $((ax + 1)R)_n$  is the sum of the  $P_k$ ,  $k = 1, 2, \dots, n$ , we have  $((ax + 1)R)_n$ .

If  $R$  is a field we have  $0 = (x^2R_n) \subseteq (x^2R)_n = R_n$ , therefore the radical of strong regularity shows that we cannot have the matrix equality for all  $(p; q)$  radicals.

REFERENCES

1. S. A. Amitsur, *Radicals in rings and bicategories*, Amer. J. Math., **76** (1954), 100-125.
2. R. Arens and I. Kaplansky, *Topological representations of algebras*, Trans. Amer. Math. Soc., **63** (1948), 457-481.
3. N. Divinsky, *Rings and Radicals*, University of Toronto Press, 1965.
4. N. Jacobson, *Structure of Rings*, Amer. Math. Soc. Colloq. Pub. Vol. **37**, Providence, 1964.
5. A. G. Kurosh, *Radicals of rings and algebras*, Mat. Sb., **33** (1953), 13-26.
6. N. McCoy, *The Theory of Rings*, MacMillan, New York, 1964.
7. J. von Neumann, *On regular rings*, Proc. Nat. Acad. Sci., **22** (1936), 707-713.

Received November 6, 1969 and in revised form October 23, 1970. The author wishes to thank J. D. McKnight, Jr., D. M. Morris, A. H. Ortiz and R. L. Snider for the use of their unpublished results as indicated in the text. The author was an N.D.E.A. Fellow at the University of Miami.

NORTHERN ILLINOIS UNIVERSITY  
 AND  
 UNIVERSITY OF MIAMI



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON  
Stanford University  
Stanford, California 94305

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

C. R. HOBBY  
University of Washington  
Seattle, Washington 98105

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
NAVAL WEAPONS CENTER

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Mohammad Shafqat Ali and Marvin David Marcus, <i>On the degree of the minimal polynomial of a commutator operator</i> .....	561
Howard Anton and William J. Pervin, <i>Integration on topological semifields</i> .....	567
Martin Bartelt, <i>Multipliers and operator algebras on bounded analytic functions</i> .....	575
Donald Earl Bennett, <i>Aposyndetic properties of unicoherent continua</i> .....	585
James W. Bond, <i>Lie algebras of genus one and genus two</i> .....	591
Mario Borelli, <i>The cohomology of divisorial varieties</i> .....	617
Carlos R. Borges, <i>How to recognize homeomorphisms and isometries</i> .....	625
J. C. Breckenridge, <i>Burkill-Cesari integrals of quasi additive interval functions</i> .....	635
J. Csima, <i>A class of counterexamples on permanents</i> .....	655
Carl Hanson Fitzgerald, <i>Conformal mappings onto <math>\omega</math>-swirly domains</i> .....	657
Newcomb Greenleaf, <i>Analytic sheaves on Klein surfaces</i> .....	671
G. Goss and Giovanni Viglino, <i>C-compact and functionally compact spaces</i> .....	677
Charles Lemuel Hagopian, <i>Arcwise connectivity of semi-aposyndetic plane continua</i> .....	683
John Harris and Olga Higgins, <i>Prime generators with parabolic limits</i> .....	687
David Michael Henry, <i>Stratifiable spaces, semi-stratifiable spaces, and their relation through mappings</i> .....	697
Raymond D. Holmes, <i>On contractive semigroups of mappings</i> .....	701
Joseph Edmund Kist and P. H. Maserick, <i>BV-functions on semilattices</i> .....	711
Shûichirô Maeda, <i>On point-free parallelism and Wilcox lattices</i> .....	725
Gary L. Musser, <i>Linear semiprime <math>(p; q)</math> radicals</i> .....	749
William Charles Nemitz and Thomas Paul Whaley, <i>Varieties of implicative semilattices</i> .....	759
Jaroslav Nešetřil, <i>A congruence theorem for asymmetric trees</i> .....	771
Robert Anthony Nowlan, <i>A study of H-spaces via left translations</i> .....	779
Gert Kjærgaard Pedersen, <i>Atomic and diffuse functionals on a <math>C^*</math>-algebra</i> .....	795
Tilak Raj Prabhakar, <i>On the other set of the biorthogonal polynomials suggested by the Laguerre polynomials</i> .....	801
Leland Edward Rogers, <i>Mutually aposyndetic products of chainable continua</i> .....	805
Frederick Stern, <i>An estimate for Wiener integrals connected with squared error in a Fourier series approximation</i> .....	813
Leonard Paul Sternbach, <i>On <math>k</math>-shrinking and <math>k</math>-boundedly complete basic sequences and quasi-reflexive spaces</i> .....	817
Pak-Ken Wong, <i>Modular annihilator <math>A^*</math>-algebras</i> .....	825