

Pacific Journal of Mathematics

A STUDY OF H -SPACES VIA LEFT TRANSLATIONS

ROBERT ANTHONY NOWLAN

A STUDY OF H -SPACES VIA LEFT TRANSLATIONS

ROBERT A. NOWLAN

H -spaces are examined by studying left translations, actions and a homotopy version of left translations to be called homolations. If (F, m) is an H -space, the map $s: F \rightarrow F^F$ given by $s(x) = L_x$, i.e. $s(x)$ is left translation by x , is a homomorphism if and only if m is associative. In general, s is an A_n -map if and only if (F, m) is an A_{n+1} space.

The action $r: F^F \times F \rightarrow F$ is given by $r(\varphi, x) = \varphi(x)$. The map s respects the action only of left translations. In general, s respects the action of homolations up to higher-order homotopies. Each homolation generates a family of maps to be called a homolation family. Denoting the set of all homolation families by $H^\infty(F)$, $s: F \rightarrow F^F$ factors through $F \rightarrow H^\infty(F)$ and this latter map is a homotopy equivalence.

By a multiplication on a space F , we mean a continuous map $m: F \times F \rightarrow F$. Let m be a given multiplication on F . For any two points x and y of F , $m(x, y)$ will be denoted by xy and is called the product of x and y . For any point x of F , the assignment $x \rightarrow yx$ and $x \rightarrow xy$ determine respectively the maps

$$L_y: F \longrightarrow F, \quad R_y: F \longrightarrow F$$

called the left and right translation of F by y .

This paper examines H -spaces with strict units by studying left translations and by the introduction of a homotopy version of left translations to be called homolations. One way to use left translations is as follows. If (F, m) is an H -space, the map

$$s: F \longrightarrow F^F$$

given by $s(x) = L_x$, i.e., $s(x)$ is left translation by x , is a homomorphism if and only if m is associative. Other properties of H -structures on a space F can also be interpreted in terms of properties of the map $s: F \rightarrow F^F$.

DEFINITION 1. A map $f: F \rightarrow Y$ is an H -map of the H -space (F, m) into the H -space (Y, w) if $w \circ (f \times f) \cong f \circ m$. (We always use " \cong " to denote "is homotopic to".)

In § II we prove that s is an H -map if and only if m is homotopy associative. In [2], and [3], Stasheff introduces the concepts of A_n -spaces and of A_n -maps, the former generalizes homotopy associativity and the latter generalizes H -maps. We will show that s

is an A_n -map if and only if (F, m) is an A_{n+1} -space.

In § III, H -spaces are studied in terms of actions. The action $r: F^F \times F \rightarrow F$ is given by $r(\varphi, x) = \varphi(x)$. The cross-section $s: F \rightarrow F^F$ respects the action only of left translations. The question arises: of which maps in F^F does s respect the action up to homotopy? This leads to the introduction of T -maps, that is maps $f: F \rightarrow F$ such that $f \circ m \cong m \circ (f \times 1)$. Such maps resemble left translations. Demanding a closer resemblance leads to the introduction of homolations which are maps f satisfying $f \circ m \cong m \circ (f \times 1)$ up to higher order homotopies.

If (F, m) is an associative H -space, a map $w: M \times F \rightarrow F$ is a transitive action if $w \circ (1 \times m) = m \circ (w \times 1)$. The action $r: s(F) \times F \rightarrow F$, where $s(F)$ is the set of all left translations is an example of a transitive action. A homotopy version of a transitive action is given as follows.

DEFINITION 2. Let (F, m) be an associative H -space. A map $w: M \times F \rightarrow F$ is a T -action if $w \circ (1 \times m) \cong m \circ (w \times 1)$.

If $T(F)$ is the maximal subset of F^F such that

$$r: T(F) \times F \longrightarrow F$$

is a T -action, then $T(F)$ consists of T -maps. Generalizing the notions of T -actions leads to the concept of T_n -actions and T_∞ -actions, that is actions $w: M \times F \rightarrow F$ satisfying $w \circ (1 \times m) \cong m \circ (w \times 1)$ up to higher order homotopies. It is then shown that a T_∞ -action of the set of homolations on F can be given such that $s: F \rightarrow F^F$ is a T_∞ -map of actions, i.e., s respects the actions of homolations up to higher order homotopies.

Each homolation generates a family of maps to be called a homolation family. Denote by $H^\infty(F)$ the set of all homolation families. In § IV, it is proven that $s: F \rightarrow F^F$ factors through $F \rightarrow H^\infty(F)$ and that this latter map is a homotopy equivalence.

Throughout this paper, we will be working in the category of k -spaces (i.e., compactly generated spaces) as developed in [5]. The reason for this is to allow unlimited use of the "exponential law." (c.f. Theorem 5, 6 in [5]).

Some of the work included in this paper is contained in my doctoral thesis [1] completed at the University of Notre Dame. Other parts of it were suggested by Professor James D. Stasheff. I deeply appreciate his suggestions and many valuable comments during the writing of this paper.

II. A_n -maps and A_n -spaces We first study H -spaces in relation

to cross-sections to evaluation maps. Let F be any space. Let the evaluation map $v: F^F \rightarrow F$ be defined by $v(\varphi) = \varphi(e)$, where φ is in F^F for some e in F . The map v has a cross-section $s: F \rightarrow F^F$ if and only if F admits a multiplication with right unit e . Given such a cross-section s we can define

$$m(x, y) = s(x)(y) \quad \text{for } x, y \text{ in } F$$

so that m has e as a right unit. Since

$$s(x)(e) = v(s(x)) = x,$$

this multiplication has a two-sided unit if s is a base point preserving map, that is $s(e) = \text{identity}$. We will make this assumption throughout this paper.

If F has a multiplication m with e as right unit, we define $s(x) = L_x$, where L_x is left translation by x . It follows that s is a homomorphism if and only if m is associative.

Thus certain properties of H -structures on a space F can be interpreted in terms of properties of the map $s: F \rightarrow F^F$. As an example we have the following proposition.

PROPOSITION 1. *The map $s: F \rightarrow F^F$ is an H -map if and only if m is homotopy associative.*

Proof. If s is an H -map of (F, m) into (F^F, c) (where c is composition of maps), there exists a homotopy

$$G: I \times F^2 \longrightarrow F^F$$

such that

$$G(0, x, y) = c \circ (s \times s)(x, y) = L_x \circ L_y$$

and

$$G(1, x, y) = s \circ m(x, y) = L_{xy}.$$

Then m can be shown to be homotopy associative by defining a homotopy

$$G': I \times F^3 \longrightarrow F$$

by

$$(1) \quad G'(t, x, y, z) = G(t, x, y)(z)$$

Conversely, if m is homotopy associative, a homotopy G' exists such that

$$G'(0, x, y, z) = x(yz)$$

and

$$G'(1, x, y, z) = (xy)z$$

and the homotopy G can be defined as in (1).

In seeking to generalize this proposition, we first need generalizations of the concepts of homotopy associativity and of H -map. In [2] and [3], Stasheff introduces the concepts of A_n -spaces and of A_n -maps; the former generalizes homotopy associativity and the latter generalizes H -maps. A space which is an A_n -space for all n is said to be an A_∞ -space. Any associative H -space is an A_∞ -space. A_∞ -spaces are homotopy equivalent to associative H -spaces.

DEFINITION 3. An A_n -structure on a space X consists of an n -tuple of maps

$$\begin{array}{ccccccc} X & = & E_1 & \subset & E_2 & \subset & \cdots \subset E_n \\ & & \downarrow p_1 & & \downarrow p_2 & & \downarrow p_n \\ * & = & B_1 & \subset & B_2 & \subset & \cdots \subset B_n \end{array}$$

such that $p_i: \pi_q(E_i, X) \rightarrow \pi_q(B_i)$ is an isomorphism for all q , together with a contracting homotopy $h: CE_{n-1} \rightarrow E_n$ of the cone of E_{n-1} , CE_{n-1} such that $h(CE_{i-1}) \subset E_i$. Such an A_n -structure will be denoted by (p_1, \dots, p_n) . If there exists an infinite collection p_1, p_2, \dots such that for each n , (p_1, \dots, p_n) is an A_n -structure, then we call (p_1, p_2, \dots) an A_∞ -structure.

Theorem 5 of [2] asserts that an A_n -structure on a space X is equivalent to an “ A_n -form”, that is a family of maps $\{M_2, \dots, M_n\}$ where each

$$M_i: I^{i-2} \times X^i \longrightarrow X$$

is suitably defined on the boundary I^{i-1} in terms of M_j for $j < i$.

DEFINITION 4. A space X together with an A_n -form will be called an A_n -space.

In this paper, we are more interested in A_n -forms than A_n -structures, so we introduce the former in some detail. It is first necessary to become acquainted with a special cell-complex K_i which is homeomorphic to I^{i-2} for $i \geq 2$. The standard cells K_i are objects

similar to standard simplices Δ^i and standard cubes I^i , having faces and degeneracies. The difference between the K_i and the simplices and the cubes is that:

(1) The index i does not refer to the dimension of the cell but rather to the number of factors X with which K_i is to be associated.

(2) K_i has degeneracy operators s_1, \dots, s_i defined on it.
and

(3) K_i has $(i(i-1)/2) - 1$ faces.

The following description of the indexing of the faces of K_i is due to Stasheff. Consider a word with i letters, and all meaningful ways of inserting one set of parentheses. To each such insertion except for (x_1, \dots, x_i) , there corresponds a cell of L_i , the boundary of K_i . If the parentheses enclose x_k through x_{k+s-1} , we regard this cell as the homeomorphic image of $K_r \times K_s$ ($r+s=i+1$) under a map which we denote by $\partial_k(r, s)$. Two such cells intersect only on their boundaries and the "edges" so formed correspond to inserting two sets of parentheses in the word. We obtain K_i by induction, starting with $K_2 = *$ (a point), supposing K_2 through K_{i-1} have been constructed. Then construct L_i by fitting together copies of $K_r \times K_s$ subject to certain conditions given in § 2 of [2], that is the fitting together of copies of $K_r \times K_s$ as dictated by the above description of the indexing. Finally, take K_i to be the cone on L_i .

The following is part of Theorem 5 of [2].

THEOREM 2. *A space X admits an A_n -structure if and only if there exist maps $M_i: K_i \times X^i \rightarrow X$ for $2 \leq i \leq n$ such that*

- (1) $M_2(*, e, x) = M_2(*, x, e) = x$ for x in X , $* = K_2$ and
- (2) For $\rho \in K_r$, $\sigma \in K_s$, $r+s=i+1$, we have

$$\begin{aligned} & M_i(\partial_k(r, s)(\rho, \sigma), x_1, \dots, x_i) \\ &= M_r(\rho, x, \dots, x_{k-1}, M_s(\sigma, x_k, \dots, x_{k+s-1}), \dots, x_i). \end{aligned}$$

We note that an A_2 -space is just an H -space. In the case $i=3$, K_3 is homeomorphic to I and (2) asserts that M_3 is a homotopy between $M_2 \circ (M_r \times 1)$ and $M_2 \circ (1 \times M_s)$, to be imprecise between $(xy)z$ and $x(yz)$. Thus M_3 is an associating homotopy and M_2 is a homotopy associative action.

In the case $i=4$, we consider the five ways of associating a product of four factors. If the multiplication M_2 is a homotopy associative multiplication, the five products are then related by the following string of homotopies:

$$x(y(zw)) \cong x((yz)w) \cong (x(yz))w \cong ((xy)z)w \cong (xy)(zw) \cong x(y(zw)).$$

Thus we have defined a map of $S^1 \times X^4 \rightarrow X$ and the map M_4 can

be regarded as an extension of the map to $I^2 \times X^4$.

If X is an associative H -space, it admits A_∞ -forms; it is only necessary to define

$$M_i(\tau, x_1, \dots, x_i) = x_1 x_2 \dots x_i \text{ for } \tau \text{ in } K_i \text{ and } 1 \leq i.$$


This will be called a trivial A_∞ -form. If X is an A_∞ -space then there is an associative H -space Y of the homotopy type of X .

DEFINITION 5. Let $(X, \{M_i\})$ be an A_n -space and (Y, w) be an associative H -space. A map $f: X \rightarrow Y$ is an A_n -map if there exists maps $h_i: K_{i+1} \times X^i \rightarrow Y$, $1 \leq i \leq n$, called sputnik homotopies, such that $h_1 = f$ and for ρ in K_r , σ in K_s ($r + s = i + 1$), we have

$$\begin{aligned} & h_i(\partial_k(r, s)(\rho, \sigma), x_1, \dots, x_i) \\ &= h_{r-1}(\rho, x_1, \dots, x_{k+1}, M_s(\sigma, x_k, \dots, x_{k+s-1}), \dots, x_i) \text{ if } k \neq r \\ &= h_{r-1}(\rho, x_1, \dots, x_{r-1}) h_{s-1}(\sigma, x_r, \dots, x_i) \text{ if } k = r. \end{aligned}$$

Note that when $n = 2$, f is just an H -map, as h_2 is a homotopy between $f \circ M_2$ and $w \circ (f \times f)$. In the case $n = 3$, since K_4 is homeomorphic to I^2 , we have a map of $S^1 \times X^3 \rightarrow Y$ and $h_3: K_4 \times X^3 \rightarrow Y$ can be thought of as an extension of this map to $I^2 \times X^4$.

Consider the following cross-section of $I^2 \times X^4$ showing a typical I^2 . Assign to the "faces" of I^2 the homotopies $h_2 \circ (M_2 \times 1)$, $w \circ (h_2 \times h_1)$, $w \circ (h_1 \times h_2)$, $h_2 \circ (1 \times M_2)$ and $h_1 \circ M_3$ as indicated



The broken line represents a point. The map h_3 then appropriately fills in the figure.

A map which is an A_n -map for all n will be called an A_∞ -map.

We are now in a position to prove the following generalization of proposition 1.

THEOREM 3. (A) Let $(F, \{M_i\})$ be an A_n -space; then $s: F \rightarrow F^F$ is an A_{n-1} map.

(B) s can be shown to be an A_n -map if and only if $(F, \{M_i\})$ can

be given the structure of an A_{n+1} space.

Proof. (A) Given that $(F, \{M_i\})$ is an A_n -space, all that is necessary to show that s is an A_{n+1} map is to define $h_1 = s$ and $h_i: K_{i+1} \times F^i \rightarrow F^F$ $1 \leq i \leq n-1$ by

$$h_i(\partial_k(r, t)(\rho, \sigma), x_1, \dots, x_i)(y) = M_{i+1}(\partial_k(r, t)(\rho, \sigma), x_1, \dots, x_i, y) .$$

(B) It is clear that $(F, \{M_i\})$ can be extended to an A_{n+1} -space (that is there exists a map $M_{n+1}: K_{n+1} \times F^{n+1} \rightarrow F$) if and only if there exists a map $h_n: K_{n+1} \times F^n \rightarrow F^F$ given by

$$h_n(\partial_k(r, t)(\rho, \sigma), x_1, \dots, x_n)(y) = M_{n+1}(\partial_k(r, t)(\rho, \sigma), x_1, \dots, x_n, y) .$$

COROLLARY 4. An A_∞ -form on F is equivalent to the existence of sputnik homotopies $h_i: K_{i+1} \times F^i \rightarrow F^F$ for all i making s an A_∞ -map.

III. T_n -maps and Homotations. We assume throughout this section that (F, m) is an associative H -space with a strict unit. In that case, the map

$$s: F \longrightarrow F^F$$

given by

$$s(f)(y) = m(f, y)$$

is a homomorphism.

We now study left translations via actions. The space F^F acts on F by

$$\begin{aligned} r: F^F \times F &\longrightarrow F \\ r(\varphi, f) &= \varphi(f) . \end{aligned}$$

The cross-section s respects the action only of left translations, for consider the diagram:

$$(1) \quad \begin{array}{ccc} F^F \times F & \xrightarrow{1 \times s} & F^F \times F^F \\ r \downarrow & & \downarrow c \\ F & \xrightarrow{s} & F^F . \end{array}$$

Suppose

$$s(\varphi(f)) = \varphi \circ s(f) .$$

Since s is left translation, we have $\varphi(fy) = \varphi(f)y$, that is the following diagram is commutative.

$$(2) \quad \begin{array}{ccc} F \times F & \xrightarrow{\quad m \quad} & F \\ \varphi \times 1 \downarrow & & \downarrow \varphi \\ F \times F & \xrightarrow{\quad m \quad} & F \end{array} .$$

In particular,

$$\varphi(y) = \varphi(ey) = \varphi(e)y$$

and φ is left translation by $\varphi(e)$. So diagram (1) commutes only on $s(F) \times F \subset F^F \times F$ where $s(F)$ is the set of left translations. Thus s is a map of spaces on which $s(F)$ acts.

The result tells us something about the action

$$r: s(F) \times F \longrightarrow F$$

namely, it is transitive.

Note that the following diagram is commutative

$$(3) \quad \begin{array}{ccc} s(F) \times F \times F & \xrightarrow{\quad 1 \times m \quad} & s(F) \times F \\ r \times 1 \downarrow & & \downarrow r \\ F \times F & \xrightarrow{\quad m \quad} & F \end{array} .$$

Let us consider the following question: what is the nature of the action r when diagram (1) is only required to be homotopy commutative. Denote by $T_2(F)$ the maximal subset of maps φ in F^F such that

$$s[\varphi(f)] \cong \varphi_0 s(f)$$

in the sense that there exists a homotopy

$$\theta_2: I \times T_2(F) \times F \longrightarrow F^F$$

such that

$$\theta_2(0, \varphi, f) = \varphi \circ s(f)$$

and

$$\theta_2(1, \varphi, f) = s[\varphi(f)] .$$

In this case, it follows that for each φ in $T_2(F)$ there exists a homotopy

$$\varphi_2: I \times F^2 \longrightarrow F$$

depending continuously on φ such that

$$\varphi_2(0, f, y) = \varphi(fy)$$

and

$$\varphi_2(1, f, y) = \varphi(fy)y .$$

DEFINITION 6. Let (F, m) be an associative H -space. A map $f: F \rightarrow F$ is a T -map if there exists a homotopy $I \times F^2 \rightarrow F$ such that $f \circ m \cong m \circ (f \times 1)$.

Thus we see that the maps in $T_2(F)$ are T -maps. The homotopy is given by

$$\varphi_2(t, f, y) = \theta_2(t, \varphi, f)(y) .$$

In particular, we note that for each φ in $T_2(F)$

$$\varphi(y) = \varphi(ey) \cong \varphi(e)y$$

indicating that up to homotopy φ acts like left translation by $\varphi(e)$. Thus the maps in $T_2(F)$ in this sense resemble left translations. We will investigate this resemblance further.

Our results show that the action

$$r: T_2(F) \times F \longrightarrow F^F$$

is a T -action in the sense that there exists a homotopy

$$\lambda_2: I \times T_2(F) \times F^2 \longrightarrow F$$

such that

$$\lambda_2: r \circ (1 \times m) \cong m \circ (r \times 1) .$$

In fact, we can take λ_2 to be adjoint to θ_2 :

$$\lambda_2(t, \varphi, f, y) = \theta_2(t, \varphi, f)(y) .$$

If φ is a true left translation, it follows that

$$\varphi(xyz) = \varphi(xy)z = \varphi(x)yz \quad \text{for } x, y, z \text{ in } F$$

however for a map φ in $T_2(F)$, the most we can claim using a rather loose notation is that:


$$\varphi(xyz) \cong \varphi(xy)z \cong \varphi(x)yz \cong \varphi(xyz) .$$

This string of homotopies defines a map

$$\dot{I}^2 \times F \longrightarrow F$$

where \dot{I}^2 is the boundary of I^2 .

This can be illustrated in the following diagram, representing $\dot{I}^2 \times F^3$ showing only \dot{I}^2 with “faces” labeled by the homotopies connecting the maps given above. Note that the edge of \dot{I}^2 represented by the broken line is just a point. (This is because F is an associative H -space. If F were only homotopy associative, this face would be labeled by the associating homotopy applied to $\varphi(x), y, z$. The following discussion could be carried out for A_n -spaces but the details are bad enough in the associative case, which is the case of interest for applications [1].)



The problem of making a map φ in $T_2(F)$ more closely “resemble” a left translation, requires that we be able to extend the map

$$\dot{I}^2 \times F^3 \longrightarrow F$$

to a map

$$I^2 \times F^3 \longrightarrow F.$$

Thus we will need higher homotopy conditions on the maps φ in $T_2(F)$. Suppose for the moment that there exists a map

$$\varphi_3: I^2 \times F^3 \longrightarrow F$$

such that

$$\varphi_3(0, t_2, x, y, z) = \varphi_2(t_2, xy, z)$$

$$\varphi_3(t_1, 0, x, y, z) = \varphi_2(t_1, x, yz)$$

$$\varphi_3(1, t_2, x, y, z) = \varphi(x)yz$$

and

$$\varphi_3(t_1, 1, x, y, z) = \varphi_2(t_1, x, y) \cdot z.$$

Let $T_3(F)$ denote the maximal subset of $T_2(F)$ such that for each φ in $T_2(F)$, there exists φ_2 and φ_3 depending continuously on φ and φ_2 subject to the conditions already mentioned. In this case, the action $r: T_3(F) \times F \rightarrow F$ is such that there exist maps

$$\lambda_2: I \times T_3(F) \times F^2 \longrightarrow F$$

such that

$$\lambda_2: r(1 \times m) \cong m(r \times 1)$$

and

$$\lambda_3: I^2 \times T_3(F) \times F^3 \longrightarrow F$$

such that

$$\lambda_3(0, t_2, \varphi, x, y, z) = \lambda_2(t_2, \varphi, xy, z)$$

$$\lambda_3(t_1, 0, \varphi, x, y, z) = \lambda_2(t_1, \varphi, x, yz)$$

$$\lambda_3(1, t_2, \varphi, x, y, z) = r(\varphi, x) \cdot yz$$

and

$$\lambda_3(t_1, 1, \varphi, x, y, z) = \lambda_2(t_1, \varphi, x, y) \cdot z.$$

This latter map is given by

$$\lambda_3(t_1, t_2, \varphi, x, y, z) = \varphi_3(t_1, t_2, x, y, z).$$

On the other hand, there exist maps

$$\theta_2: I \times T_3(F) \times F \longrightarrow F^F$$

such that

$$\theta_2: \varphi \circ s(f) \cong s[\varphi(f)]$$

and

$$\theta_3: I^2 \times T_3(F) \times F^2 \longrightarrow F^F$$

such that

$$\theta_3: (t_1, t_2, \varphi, x, y)(z) = \lambda_3(t_1, t_2, \varphi, x, y, z).$$

Parallel to every demand that a map $\varphi: F \rightarrow F$ more closely resemble a left translation by satisfying higher homotopy conditions will be the requirement of higher homotopy conditions on the action r and similar higher homotopy conditions on the map s .

DEFINITION 7. Let (X, m) be an associative H -space. A map $\varphi: X \rightarrow X$ is a T_n -map of X into itself if there exists a family of maps

$$\varphi_i: I^{i-1} \times X^i \longrightarrow X \qquad 1 \leq i \leq n$$

such that $\varphi_1 = \varphi$ and

$$\begin{aligned}
& \varphi_i(t_1, \dots, t_{i-1}, x_1, \dots, x_i) \\
&= \varphi_{i-1}(t_1, \dots, \hat{t}_k, \dots, t_{i-1}, x_1, \dots, x_k x_{k+1}, \dots, x_i) & \text{if } t_k = 0 \\
&= \varphi_k(t_1, \dots, t_{k-1}, x_1, \dots, x_k) \cdot (x_{k+1}, x_{k+2} \dots x_i) & \text{if } t_k = 1.
\end{aligned}$$

In case φ_i exists for all i , we call φ a homolation, that is, a homotopy translation. Denote the set of all homolations by $T_\infty(F)$.

DEFINITION 8. Let (F, m) be an associative H -space. A homolation family on F is a collection of maps $\{\varphi_i: I^{i-1} \times F^i \rightarrow F, \forall i \geq 1\}$ where φ_1 is a homolation and $\varphi_1: F \rightarrow F$ is a homotopy equivalence. We will denote by $H^\infty(F)$, the set of all homolation families. $H^\infty(F)$ is a subspace of $C(F; F) \times C(I \times F^2; F) \times \dots$ where $C(I^j \times F^{j+1}; F)$ is the set of all continuous maps $f: I^j \times F^{j+1} \rightarrow F$ (with the k -topology derived from the compact-open topology).

DEFINITION 9. Let (X, m) be an associative H -space. A map

$$w: M \times X \longrightarrow X$$

of M on X is said to be a T_n -action if there exist maps

$$w_i: I^{i-1} \times M \times X^i \longrightarrow X \quad 1 \leq i \leq n$$

such that $w_1 = w$ and

$$\begin{aligned}
& w_i(t_1, \dots, t_{i-1}, g, x_1, \dots, x_i) \\
&= w_{i-1}(t_1, \dots, \hat{t}_k, \dots, t_{i-1}, g, \dots, x_k x_{k+1}, \dots, x_i) & \text{if } t_k = 0 \\
&= w_k(t_1, \dots, t_{k-1}, g, x_1, \dots, x_k) \cdot (x_{k+1}, x_{k+2} \dots x_i) & \text{if } t_k = 1.
\end{aligned}$$

If a map $w: M \times X \rightarrow X$ is a T_n -action for all n , then w is said to be a T_∞ -action.

THEOREM 5. Let $T_n(F)$ denote the maximal subset of F^F such that there exist maps $\lambda_i: I^{i-1} \times T_n(F) \times F^i \rightarrow F$ for $1 \leq i \leq n$ making $r: T_n(F) \times F \rightarrow F$ a T_n -action; then $T_n(F)$ consists of T_n -maps.

Proof. We may define the maps

$$\varphi_i: I^{i-1} \times F^i \longrightarrow F \quad 1 \leq i \leq n$$

by

$$\varphi_i(t_1, \dots, t_{i-1}, f_1, \dots, f_i) = \lambda_i(t_1, \dots, t_{i-1}, \varphi, f_1, \dots, f_i).$$

DEFINITION 10. Let (X, m) and (M, v) be associative H -spaces and $w: M \times X \rightarrow X$ be a T_n -action. A homomorphism $f: X \rightarrow M$ is said to be a T_n -map of actions if there exist maps

$$\theta_i: I^{i-1} \times M \times X^{i-1} \longrightarrow M$$

such that $\theta_1 = 1_M$ and

$$\begin{aligned} & \theta_i(t_1, \dots, t_{i-1}, g, x_1, \dots, x_{i-1}) \\ &= \theta_{i-1}(t_1, \dots, \hat{t}_k, \dots, t_{i-1}, g, \dots, x_k x_{k+1}, \dots, x_{i-1}) \\ & \hspace{15em} \text{if } t_k = 0, k \neq i-1 \\ &= v[\theta_{i-1}(t_1, \dots, t_{i-2}, g, x_1, \dots, x_{i-2}), f(x_{i-1})] \hspace{2em} \text{if } t_{i-1} = 0 \\ &= f[m(w_k(t_1, \dots, t_{k-1}, g, x_1, \dots, x_k), x_{k+1} x_{k+2} \dots x_i)] \hspace{2em} \text{if } t_k = 1. \end{aligned}$$

If θ_i exists for all i , then f is said to be a T_∞ -map of actions.

COROLLARY 6. *The map $r: T_\infty(F) \times F$ is a T_∞ -action and s is then a T_∞ -map of actions.*

Proof. Define $\lambda_i: I^{i-1} \times T_\infty(F) \times F^i \rightarrow F$ by

$$\lambda_i(t_1, \dots, t_{i-1}, \varphi, f_1, \dots, f_i) = \varphi_i(t_1, \dots, t_{i-1}, f_1, \dots, f_i)$$

and

$$\theta_i: I^{i-1} \times T^\infty(F) \times F^{i-1} \longrightarrow F^F$$

by

$$\theta_i(t_1, \dots, t_{i-1}, \varphi, f_1, \dots, f_{i-1})(f_i) = \lambda_i(t_1, \dots, t_{i-1}, \varphi, f_1, \dots, f_i).$$

IV. The homotopy equivalence of F and $H^\infty(F)$. As we have seen, we can identify an associative H -space with the set of left translations of that space. We note that this identification of F in F^F as left translation is not homotopy invariant: $\varphi(fx) = \varphi(f)x$ is not a homotopy statement. Our definition of homolation is homotopy invariant and it characterizes $F \rightarrow F^F$ from a homotopy point of view.

We are now in a position to prove the following theorem. Recall that $H^\infty(F)$ is the set of all homolation families.

THEOREM 7. *If (F, m) is a connected associative H -space, the map $s: F \rightarrow F^F$ factors through $H^\infty(F)$, and the factor $F \rightarrow H^\infty(F)$ is a homotopy equivalence.*

Proof. Define a map

$$\tau: F \longrightarrow H^\infty(F)$$

as follows:

$$\tau(f) = \Phi_f = \{\varphi_1^f, \varphi_2^f, \dots\}$$

where

$$\varphi_1^f: F \longrightarrow F$$

is given by

$$\varphi_1^f(g) = fg$$

that is left translation of F . φ_1^f is a homotopy equivalence since F is connected (see [4]).

The remaining maps are given by

$$\varphi_k^f(t_1, \dots, t_{k-1}, f_1, \dots, f_k) = ff_1 \cdots f_k \quad \text{for all } k.$$

The map τ is continuous, since the composition of maps

$$F \xrightarrow{\tau} C(F; F) \times C(I \times F^2; F) \cdots \xrightarrow{p^{(k)}} C(I^{k-1} \times F^k; F)$$

is continuous for each k and $p^{(k)}$ is projection onto the corresponding factor.

On the other hand, define the map

$$\mu: H^\infty(F) \longrightarrow F$$

by

$$\mu(I) = \gamma_1(e)$$

where $I = \{\gamma_1, \gamma_2, \dots\}$ is in $H^\infty(F)$ and e is the unit of F .

The map μ is continuous, since it is the composition of maps

$$H^\infty(F) \xrightarrow{p_1} H^\infty(F)_1 = T_\infty(F) \xrightarrow{w_e} F$$

where p_1 is projection of $H^\infty(F)$ on that part of $H^\infty(F)$ contained in F^F , namely the set of homotations, here denoted by $H^\infty(F)_1$, and the map w_e is the evaluation map at e (continuous in the k -topology).

Note that $\mu(\tau(f)) = \mu(\Phi_f) = \varphi_1^f(e) = fe = f$ so that $\mu \circ \tau = 1_F$.

On the other hand

$$\tau \circ \mu(I) = \tau(\gamma_1(e)) = \Phi_{\gamma_1(e)} = \{\varphi_1^{\gamma_1(e)}, \varphi_2^{\gamma_1(e)}, \dots\}.$$

We claim that $\tau \circ \mu \cong 1_{H^\infty(F)}$, that is there exists a map

$$H_i: H^\infty(F) \longrightarrow H^\infty(F)$$

such that $H_0 = 1_{H^\infty(F)}$ and $H_1 = \tau \circ \mu$.

To see this, let $H^\infty(F)_k$ be the subspace of $H^\infty(F)$ which is contained in $C(I^{k-1} \times F^k; F)$. The map $H_i = \{H_i^1, H_i^2, \dots\}$ will consist of homotopies

$$\{H_t^k\}: H^\infty(F) \longrightarrow H^\infty(F)_k \quad \text{for each } k$$

such that $H_0^k = 1_{H^\infty(F)_k}$ and $H_1^k = \tau \circ \mu \mid H^\infty(F)_k$ and the H_t^k are compatible.

Define $H_t^k: H^\infty(F) \rightarrow H^\infty(F)_k$ as follows:

$$H_t^k(\Gamma)(t_1, \dots, t_{k-1}, f_1, \dots, f_k) = \gamma_{k+1}(t, t_1, \dots, t_{k-1}, e, f_1, \dots, f_k) .$$

The map is continuous as each γ_{k+1} in Γ is continuous and $\Gamma \rightarrow \gamma_{k+1}$ is continuous being projection.

Note if $t_j = 0$

$$\begin{aligned} & H_t^k(\Gamma)(t_1, \dots, t_{k-1}, f_1, \dots, f_k) \\ &= \gamma_k(t, t_1, \dots, \hat{t}_j, \dots, t_{k-1}, e, f_1, \dots, f_j f_{j+1}, \dots, f_k) \\ &= H_t^{k-1}(\Gamma)(t_1, \dots, \hat{t}_j, \dots, t_{k-1}, f_1, \dots, f_j f_{j+1}, \dots, f_k) \end{aligned}$$

while if $t_j = 1$

$$\begin{aligned} & H_t^k(\Gamma)(t_1, \dots, t_{k-1}, f_1, \dots, f_k) \\ &= \gamma_{j+1}(t, t_1, \dots, t_{j-1}, e, f_1, \dots, f_j)(f_{j+1}, \dots, f_k) \\ &= H_t^j(\Gamma)(t_1, \dots, t_{j-1}, f_1, \dots, f_j)(f_{j+1}, \dots, f_k) . \end{aligned}$$

Thus $\{H_t^k\}$ is in $H^\infty(F)$. Further

$$\begin{aligned} H_0^k(\Gamma)(t_1, \dots, t_{k-1}, f_1, \dots, f_k) &= \gamma_k(t_1, \dots, t_{k-1}, e f_1, \dots, f_k) \\ &= \gamma_k(t_1, \dots, t_{k-1}, f_1, \dots, f_k) . \end{aligned}$$

Thus $H_0^k = 1_{H^\infty(F)_k} \{H_0^k(\Gamma)\} = \Gamma$ and

$$\begin{aligned} & H_1^k(\Gamma)(t_1, \dots, t_{k-1}, f_1, \dots, f_k) \\ &= \gamma_1(e) f_1 \dots f_k \\ &= \varphi_1^{i_1(e)}(t_1, \dots, t_{k-1}, f_1, \dots, f_k) \\ &= \tau \circ \mu(\Gamma)(t_1, \dots, t_{k-1}, f_1, \dots, f_k) . \end{aligned}$$

Thus $H_1^k = \tau \circ \mu \mid H^\infty(F)_k$, $\{H_1^k(\Gamma)\} = \tau \circ \mu(\Gamma)$. This completes the proof that F and $H^\infty(F)$ are homotopy equivalent.

Now $H^\infty(F)$ is itself an H -space; we can define composition of families as well as just maps $F \rightarrow F$ (see [1]). The map $F \rightarrow H^\infty(F)$ is an A_∞ -map and hence induces $B_F \rightarrow B_{H^\infty(F)}$ which is again a homotopy equivalence if F is a CW -complex.

In my thesis [1], I show that $B_{H^\infty(F)}$ is a classifying space for fibrations with A_∞ -actions of F on the total space. The above homotopy equivalence then shows a fibre space admits such an A_∞ -action if and only if it admits an associative action.

REFERENCES

1. R. A. Nowlan, *A_n -actions on Fibre Spaces*, Thesis, University of Notre Dame, 1969.
2. J. D. Stasheff, *Homotopy Associativity of H -spaces (I)*, Trans. Amer. Math. Soc., **108** (1963), 275-292.
3. ———, *Homotopy Associativity of H -spaces (II)*, Trans. Amer. Math. Soc., **108** (1963), 293-312.
4. ———, *On Homotopy Abelian H -spaces*, Proc. Cambridge Phil. Soc., **57** (1961), 734-745.
5. N. E. Steenrod, *A convenient category of topological spaces*, Michigan Math. J., (1967), 133-152.

Received April 6, 1970 and in revised form July 9, 1970.

SOUTHERN CONNECTICUT STATE COLLEGE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics

Vol. 37, No. 3

March, 1971

Mohammad Shafqat Ali and Marvin David Marcus, <i>On the degree of the minimal polynomial of a commutator operator</i>	561
Howard Anton and William J. Pervin, <i>Integration on topological semifields</i>	567
Martin Bartelt, <i>Multipliers and operator algebras on bounded analytic functions</i>	575
Donald Earl Bennett, <i>Aposyndetic properties of unicoherent continua</i>	585
James W. Bond, <i>Lie algebras of genus one and genus two</i>	591
Mario Borelli, <i>The cohomology of divisorial varieties</i>	617
Carlos R. Borges, <i>How to recognize homeomorphisms and isometries</i>	625
J. C. Breckenridge, <i>Burkill-Cesari integrals of quasi additive interval functions</i>	635
J. Csima, <i>A class of counterexamples on permanents</i>	655
Carl Hanson Fitzgerald, <i>Conformal mappings onto ω-swirly domains</i>	657
Newcomb Greenleaf, <i>Analytic sheaves on Klein surfaces</i>	671
G. Goss and Giovanni Viglino, <i>C-compact and functionally compact spaces</i>	677
Charles Lemuel Hagopian, <i>Arcwise connectivity of semi-aposyndetic plane continua</i>	683
John Harris and Olga Higgins, <i>Prime generators with parabolic limits</i>	687
David Michael Henry, <i>Stratifiable spaces, semi-stratifiable spaces, and their relation through mappings</i>	697
Raymond D. Holmes, <i>On contractive semigroups of mappings</i>	701
Joseph Edmund Kist and P. H. Maserick, <i>BV-functions on semilattices</i>	711
Shûichirô Maeda, <i>On point-free parallelism and Wilcox lattices</i>	725
Gary L. Musser, <i>Linear semiprime $(p; q)$ radicals</i>	749
William Charles Nemitz and Thomas Paul Whaley, <i>Varieties of implicative semilattices</i>	759
Jaroslav Nešetřil, <i>A congruence theorem for asymmetric trees</i>	771
Robert Anthony Nowlan, <i>A study of H-spaces via left translations</i>	779
Gert Kjærgaard Pedersen, <i>Atomic and diffuse functionals on a C^*-algebra</i>	795
Tilak Raj Prabhakar, <i>On the other set of the biorthogonal polynomials suggested by the Laguerre polynomials</i>	801
Leland Edward Rogers, <i>Mutually aposyndetic products of chainable continua</i>	805
Frederick Stern, <i>An estimate for Wiener integrals connected with squared error in a Fourier series approximation</i>	813
Leonard Paul Sternbach, <i>On k-shrinking and k-boundedly complete basic sequences and quasi-reflexive spaces</i>	817
Pak-Ken Wong, <i>Modular annihilator A^*-algebras</i>	825