

# Pacific Journal of Mathematics

**NOTE ON THE OPEN MAPPING THEOREM**

LAWRENCE GERALD BROWN

## NOTE ON THE OPEN MAPPING THEOREM

LAWRENCE G. BROWN

**The open mapping and closed graph theorems are usually stated in terms of metrizable topological groups which are complete in a one-sided uniformity. It would be desirable to use only the weaker hypothesis of completeness in the two-sided uniformity. Important results of this sort are already known, and it is our purpose to strengthen those results by removing the separability hypotheses.**

There are three uniformities normally used in connection with a topological group: the left, right, and two-sided uniformities. Completeness in the left uniformity is equivalent to completeness in the right uniformity, and stronger than completeness in the two-sided uniformity. If the group is metrizable, then it always has at least one left invariant metric; completeness in the left uniformity is equivalent to completeness in this metric (it does not matter which left-invariant metric is chosen). Still in the case of a metrizable group, completeness in the two-sided uniformity is equivalent to topological completeness, i.e., the existence of at least one metric in which the group is complete as a metric space (see [3], Exercise  $Q(d)$ , p. 212 for proof). Every topological group has a completion in the two-sided sense, but not necessarily in the one-sided sense (more precisely, the completion in the left-uniformity, though it is a complete uniform space, is not in general a group). When the one-sided completion does exist, it coincides with the two-sided completion. Two important examples of metrizable topological groups which are complete in the two-sided but not the one-sided sense are the full permutation group of a countably infinite set and the unitary group of an infinite-dimensional, separable Hilbert space (with the strong operator topology). Both of these examples are separable. Nonseparable examples of less naturalness can be obtained by taking the direct product of one of the above with a nonseparable Banach space.

Lemma 1 below, which is the separable case of our theorem, was essentially proved by Banach [1], though his statement of it (Satz 8) was weaker. Its present statement and brief proof are the same as in Corollary 3.2 of Pettis [5]. Corollary 1 of the theorem and the corollary to Lemma 1 have some independent interest. They state that if  $N$  is a closed normal subgroup of the group  $G$ , which is metrizable and complete in its two-sided uniformity, then  $G/N$  is also complete. The corresponding result for one-sided completeness is trivial and is true even if  $N$  is not normal. (This case where  $N$  is

not normal is, so far as we know, the only case where the results for one-sided completeness remain significantly superior to those for two-sided completeness.) Lemma 2, which is corollary 2 to the theorem, may also have some independent interest. Our theorem itself, when strengthened by the addition of the accompanying remark, differs from the "standard"<sup>1</sup> version of the open mapping theorem ([2], Theorem 3, p. 90) only in the type of completeness used. Our version of the closed graph theorem, Corollary 3, is slightly weaker than the analog of the "standard" version in that both groups are assumed metrizable.

LEMMA 1. *Let  $G$  and  $H$  be separable metrizable topological groups which are complete in their two-sided uniformities and  $\pi: G \rightarrow H$  a continuous homomorphism such that  $\overline{\pi(U)}$  is a neighborhood of the identity for each neighborhood  $U$  of the identity in  $G$ . Then  $\pi$  is open.*

REMARK. It is easy to see that the hypothesis on  $\overline{\pi(U)}$  will be satisfied if  $\pi$  is surjective. This special case is Satz 8 of [1].

*Proof.* Let  $U$  be an open neighborhood of the identity in  $G$ . Then  $\pi(U)$  is analytic and hence ([4], p. 94-95) almost open. But  $\pi(U)$  is second category in  $H$ . (If  $\pi(U) \subseteq \bigcup_n F_n$ ,  $F_n$  closed, then some  $\pi^{-1}(F_n) \cap U$  has interior. Hence  $\overline{\pi^{-1}(F_n)} \subseteq F_n$  has interior.) Hence (see [5] or [3], Exercise P(b), p. 211)  $\pi(U) \cdot \pi(U)^{-1}$  is a neighborhood of the identity in  $H$ .

COROLLARY. *If  $G$  is a polonais group (i.e., a separable, metrizable, and topologically complete topological group) and  $N$  is a closed normal subgroup, then  $G/N$  is again complete and hence polonais.*

*Proof.* Let  $H$  be the completion of  $G/N$ .

LEMMA 2. *Let  $G$  and  $H$  be metrizable topological groups which are complete in their two-sided uniformities and  $\pi: G \rightarrow H$  a continuous homomorphism such that  $\overline{\pi(U)}$  is a neighborhood of the identity in  $H$  for each neighborhood  $U$  of the identity in  $G$  and such that  $\overline{\pi(G)} = H$ . Then for each closed separable subgroup  $H_0$  of  $H$ , there is a closed separable subgroup  $G_0$  of  $G$  such that  $\pi$  is a relatively open map of  $G_0$  onto  $H_0$ .*

<sup>1</sup> A perhaps better choice for the "standard" version appears in [3; p. 213]. It has a weaker hypothesis.

*Proof.* Let  $U_1 \supseteq U_2 \supseteq U_3 \cdots$  be a fundamental system of neighborhoods of the identity in  $G$ . Let  $V_j (j = 1, 2, \dots)$  be open neighborhoods of the identity in  $H$  such that  $\overline{\pi(U_j)} \supseteq V_j$ . Let  $G_1$  be a closed separable subgroup of  $G$  such that  $\overline{\pi(G_1)} \supseteq H_0$  and

$$\overline{\pi(G_1 \cap U_j)} \supseteq H_0 \cap V_j.$$

Let  $H_1 = \overline{\pi(G_1)}$ . Similarly, define  $G_n (n = 2, 3, \dots)$  and  $H_n (n = 2, 3, \dots)$  so that for  $n \geq 1$ :

- (i)  $G_n \subseteq G_{n+1}, H_n \subseteq H_{n+1},$
- (ii)  $\overline{\pi(G_n)} = H_n,$
- (iii)  $\overline{\pi(G_n \cup U_j)} \supseteq H_{n-1} \cap V_j,$
- (iv)  $G_n, H_n$  are closed and separable.

Let  $G_\infty = \bigcup_{n=1}^\infty G_n$  and  $H_\infty = \bigcup_{n=1}^\infty H_n$ . Then by Lemma 1,  $\pi$  is a relatively open map of  $G_\infty$  onto  $H_\infty$ . Let  $G_0 = \pi^{-1}(H_0) \cap G_\infty$ .

**THEOREM.** *Let  $G$  and  $H$  be metrizable topological groups which are complete in their two-sided uniformities and  $\pi: G \rightarrow H$  a continuous homomorphism such that  $\overline{\pi(U)}$  is a neighborhood of the identity in  $H$  for each neighborhood  $U$  of the identity in  $G$ . Then  $\pi$  is open.*

**REMARK.** It is easy to remove all hypotheses on  $H$  (except that  $H$  be a Hausdorff topological group) from the statement of the theorem. (It is easy to construct a countable base at  $e$  and hence prove  $H$  metrizable.  $H$  can then be completed.)

*Proof.* Since  $\overline{\pi(G)}$  is an open subgroup of  $H$ , we may assume  $\pi(G)$  dense. Let  $y_n$  be a sequence in  $H$  so that  $\lim y_n = e$ . Then  $\{y_n\}$  is imbedded in a closed separable subgroup  $H_0$  of  $H$ . Choose  $G_0$  as in Lemma 2. Then there are  $x_n$  in  $G_0$  such that  $\pi(x_n) = y_n$  and  $\lim x_n = e$ .

**COROLLARY 1.** *If  $G$  is a metrizable topological group which is complete in its two-sided uniformity and  $N$  a closed normal subgroup, then  $G/N$  is complete in its two-sided uniformity.*

**COROLLARY 2.** *With  $G, N$  as in Corollary 1, any closed separable subgroup of  $G/N$  is the image of a closed separable subgroup of  $G$ .*

**COROLLARY 3.** *If  $G$  and  $H$  are as in the theorem, and  $f: G \rightarrow H$  is a homomorphism such that  $f$  has a closed graph, and  $f^{-1}(U)$  is a neighborhood of the identity in  $G$  for each neighborhood of the identity  $U$  in  $H$ , then  $f$  is continuous.*

*Proof.* Apply the theorem to the projection of  $K$  onto  $G$ , where  $K$  is the graph of  $f$ .

#### REFERENCES

1. S. Banach, *Metrische Gruppen*, *Studia Math.*, **3** (1931), 101-113.
2. T. Husain, *Introduction to Topological Groups*, W. B. Saunders, Philadelphia, 1966.
3. J. Kelley, *General Topology*, Van Nostrand, Princeton, 1955.
4. K. Kuratowski, *Topology Vol. I* (translated from the French by J. Jawarowki), Academic Press, New York, 1966.
5. B. Pettis, *On continuity and openness of homomorphisms in topological groups*, *Ann. of Math.*, **54** (1950), 293-308.

Received February 4, 1970.

STANFORD UNIVERSITY

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON  
Stanford University  
Stanford, California 94305

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

C. R. HOBBY  
University of Washington  
Seattle, Washington 98105

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
NAVAL WEAPONS CENTER

# Pacific Journal of Mathematics

Vol. 38, No. 1

March, 1971

Bruce Alan Barnes, <i>Banach algebras which are ideals in a Banach algebra</i> . . . . .	1
David W. Boyd, <i>Inequalities for positive integral operators</i> . . . . .	9
Lawrence Gerald Brown, <i>Note on the open mapping theorem</i> . . . . .	25
Stephen Daniel Comer, <i>Representations by algebras of sections over Boolean spaces</i> . . . . .	29
John R. Edwards and Stanley G. Wayment, <i>On the nonequivalence of conservative Hausdorff methods and Hausdorff moment sequences</i> . . . . .	39
P. D. T. A. Elliott, <i>On the limiting distribution of additive functions (mod 1)</i> . . . . .	49
Mary Rodriguez Embry, <i>Classifying special operators by means of subsets associated with the numerical range</i> . . . . .	61
Darald Joe Hartfiel, <i>Counterexamples to a conjecture of G. N. de Oliveira</i> . . . . .	67
C. Ward Henson, <i>A family of countable homogeneous graphs</i> . . . . .	69
Satoru Igari and Shigehiko Kuratsubo, <i>A sufficient condition for <math>L^p</math>-multipliers</i> . . . . .	85
William A. Kirk, <i>Fixed point theorems for nonlinear nonexpansive and generalized contraction mappings</i> . . . . .	89
Erwin Kleinfeld, <i>A generalization of commutative and associative rings</i> . . . . .	95
D. B. Lahiri, <i>Some restricted partition functions. Congruences modulo 11</i> . . . . .	103
T. Y. Lin, <i>Homological algebra of stable homotopy ring <math>\pi_*</math> of spheres</i> . . . . .	117
Morris Marden, <i>A representation for the logarithmic derivative of a meromorphic function</i> . . . . .	145
John Charles Nichols and James C. Smith, <i>Examples concerning sum properties for metric-dependent dimension functions</i> . . . . .	151
Asit Baran Raha, <i>On completely Hausdorff-completion of a completely Hausdorff space</i> . . . . .	161
M. Rajagopalan and Bertram Manuel Schreiber, <i>Ergodic automorphisms and affine transformations of locally compact groups</i> . . . . .	167
N. V. Rao and Ashoke Kumar Roy, <i>Linear isometries of some function spaces</i> . . . . .	177
William Francis Reynolds, <i>Blocks and <math>F</math>-class algebras of finite groups</i> . . . . .	193
Richard Rochberg, <i>Which linear maps of the disk algebra are multiplicative</i> . . . . .	207
Gary Sampson, <i>Sharp estimates of convolution transforms in terms of decreasing functions</i> . . . . .	213
Stephen Scheinberg, <i>Fatou's lemma in normed linear spaces</i> . . . . .	233
Ken Shaw, <i>Whittaker constants for entire functions of several complex variables</i> . . . . .	239
James DeWitt Stein, <i>Two uniform boundedness theorems</i> . . . . .	251
Li Pi Su, <i>Homomorphisms of near-rings of continuous functions</i> . . . . .	261
Stephen Willard, <i>Functionally compact spaces, <math>C</math>-compact spaces and mappings of minimal Hausdorff spaces</i> . . . . .	267
James Patrick Williams, <i>On the range of a derivation</i> . . . . .	273