

Pacific Journal of Mathematics

**HOMOLOGICAL ALGEBRA OF STABLE HOMOTOPY RING π_*
OF SPHERES**

T. Y. LIN

HOMOLOGICAL ALGEBRA OF STABLE HOMOTOPY RING π_* OF SPHERES

T. Y. LIN

The stable homotopy groups are studied as a graded ring π_* via homological algebra. The main object is to show that the projective (and weak) dimension of a finite type π_* -module is ∞ unless the module is free. As a corollary, a partial answer to Whithead's corollary to Freyd's generating hypothesis is obtained.

1. Introduction and statement of main results. It is well-known that the stable homotopy groups of spheres form a commutative graded ring π_* [20]. This paper is our first effort towards the investigation of the homological properties of the stable homotopy ring π_* . In this paper we have completed the computations of all the homological numerical invariants of finitely generated type. The nonfinitely generated type will be taken up in forth-coming papers.

The paper is organized as follows: The introduction is § 1. In § 2 we give a brief exposition about the theory of graded rings, which are needed in later sections. § 3 is primarily a preparation for § 4. Here the finitistic global dimensions of the " p -primary component" A_p of π_* (precisely, A_p is a ring obtained by localizing π_* at a maximal ideal) are computed, and a geometric realization of A_p is constructed. § 4 is the mainbody of this paper, here we prove Theorem 2 and derive from it Theorems 1, 3, and 4. We would like to suggest that the reader, after § 1, go directly to § 4 and refer to the rest of the sections when necessary.

The author would like to thank Professor W. Massey for his encouragement and fruitful discussions.

The main results are:

THEOREM 1. *Let A be a locally finitely generated (i.e., there are only finitely many generators at each degree) π_* -module having finite projective, as well as weak, dimension. Then A is a free π_* -module and hence is realizable as a stable homotopy module $\pi_*(Y)$ by a wedge Y of spheres.*

In [7] Freyd propose a conjecture, known as the generating hypothesis, which asserts that a map between finite CW -complexes, which induces the zero map on stable homotopy groups, is stably null-homotopic. A consequence of this conjecture, due to G. Whithead, asserts that the finitely generated stable homotopy module $\pi_*(X)$ of

a finite spectrurum is a stable homotopy module of a wedge of spheres. Theorem 1 implies that a finitely generated stable homotopy module of finite projective dimension is a stable homotopy module of a wedge of spheres. Thus Theopem 1 proves “Whithead’s corollary to Freyd’s generating hypothesis” with one additional condition.

THEOREM 2. *Let Y be a connected spectrum having a finite skeleton in each dimension. Then the projective, as well as weak, dimension of $\pi_*(Y)$ as a π_* -module is infinite unless Y is a wedge of spheres.*

In the notation of § 2, 3 below, we have.

THEOREM 3. (i) $f.P.D(\pi_*) = f\angle.P.D(\pi_*) = 0 < F.P.D(\pi_*) \neq 0$
(ii) $f.W.D(\pi_*) = f\angle.W.D(\pi_*) = 0 \leq F.W.D(\pi_*)$.

It is well known that the projective dimension of a module over the Steenrod Algebra (mod p) is either 0 or ∞ . Theorem 3 implies that an analogous situation also occurs in the stable homotopy ring, when a finiteness condition is imposed. For $f\angle.P.D(\pi^*) = 0$ means that the projective dimension of a locally finitely generated π_* -module is either 0 or ∞ . Note that the finiteness condition cannot be removed. For $F.P.D(\pi_*) > 0$ implies that there are some π_* -modules having projective dimension ≥ 1 . In fact, in [14] we show that the only projective dimension of π_* -module is either 0, 1, or ∞ , equivalently $F.P.D(\pi_*) = 1$.

THEOREM 6. *The projective, as well as weak, dimension of an ideal in π_* is infinite.*

This theorem shows that π_* is a very nonhereditary ring.

Since we are working with the stable theory, a good category will be helpful. Therefore we will freely use Boardman’s stable category of (pointed) CW -spectra. [6].

CONVENTION:

- (i) All the spectra X are assumed to be connected, i.e., there is some integer n_0 such that $\pi_n(X) = 0$ for $n < n_0$.
- (ii) All the modules are assumed to be bounded from below, i.e. there is some integer n_0 such that $A_n = 0$ for all $n < n_0$; see § 2.3.
- (iii) We will not distinguish between a map or its homotopy class and a spectrum or its homotopy type.

2. Graded rings. The purpose of this section is to give a brief

exposition about the theory of graded rings, which are needed in later sections. First, we define a graded ring and set up the category of its modules; also we recall here a few terms and facts about homological algebra. Secondly, we briefly discuss a graded version of the theory of Noetherian rings: Since the finitely generated condition is too strong for homotopy [10], we relax the finiteness condition and arrive at a generalization-called a locally Noetherian ring [16]. Then we prove some folk theorems about such rings. These theorems are standard in the ordinary Noetherian theory of rings, yet the graded versions seem never to have appeared in print. Thirdly, we review some facts about the localization. Finally we introduce an operation on graded modules—the lifting of a module. This operation gives us a clearer view of graded free modules, and implies clearly the realizability of free π_* -modules and mappings between them.

I. First let us recall some definitions.

DEFINITIONS 2.1. A graded ring R is a sequence of abelian groups R_n for all *NONNEGATIVE* integers $n \geq 0$ together with homomorphisms $R_n \otimes_z R_m \rightarrow R_{n+m}$ (which we call multiplication) satisfying associativity and distributivity. We always assume the existence of a multiplicative identity element 1 in R_0 .

A graded left R -module A is a sequence of abelian groups A_n for *ALL* integers n , together with homomorphisms $R_n \otimes_z A_m \rightarrow A_{n+m}$ (which we call scalar multiplication) satisfying the usual module identities.

A graded ring is said to be commutative (in the graded sense) if $\alpha \cdot \beta = (-1)^{n \cdot m} \beta \cdot \alpha$ for $\alpha \in R_n$ and $\beta \in R_m$. The integer n will be called the degree of α and denoted by $\deg \alpha$.

From now on, all rings will be assumed to be commutative (in the graded sense); all modules, ideals and properties will be assumed to be two sided unless they are specifically called otherwise. All modules, rings and ideals are graded.

An R -module map $f: A \rightarrow B$ of degree d is a sequence of maps $f: A_n \rightarrow B_{n+d}$ such that for $\alpha \in R_n$ and $x \in A_m$ $f(\alpha x) = (-1)^{n \cdot d} \alpha f(x)$.

Topological Examples. 2.2. Let $\pi_n = \pi_n(S^0)$ be the n th stable homotopy group of the 0-sphere spectrum. Then the collection $\pi_* = \{\pi_n, n \geq 0\}$, with composition of maps as multiplication, is a graded commutative ring [20].

Let X be a spectrum, then the collection of stable homotopy groups $\pi_*(X) = \{\pi_n(X), -\infty < n < \infty\}$ is a graded π_* -module with composition of maps as scalar multiplication. Similarly a stable cohomotopy group $\pi^*(X) = \{\pi^n(X), -\infty < n < \infty\}$ is a graded left π_* -module. Moreover, since π_* is commutative, a right π_* -module can be changed into a left π_* -module by standard sign convention. Thus throughout the whole paper we will omit the term left or right, and leave the reader to interpret it properly.

Next we will recall some notions from homological algebra.

DEFINITIONS. 2.3. Let $\mathcal{M}(R)$ be the category of R -modules bounded from below; that is, (i) the objects are the R -modules $A = \{A_n; -\infty < n < \infty\}$ such that there is an integer n_0 with $A_n = 0$ for all $n < n_0$, and (ii) the morphisms are R -module maps of degree 0. Let $P.d_R A$ and $W.d_R A$ be the projective and weak dimensions, respectively, of the R -module A in the category $\mathcal{M}(R)$ (Note that the notions of projective and weak dimension are categorical in nature, so it is important to require that everything is *INSIDE THE CATEGORY* $\mathcal{M}(R)$). Following [3, pp. 478] or [9, pp. 48] we write

$$\begin{aligned} G.D(R) &= \text{Sup}\{P.d_R A \mid A \in \mathcal{M}(R)\} \\ F.P.D(R) &= \text{Sup}\{P.d_R A \mid A \in \mathcal{M}(R), P.d_R A < \infty\} \\ f\angle.P.D(R) &= \text{Sup}\{P.d_R A \mid A \in \mathcal{M}(R), P.d_R A < \infty \\ &\text{and } A \text{ is locally finitely generated}\} \end{aligned}$$

$$f.P.D(R) = \text{Sup}\{P.d_R A \mid A \in \mathcal{M}(R), P.d_R A < \infty$$

and A is finitely generated}, where “Sup” stands for the supremum, and “locally finitely generated” means that there is a set of generators which consists of only *FINITELY MANY* generators *EACH DEGREE*. (see 2.6 below). Similarly we can define $F.W.D(R)$, $f\angle.W.D(R)$ and $f.W.D(R)$ by replacing $P.d_R A$ with $W.d_R A$. (The notations $G.D$, $F.P.D$, $f.P.D$ are standard, while $f\angle.P.D$ is introduced here; the “ $f\angle$ ” stands for “finite dimension and locally finitely generated”).

Let us recall

PROPOSITION 2.4. *Let $0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0$ be a short exact sequence of R -modules. Then if A is projective and A'' is not, $P.d_R A' = P.d_R A'' - 1$.*

COROLLARY 2.5. *$F.P.D(R) \neq 0$ (or $f\angle.P.D(R) \neq 0$) implies that there is a module A with $P.d_R A = 1$.*

Proof. By definition $F.P.D(R) \neq 0$ (or $f\angle.P.D(R) \neq 0$) implies that

there is a module, say B , such that $P.d_R B = n < \infty$. Let

$$0 \leftarrow B \leftarrow R_0 \leftarrow P_1 \leftarrow \cdots \leftarrow P_n \leftarrow 0$$

be a projective resolution. Then we can break the resolution into short exact sequences

$$\begin{aligned} 0 \leftarrow B \leftarrow P_0 \leftarrow K_0 \leftarrow 0 \\ 0 \leftarrow K_i \leftarrow P_{i+1} \leftarrow K_{i+1} \leftarrow 0, \quad 0 \leq i \leq n-2 \\ 0 \leftarrow K_{n-2} \leftarrow P_{n-1} \leftarrow p_n \leftarrow 0. \end{aligned}$$

By Proposition 2.4 we get $P.d_R K_0 = 1$. Taking $A = K_0$ finishes the proof.

II. We also need a graded version of Noetherian rings.

DEFINITIONS. 2.6. An R -module A is said to be locally finitely generated if there is a set X of generators such that (i) $X \cap A_n$ is a finite set and (ii) there exists an integer n_0 such that $X \cap A_m = \emptyset$ for $m < n_0$. (Note that if $A \in \mathcal{M}(R)$, i.e., A is bounded from below, then condition (ii) is automatically satisfied)

An R -module A is said to be locally Noetherian if every submodule is locally finitely generated. A ring is locally Noetherian if it is locally Noetherian as a module.

There is a simple criterion for locally Noetherianness

PROPOSITION 2.7. R is locally Noetherian if and only if R_0 is an ungraded Noetherian ring and each R_n is a finitely generated R_0 -module.

Proof. To prove necessity, let I_0 be an ideal of R_0 . Then $I = I_0 \cup (\cup_{n \geq 1} R_n)$ is an ideal of R and hence, by assumption, is locally finitely generated; therefore I_0 and each R_n , $n \geq 1$ are finitely generated. Thus we establish the necessity. Sufficiency holds because if I is an ideal of R , then, by locally Noetherianness, each I_n is a finitely generated R_0 -module.

Topological example 2.8. The stable homotopy ring π_* is a locally Noetherian ring (by Serre's theorem and 2.7). The stable homotopy module is locally Noetherian if X is a connected spectrum having a finite skeleton in each dimension. (see 2.9 (i) below).

Most of the (ordinary) Noetherian theorems can easily be rephrased to get valid graded versions. The following proposition is a graded

version of a standard Noetherian theorem.

PROPOSITION 2.9. (i) *A locally finitely generated module over a locally Noetherian ring is locally Noetherian.* (ii) *A locally finitely generated flat module of a locally Noetherian ring is projective.*

The author learned this proposition from P. May in a course on homological algebra [16]. Since the proposition seems not to appear in print, we reproduce the proof here.

Proof. (i) The proof is almost the same as in the ungraded case. First let us note that a quotient of a locally Noetherian module is locally Noetherian; also let us observe that a locally finitely generated free module of a locally Noetherian ring is locally Noetherian. Then, since every locally finitely generated R -module A is a quotient of a locally finitely generated free R -module, so the module A is locally Noetherian.

(ii) Let A be a locally finitely generated R -module, then there is a locally finitely generated free R -module F such that

$$0 \rightarrow K \rightarrow F \rightarrow A \rightarrow 0$$

is exact. Since R is locally Noetherian, so by (i) F is locally Noetherian; hence K is locally finitely generated. Thus, in order to prove that A is projective, it is sufficient to prove the following Lemma.

LEMMA 2.10. *Let $0 \rightarrow K \xrightarrow{\psi} F \rightarrow A \rightarrow 0$ be a short exact sequence of R -modules, where A is flat, F is free and K is locally finitely generated. Then the sequence splits, hence A is projective.*

Proof. Without loss of generality, we can assume, since A , F and K are bounded from below, that A , F and K have no negative degrees (i.e. $F_i = K_i = A_i = 0$ for $i < 0$). We will prove this lemma by constructing $\phi: F \rightarrow K$ $\ni \phi\psi = 1$: Let $F(q)$ be the submodule generated by F_i , $i \leq q$ and define $K(q)$ similarly. Observe that $\psi(K(q)) \subset F(q)$. We will define $\phi(q): F(q) \rightarrow K(q)$ inductively to satisfy $\phi(q)\psi = 1$ on $K(q)$. We start with $\phi(-1) = 0$; we suppose that $n \geq 0$ and $\phi(n-1)$ has been defined. In order to construct $\phi(n)$, first we remark that if $x \in K_n$, then there exists a map $\lambda: F(n) \rightarrow K(n)$ such that (i) $\lambda\psi = 1$ on $K(n-1) + Rx$ and (ii) $\lambda = \phi(n-1)$ on $F(n-1)$. To prove this remark, let $\{y_j\}$ be a basis for F and suppose that $\psi(x) = \sum_h \gamma_{jh} y_{jh}$, where $\gamma_{jh} \in R$ and $y_{jh} \in \{y_j\}$. Let I be the left ideal generated by $\{\gamma_{jh}\}$. Then, since A is flat, we

have $IF \cap K = IK$ (see the Sublemma 2.11 below). So $x = \sum \gamma_{j_h} \cdot k_{j_h}$ for some $k_{j_h} \in K$. If $\gamma_{j_h} \cdot k \in K(n-1)$, then $\gamma_{j_h} \cdot k_{j_h} = \phi(n-1)\psi(\gamma_{j_h}k_{j_h})$. Define λ by

$$\lambda = \phi(n-1) \text{ on } F(n-1) ;$$

$\lambda(y_{j_h}) = k_{j_h}$, if $y_{j_h} \in F_n$, $\gamma_{j_h} \in R_0$ and $\gamma_{j_h} \neq 0$; and $\lambda(y_j) = 0$, if $y_j \in F_n$ and does not appear in $\{y_{j_h}\}$. Then $\lambda\psi(x) = x$, and this proves our remark.

Proceeding inductively, suppose that $\lambda_{\ell-1}: F(n) \rightarrow K(n)$ can be constructed so as to satisfy

$$\begin{aligned} \lambda_{\ell-1} &= \phi(n-1) \text{ on } F(n-1); \text{ and for any } x_1, \dots, x_{\ell-1} \in K_n, \\ \lambda_{\ell-1} \cdot \psi &= 1 \text{ on } K(n-1) + Rx_1 + \dots + Rx_{\ell-1}. \end{aligned}$$

Suppose given $x_1, x_2, \dots, x_{\ell} \in K_n$. Construct λ as above for $K(n-1) + Rx_{\ell}$ and construct $\lambda_{\ell-1}$, as assumed possible, for $K(n-1) + Rx_1 + \dots + Rx_{\ell-1}$, where $w_i = x_i - \lambda\psi(x_i)$. Now define $\lambda_{\ell} = \lambda_{\ell-1} + \lambda - \lambda_{\ell-1}\psi\lambda$. Then

$$\lambda_{\ell} = \phi(n-1) \text{ on } F(n-1) ;$$

and

$$\lambda_{\ell}\psi(x_i) = x_i \quad 0 \leq i \leq \ell.$$

Since K is locally finitely generated, it follows that we can construct $\phi(n)$. Since $F = \bigcup_n F(n)$, we can find $\phi: F \rightarrow K$ such that $\phi\psi = 1$. This completes the Lemma.

The only thing left is to show

SUBLEMMA 2.11. *The following are equivalent*

- (i) *A is a flat R-module*
- (ii) *For every ideal I of R, and for every short exact sequence $0 \rightarrow K \rightarrow F \rightarrow A \rightarrow 0$ with F free, we have $IF \cap K = IK$.*

Proof. Consider the following commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Tor}_1^R(A, R/I) & \longrightarrow & K \otimes_R R/I & \longrightarrow & F \otimes_R R/I \longrightarrow A \otimes_R R/I \longrightarrow 0 \\ & & & & \downarrow \cong & & \downarrow \cong \\ 0 & \longrightarrow & \frac{K \cap IF}{IK} & \longrightarrow & \frac{K}{IK} & \longrightarrow & \frac{F}{IF} \longrightarrow \frac{A}{IA} \longrightarrow 0. \end{array}$$

Note that the top horizontal line is exact since it is a part of a long exact sequence of the functor Tor ; and the bottom line is easily

seen to be exact. The right three vertical arrows are the natural isomorphisms induced by scalar multiplication. Hence

$$\frac{K \cap IF}{IK} \cong \text{Tor}_1^R(A, R/I).$$

Thus

$$\frac{K \cap IF}{IK} \cong \text{Tor}_1^R(A, R/I) = 0$$

if and only if A is flat. Or equivalently, $K \cap IF = IK$ if and only if A is flat.

The next propositions are also standard in Noetherian theory (though the ring need not be locally Noetherian, the locally finiteness conditions are essential).

PROPOSITION 2.12. (graded Nakayama Lemma)

The following conditions on an ideal I in R are equivalent

- (i) $I \subset J = \text{the Jacobson radical of } R$.
- (ii) *Let A be a locally finitely generated R -module and H be a submodule of A . Then $H + IA = A$ implies that $H = A$.*
- (iii) *Let A be a locally finitely generated R -module. Then $IA = A$ implies that $A = 0$.*

Proof. Before we start to prove the equivalence, let us first observe that every proper submodule H of a locally finitely generated R -module A is always contained in a proper maximal submodule: Suppose $H \neq A = \{A_n; -\infty < n < \infty\}$. Then we will construct a proper maximal submodule $G \ni H \subset G \subsetneq A$. Let $m-1$ be the highest degree such that $H_{m-1} = A_{m-1}$. Since A is locally finitely generated, A_m is finitely generated over R_0 (= the 0th component of R). Then, by Zorn's Lemma, we can find a maximal proper R_0 -submodule E (of A_m) containing H_m . Let G be an R -submodule defined by

$$G_n = \begin{cases} A_n, & \text{if } n \neq m \\ E, & \text{if } n = m. \end{cases}$$

It is not difficult to see that G is, indeed, an R -submodule (Let $H^{(m-1)}$ be the subset consisting of all H^n , $n \leq m-1$. Then, by construction, $H^{(m-1)} = G^{(m-1)} = A^{(m-1)}$. Since $R \cdot H^{(m-1)} \subset H$, we have $R \cdot G^{(m-1)} \subset H \subset G$. Therefore $R \cdot G \subset G$. Thus G is, indeed, a submodule). The maximality of G (as R -submodule) follows immediately from the maximality of E as R_0 -submodule of A_m .

Now with this observation, we are ready for proving the

equivalences.

(i) \Rightarrow (ii). First we note that $J \cdot A \subset J(A)$ = the intersection of all maximal submodules (Since $J(A)$ is the set of elements of A belonging to the kernels of all homomorphisms of A into simple R -modules, and J annihilates all simple R -modules). Next let us assume, to the contrary, that $H \neq A$. Since A is locally finitely generated, by the previous remarks, there is a proper maximal R -submodule G containing H . Since $I \cdot A \subset J \cdot A \subset J(A)$, we have $I \cdot A \subset G$. Therefore $IA + H \subset G \neq A$; contradiction. This establishes the implication (i) \Rightarrow (ii).

(ii) \Rightarrow (iii). Take $H = 0$ in (ii); we get the conclusion.

(iii) \Rightarrow (i). If B is a maximal ideal, then, by (iii), $I \cdot (R/B) = 0$. In other words, $B \supset I \cdot R = I$. Since B is arbitrary, $I \subset J$ = the Jacobson radical of R = the intersection of all the maximal ideals.

PROPOSITION 2.13. *Let $C(R) \subseteq \mathcal{M}(R)$ be the full subcategory of locally finitely generated projective R -modules. Let I be an ideal of R contained in the Jacobson radical of R . Set $R' = R/I$ and write $A' = A \otimes_R R' = A/IA$ for $A \in \mathcal{M}(R)$. Then*

$$': C(R) \longrightarrow C(R')$$

is a full additive functor satisfying the following property: If $f: P \rightarrow Q$ is a morphism in $C(R)$ such that $f': P' \rightarrow Q'$ is an isomorphism in $C(R')$, then f is an isomorphism. (This proposition is the graded version of [5, proposition 2.12, p. 90] and the proof is essentially the same.)

Proof. Given $f': P' \rightarrow Q'$ there is an $f: P \rightarrow Q$ making the diagram

$$\begin{array}{ccc} P & \xrightarrow{f} & Q \\ \downarrow & & \downarrow \\ P' & \xrightarrow{f'} & Q' \end{array}$$

commute. This is because P is projective and $Q \rightarrow Q'$ is onto. Thus the functor $'$ is full. If f' is onto, then it follows from the Nakayama Lemma 2.12 (ii) (since Q is locally finitely generated) that f is onto. The projectivity of Q now implies that f is a split epimorphism. It follows that $H = \text{Ker } f$, being a direct summand of P , is locally finitely generated. Now since $\text{Ker } f' = 0$, $H' = 0$; or equivalently $H/I \cdot H = 0$. By the Nakayama Lemma 2.12 (iii), we get $H = 0$. This proves the proposition.

COROLLARY 2.14. (i) *A locally finitely generated projective (or*

flat) π_* -module A is free.

(ii) A locally finitely generated projective (or flat) module A over a local ring (= locally Noetherian ring with unique maximal ideal) is free.

Proof. (i) Observe that the Jacobson radical of π_* is $\pi_+ = \{\pi_n, n > 0\}$. Take $I = \pi_+$ as above. Then $\pi'_* = \pi_*/\pi_+ \cong \pi_0$ is isomorphic to the ring Z of integers, so $A' \in C(Z)$ is free. By Proposition 2.13, A is free.

(ii) Take I = the Jacobson radical (= the unique maximal ideal) above. Then R' is field, so A' is free. By Proposition 2.13, A is free.

THEOREM 2.15. (i) If $A \in \mathcal{M}(R)$ and $P.d_R A = n < \infty$, then there is a free resolution of length n , i.e., the sequence

$$0 \longleftarrow A \longleftarrow F_0 \longleftarrow F_1 \longleftarrow \cdots \longleftarrow F_n \longleftarrow 0$$

is exact and each F_i is free.

(ii) Let R be the stable homotopy ring π_* or an arbitrary local ring; and let A be a locally finitely generated module. Then $P.d_R A = n < \infty$ implies that there is a locally finitely generated free resolution of length n , i.e., the sequence

$$0 \longleftarrow A \longleftarrow F_0 \longleftarrow F_1 \longleftarrow \cdots \longleftarrow F_n \longleftarrow 0$$

is exact and each F_i is a locally finitely generated free R -module.

Proof. (i) By assumption $P.d_R A = n < \infty$, there is an exact sequence

$$0 \longleftarrow A \longleftarrow F_0 \longleftarrow F_1 \longleftarrow \cdots \xleftarrow{d_{n-1}} F_{n-1} \xleftarrow{d_n} K_n \longleftarrow 0$$

such that F_i are free and K_n is projective. By the Eilenberg Lemma [4, pp. 24], there is a very large free module, say F , such that $K_n \oplus F$ is a free module. Then the following sequence

$$0 \longleftarrow A \longleftarrow F_0 \longleftarrow \cdots \xleftarrow{d'_{n-1}} (F_{n-1} \oplus F) \xleftarrow{d'_n} (K_n \oplus F) \longleftarrow 0$$

is a free resolution for A , where $d'_n = d_n \oplus 1_F$ (1_F is the identity map on F), $d'_{n-1} = d_{n-1} \oplus 0$ (0 is the zero map $F \rightarrow F_{n-2}$); and $d'_i = d_i$, $0 \leq i \leq n-2$. This proves (i).

(ii) By Corollary 2.14, any projective resolution is a free resolution. The proof that the resolution is locally finitely generated is similar to the ordinary Noetherian case.

III. The localization can also be carried over to the graded commutative ring R . (Note that since the ring R is not strictly

commutative, some special care has to be taken [18]). For simplicity we will assume that the elements of the multiplicative system S (containing 1 but not containing 0) are of degree 0, i.e. $S \subset R_0$. Let $R \otimes S$ be the Cartesian product, then we will write

$$(a, s) \sim (a', s') \quad \text{if} \quad \exists t \in S \ni t \cdot (as' - a's) = 0$$

where (a, s) and (a', s') are elements of $R \times S$. It is easy to verify that this is, in fact, an equivalence relation. Let the equivalence class to which (a, s) belongs be denoted by a/s and be assigned a degree, $\deg a/s = \deg a$. We will denote by R_s the totality of the equivalence classes. As in the ordinary case, we can make R_s into a ring by introducing the sum and product with the usual formulas. Moreover there is a natural map

$$R \longrightarrow R_s$$

which sends a to $a/1$ (Note that this map, in general, is not a monomorphism). We will call this map a natural embedding (abuse of language). With this natural embedding we can turn R_s into an R -module.

We also can consider the localization of a module. Let A be an R -module, then the localized module is defined to be $A_s = A \otimes_s R_s$.

PROPOSITION 2.16. *If A is a flat R -module, then A_s is flat R -module, in particular R_s is a flat R -module.*

The proof of this proposition can be found in [17, pp. 170, Lemma 2].

IV. Finally we recall

DEFINITION 2.17. Let $A = \{A_n, -\infty < n < \infty\} \in \mathcal{M}(R)$. Let $LA = \{(LA)_n, -\infty < n < \infty\}$ to be the R -module such that (i) $(LA)_n = A_{n-1}$ as Abelian group and (ii) $\gamma \cdot La = (-1)^{\deg \gamma} L\gamma a$, where $\gamma \in R$ and $a \in A$. Inductively we define $L_{\angle} A = L(L_{\angle^{-1}} A)$.

It is easy to see that $L_{\angle} R$ is a free R -module with one single generator $L_{\angle} 1$ (1 is the identity in R) at degree \angle . Note also that a free R -module F is a direct sum of the form $L_{\angle} R$ i.e., $F \cong \sum_i \oplus (L_{\angle_i} R)$.

In the case π_* is the ring, then $L_{\angle_i} \pi_*$ is just the stable homotopy module $\pi_*(S^{\angle_i})$ of the \angle_i -sphere. Thus we have a theorem analogous to [2, § 3, Theorem 3], namely,

THEOREM 2.18. *Let $F \in \mathcal{M}(\pi_*)$ be a locally finitely generated*

free π_* -module. Then there exists a wedge N of spheres, which has only finitely many spheres at each dimension, such that

- (i) $\pi_*(N) \cong F$ as π_* -module; and
- (ii) $\pi_0(N, X) \cong \text{hom}_{\pi_*}(F, \pi_*(X))$ for any spectrum X , where $\pi_0(N, X)$ is the homotopy classes of maps from N to X , and $\text{hom}_{\pi_*}(F, \pi_*(X))$ is the homomorphism of degree 0 from F to $\pi_*(X)$.

3. The \mathfrak{p} -primary component of π_* . Identify π_0 with ring Z of integers. Let M_p be the maximal ideal of π_* generated by the prime p and π_+ , where π_+ are the elements of degree ≥ 1 . The complement of M_p is obviously a multiplicative system of degree 0; hence, by §1. III, upon localizing at this multiplicative system we get a local ring A_p which, as π_* -module, is flat. The main objects of this section are (i) to compute the finitistic global dimension of A_p ; and (ii) to realize A_p as a stable homotopy module of some spectrum L_p .

I. The finitistic global dimension of A_p .

Before we state the main theorem, recall the theorem of Kervaire-Milnor [11].

THEOREM 3.1. *For each prime p and positive integer j , there exists $\alpha \in A_p$ such that $p^j \cdot \alpha = 0$ and $p^{j-1} \cdot \alpha \neq 0$.*

Let the notation be as in Definition 2.3. Then our main theorem in this section is the following

THEOREM 3.2. (i) $f.P.D(A_p) = f\angle.P.D(A_p) = 0 < F.P.D(A_p) \neq 0$
(ii) $f.W.D(A_p) = f\angle.W.D(A_p) = 0 \leq F.W.D(A_p)$.

This theorem shows that the conjecture stated in [9, pp. 62, (2)] should exclude the case $f.P.D = 0$.

The proof of this theorem follows that of Lemma 3.3 and 3.4 below.

LEMMA 3.3. $f.P.D(A_p) = f\angle.P.D(A_p) = 0$.

Proof. By definition, it suffices to prove $f\angle.P.D(A_p) = 0$. In order to prove this, let us assume, to the contrary, that $f\angle.P.D(A_p) > 0$. Then from Corollary 2.5 there is a locally finitely generated A_p -module A of projective dimension one. Moreover since A_p is a (locally Noetherian) local ring, by Theorem 2.15. (ii), we have a free resolution

$$0 \longleftarrow A \xleftarrow{\epsilon} F \longleftarrow K \longleftarrow 0$$

with K and F being locally finitely generated. Also we can choose K and F so that $K \subset M \cdot F$, where M is the Jacobson radical (= the unique maximal ideal) of A_p . For, let $\{a_i\}$ be a minimal generating system of A , let F be a free module with generators $\{x_i\}$ (where $\deg x_i = \deg a_i$), and let ε be a map sending x_i to a_i . Now $\sum \alpha_i x_i$ is in K if and only if $\sum \alpha_i a_i = 0$. Suppose α_1 is not in M . Then α_1 is a unit. Thus $a_1 = -\alpha_1^{-1}(\sum_{i>1} \alpha_i a_i)$, which contradicts the minimality of $\{a_i\}$. Hence α_1 and similarly α_i must be in M . Therefore K is contained in $M \cdot F$ which is a direct sum $\bigoplus_{i=1}^n \{M_i \mid M_i = M\}$. Since K is free, then $K \subset M \cdot F$ implies that $L_{\not\sim} A_p \subset \bigoplus_{i=1}^n \{M_i \mid M_i = M\}$, where $L_{\not\sim} A_p$ is the ring A_p with degree lifted by $\not\sim$ (see Definition 2.17). Then we have

$$L_{\not\sim} 1 = (m_1, m_2, \dots, m_n), \quad m_i \in M.$$

This implies that the annihilator of $m_i (i = 1, 2, \dots, n)$ is also an annihilator of $L_{\not\sim} 1$, and therefore is zero. We will now show that the annihilator of $\{m_i\}$ is not zero.

Let m_{i_1}, \dots, m_{i_s} be those m_i with degree ≥ 1 . Let p^j be the maximum additive order of m_{i_1}, \dots, m_{i_s} . From Theorem 3.1, there is $\beta \in A_p$ such that $p^j \cdot \beta \neq 0$ and $p^{j+1} \cdot \beta = 0$. Write $p^j \cdot \beta = \beta_0$. Then obviously $\beta_0 \cdot m_{i_k} = \beta \cdot p^j \cdot m_{i_k} = 0$, for $k = 1, \dots, s$, and $p \cdot \beta_0 = 0$. Note that the elements of degree 0 in the Jacobson radical M of A_p are of the form tp/q , where p and q are relatively prime. Thus $\beta_0 \cdot m_j = \beta_0 \cdot tp/q = t/q \cdot p \cdot \beta_0 = 0$ for $j \neq i_1, \dots, i_s$. This shows that $\beta_0 \neq 0$ is a nonzero annihilator of m_i , $i = 1, \dots, n$. Hence the annihilator of $L_{\not\sim} 1$ is nonzero; contradiction. This concludes the proof of the lemma.

To prove the strict inequality of the theorem, we prove

LEMMA 3.4. *Let $R = \{R_n, n \geq 0\}$ be a nonnegatively graded commutative ring. Then $F.P.D(R) = 0$ implies that R_0 , the 0th component of R , is an ungraded perfect ring.*

Proof. From the proof of [3, Lemma 1.3], we see that $F.P.D(R) = 0$ implies that every decreasing sequence of principal ideals $\alpha_1 R \supseteq \alpha_1 \alpha_2 R \supseteq \dots \supseteq \alpha_1 \dots \alpha_n R \supseteq \dots$ is finite, ($\deg \alpha_i = 0$). But observe that since we have chosen $\deg \alpha_i = 0$, every principal ideal $\alpha_1 \dots \alpha_n R$ of R gives rise to a principal $\alpha_1 \dots \alpha_n R_0$ of R_0 , and vice versa. Thus the finiteness of the decreasing sequence $\{\alpha_1 \dots \alpha_n R\}$ implies that of $\{\alpha_1 \dots \alpha_n R_0\}$. Then, by [3, Theorem P], R_0 is an ungraded perfect ring.

Proof of Theorem 3.2. The 0th component of A_p is just the local ring $Z(p)$ obtained by localizing the ring Z of the integers at the

prime ideal pZ , and $Z(p)$ is obviously not a perfect ring. Thus we conclude from Lemma 3.4 that $F.P.D(A_p) \neq 0$. Since A_p is locally Noetherian, by Proposition 2.9. (ii),

$$f \not\leq W.D(A_p) = f.W.D(A_p) = f \not\leq P.D(A_p) = 0 ,$$

while from its definition $F.W.D(A_p) \geq 0$. Thus we prove the theorem.

II. Geometric realization.

The purpose of this section is to prove the following theorem.

THEOREM 3.5. *There exists a spectrum L_p such that $\pi_*(L_p) = A_p$ as a π_* -module.*

The proof of this theorem is broken into lemmas.

Let $q_1, q_2, \dots, q_i, \dots$ be all the prime integers except p . Let Q_i be the multiplicative system $Q_i = \{1, q_i, q_i^2, \dots, q_i^n, \dots\}$, and (Q_1, \dots, Q_n) be a multiplicative system generated by Q_i , $i = 1, 2, \dots, n$. Since we have identified the ring Z of integers with π_0 , each Q_i and hence (Q_1, \dots, Q_n) are multiplicative systems (of degree 0) in π_* . Hence, by § 1. III, we have a sequence of local rings i.e.,

$$\begin{aligned} \pi_*(Q_1) & \text{ is } \pi_* \text{ localized at } Q_1 \\ \pi_*(Q_1, Q_2) & \text{ is } \pi_* \text{ localized at } (Q_1, Q_2) \\ & \vdots \\ \pi_*(Q_1, Q_2, \dots, Q_n) & \text{ is } \pi_* \text{ localized at } (Q_1, Q_2, \dots, Q_n) . \end{aligned}$$

Note that the ring $Z = \pi_0$ is still naturally embedded in each $\pi_*(Q_1, Q_2, \dots, Q_n)$. Thus it is legitimate to regard, say Q_{n+1} , as a multiplicative system (of degree 0) in $\pi_*(Q_1, Q_2, \dots, Q_n)$. Hence we have the following lemma:

LEMMA 3.6. *$\pi_*(Q_1, Q_2, \dots, Q_{n+1})$ is the local ring obtained by localizing $\pi_*(Q_1, \dots, Q_n)$ at Q_{n+1} , and hence*

$$(1) \quad \pi_*(Q_1, \dots, Q_{n+1}) = \lim_{\rightarrow} \pi_*(Q_1, \dots, Q_n)$$

where the direct limit “ \lim_{\rightarrow} ” is taken over the direct system:

$$\pi_*(Q_1, \dots, Q_n) \xrightarrow{q_{n+1}} \pi_*(Q_1, \dots, Q_n) \xrightarrow{q_{n+1}} \dots ,$$

in which q_{n+1} is multiplication by the prime q_{n+1} .

Note also that there are natural “embedding” arising from

localizations (see § 1. III)

$$f_n: \pi_*(Q_1, Q_2, \dots, Q_n) \longrightarrow \pi_*(Q_1, Q_2, \dots, Q_{n+1}), \quad n \geq 0$$

where, $n = 0$, $\pi_*(Q_1, Q_2, \dots, Q_n)$ is interpreted as π_* . (The f_n are not necessarily monomorphisms; see § 1. III).

From the construction of localization it is not difficult to see that we have the following lemma.

LEMMA 3.7. *The sequence $\{\pi_*(Q_1, \dots, Q_n), f_n | n = 0, 1, 2, \dots\}$ is a direct system of π_* -modules, with direct limit*

$$(2) \quad A_p = \lim \pi_*(Q_1, \dots, Q_n).$$

Next we will construct geometric objects realizing these π_* -modules, namely,

LEMMA 3.8. *There exist a sequence of spectra $\{\mathcal{L}_n\}$ and a sequence of maps $\{g_n: \mathcal{L}_n \rightarrow \mathcal{L}_{n+1}\}$ such that, for $n \geq 0$*

$$\pi_*(\mathcal{L}_n) = \pi_*(Q_1, Q_2, \dots, Q_n);$$

and

$$(g_n)_* = f_n: \pi_*(Q_1, Q_2, \dots, Q_n) \longrightarrow \pi_*(Q_1, \dots, Q_{n+1})$$

where f_n is the map in Lemma 3.7.

Proof. Take $\mathcal{L}^0 = S^0$ be the 0-sphere spectrum. Assume inductively that the spectrum \mathcal{L}_n is constructed such that

$$\pi_*(\mathcal{L}_n) = \pi_*(Q_1, \dots, Q_n).$$

Then we are going to construct \mathcal{L}_{n+1} such that

$$\pi_*(\mathcal{L}_{n+1}) = \pi_*(Q_1, \dots, Q_{n+1}).$$

Namely, consider the following directed system

$$(3) \quad \mathcal{L}_n \xrightarrow{q'_n} \mathcal{L}_n \xrightarrow{q'_n} \dots \xrightarrow{q'_n} \mathcal{L}_n \longrightarrow \dots$$

where q'_n is the identity map multiplied by the prime q_n (which is the n th prime listed in the beginning of this section § 3. II). Then, by [6, pp. 22, 4.5], the directed system (3) gives rise to a spectrum \mathcal{L}_{n+1} and a filtration $\{Y_i\}$ of \mathcal{L}_{n+1} such that, for each i , there is a homotopy equivalence

$$Y_i \simeq \mathcal{L}_n$$

and such that there exists a homotopy commutative diagram

$$(4) \quad \begin{array}{ccccccc} \mathcal{L}_n & \xrightarrow{q'_n} & \mathcal{L}_n & \xrightarrow{q'_n} & \cdots & \xrightarrow{q'_n} & \mathcal{L}_n \cdots \\ \downarrow \sim & & \downarrow \sim & & & & \downarrow \sim \\ Y_1 & \subset & Y_2 & \subset & \cdots & \subset & Y_i \cdots \end{array}$$

Thus, since the direct limit behaves well with respect to the stable homotopy group [6, pp. 21, 4.5], we have

$$(5) \quad \pi_*(\mathcal{L}_{n+1}) = \pi_* \lim_{\rightarrow} Y_i = \lim_{\rightarrow} \pi_*(Y_i) = \lim_{\rightarrow} \pi_*(\mathcal{L}_n)$$

where the first direct limit “lim” is taken over the bottom line of the above diagram (4), and the last “lim” is taken over (3). Note that we are assuming inductively that $\pi_* \overrightarrow{(\mathcal{L}_n)} = \pi_*(Q_1, \dots, Q_n)$. Hence (5) and Lemma 3.6, (1) imply that

$$\pi_*(\mathcal{L}_{n+1}) = \pi_*(Q_1, \dots, Q_{n+1}).$$

Thus we have constructed \mathcal{L}_{n+1} .

Note that there is a natural projection

$$g_n: \mathcal{L}_n = Y_1 \longrightarrow \mathcal{L}_{n+1} = \lim_{\rightarrow} Y_i$$

which obviously induces a map

$$\pi_*(\mathcal{L}_n) = \pi_*(Y_1) \longrightarrow \pi_*(\mathcal{L}_{n+1}) = \lim_{\rightarrow} \pi_*(\mathcal{L}_n)$$

that is just the map f_n . Thus we complete the proof of this lemma.

3.9. The proof of the Theorem 3.5.

Consider the directed system

$$(6) \quad \mathcal{L}_1 \xrightarrow{g_0} \mathcal{L}_1 \xrightarrow{g_1} \cdots \longrightarrow \mathcal{L}_n \xrightarrow{g_n} \cdots$$

which is constructed in Lemma 3.8. Then again by [6, pp. 22, 4.5], this directed system gives rises to a spectrum L_p such that

$$(7) \quad \pi_*(L_p) = \lim_{\rightarrow} \pi_*(\mathcal{L}_n)$$

where \lim_{\rightarrow} is taken over (6). Then, by Lemma 3.8,

$$\pi_*(L_p) = \lim_{\rightarrow} \pi_*(Q_1, \dots, Q_n),$$

and by Lemma 3.7

$$\pi_*(L_p) = A_p.$$

Thus we have constructed a spectrum L_p such that $\pi_*(L_p) = A_p$, and this completes the proof.

4. Stable homotopy ring. This section is the main body of the whole paper. The main theorems will be proved in first part, modulo a key computation. The key computation is in Theorem 2; we will prove this theorem in the second part.

I. First let us prove

THEOREM 4.1. *Let A be a locally finitely generated π_* -module having finite projective, as well as weak, dimension. Then A is a free π_* -module and hence is realizable as a stable homotopy module $\pi_*(Y)$ of a wedge Y of spheres.*

REMARK. The condition that A is locally finitely generated is necessary, see Theorem 4.4 below.

Proof. This theorem essentially depends on the next theorem, so we will give a proof modulo Theorem 2:

Let A be a locally finitely generated π_* -module. Then we will show that $P.d_{\pi_*}A < \infty$ implies that A is a free π_* -module. From Corollary 2.5, we see that it is sufficient to prove this for the case $P.d_{\pi_*}A \leq 1$. Let

$$0 \longleftarrow A \longleftarrow F_0 \xleftarrow{d_1} F_1 \longleftarrow 0$$

be a locally finitely generated free resolution (by Theorem 2.15 (ii)). Then, by Theorem 2.18, we can realize F_0, F_1 and d_1 by wedges of spheres N_0, N_1 and a map $f: N_0 \rightarrow N_1$; that is, $\pi_*(N_0) \cong F_0$, $\pi_*(N_1) \cong F_1$ and $d_1 = f_*$. Let C_f be the mapping cone of f , then C_f has a finite skeleton for each dimension (since N_0 and N_1 were chosen to have only finitely many spheres at each dimension). Since the map $d_1 = f_*$ is a monomorphism, the Puppe sequence (of the mapping cone sequence $N_0 \rightarrow N_1 \rightarrow C_f$) reduces to a short exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & \pi_*(C_f) & \longrightarrow & \pi_*(N_1) & \xrightarrow{f_*} & \pi_*(N_1) \longrightarrow 0 \\ & & & & \wr & & \wr \\ & & & & F_1 & & F_0 \end{array} \quad .$$

Comparing this with the other sequence above (i.e. the free resolution of A), we conclude that $\pi_*(C_f) \cong A$; hence $P.d_{\pi_*}(\pi_*(C_f)) = P.d_{\pi_*}A \leq 1$. On the other hand, from the next theorem (See Theorem 2 below),

$P.d_{\pi_*}A = P.d_{\pi_*}(\pi_*(C_f)) = 0$ or ∞ . Therefore we have $P.d_{\pi_*}(\pi_*C_f) = P.d_{\pi_*}A = 0$; hence, by Corollary 2.14 (1), A is free; and by Theorem 2.18 it is realizable by a wedge of spheres. This proves the theorem.

The proof of the next theorem is very long so we will only state the theorem here; the proof will be given later.

THEOREM 4.2. *Let Y be a connected spectrum having a finite skeleton at each dimension. Then the projective, as well as weak, dimension of $\pi_*(Y)$ as π_* -module is infinite unless Y is a wedge of spheres.*

THEOREM 4.3. (i) $f.P.D(\pi_*) = f\angle.P.D(\pi_*) = 0 < F.P.D(\pi_*) \neq 0$
(ii) $f.W.D(\pi_*) = f\angle.W.D(\pi_*) = 0 \leq F.W.D(\pi_*)$.

Proof. $f.P.D(\pi_*) = f\angle.P.D(\pi_*) = f.W.D(\pi_*) = f\angle.W.D(\pi_*) = 0$ follows from Theorem 4.1. $F.W.D(\pi_*) \geq 0$ follows from the definition. $F.P.D(\pi_*) > 0$ follows from Lemma 3.4.

THEOREM 4.4. *There is a stable homotopy module $\pi_*(X)$ such that $P.d_{\pi_*}(\pi_*(X)) = 1$.*

Proof. By Theorem 4.3, $F.P.D(\pi_*) > 0$. It then follows from Corollary 2.5 that there is a π_* -module, say A , such that $P.D_{\pi_*}(A) = 1$. Let

$$0 \longleftarrow A \longleftarrow F_0 \longleftarrow F_1 \longleftarrow 0$$

be a free resolution of A , which exists, by Theorem 2.15, (ii). Then, by Theorem 2.18, we can realize F_0 , F_1 and d_1 by wedges of spheres N_0 , N_1 and a map $f: N_1 \rightarrow N_0$, i.e., $\pi_*(N_0) \cong F_0$, $\pi_*(N_1) \cong F_1$ and $f_* = d_1$. Let $L_1 \xrightarrow{f} N_0 \rightarrow C_f$ be the mapping cone sequence of f , then the Puppe sequence reduces to a short exact sequence

$$0 \longleftarrow \pi_*(C_f) \longleftarrow \pi_*(N_0) \xleftarrow{f_*} \pi_*(N_1) \longleftarrow 0,$$

since $f_* = d_1$ is a monomorphism. By comparing the two short exact sequences, we see that $\pi_*(C_f) \cong A$. Setting $X = C_f$ finishes the proof.

REMARK. From [14], we can show that the spectrum L_p , which is constructed in Theorem 3.5, is such a spectrum.

From Theorem 4.3 and [3, Corollary 5.6], we have

THEOREM 4.5. (1) *Every finitely generated proper ideal of π_* has nonzero annihilator*

(ii) *A finitely generated projective submodule of a projective module is always a direct summand.*

THEOREM 4.6. *The projective as well as weak, dimension of an ideal in π_* is infinite.*

Proof. Let $I \subset \pi_*$ be a proper ideal. Let us assume that $W.d_{\pi_*} I \leq P.d_{\pi_*} I < \infty$. Then, by Theorem 4.3, $P.D_{\pi_*} I = 0$; and by Corollary 2.14, I is free. Thus the annihilator of I has to be zero and this contradicts Theorem 4.5. Therefore $P.d_{\pi_*} I = \infty$.

THEOREM 4.7. *Let ${}_p\pi_*$ be the subring of π_* consisting of π_0 and p -primary component of π_n , $n \geq 1$. Then the global dimension of ${}_p\pi_*$ is infinite.*

Proof. From the “slide product” (e.g. see [1] or [12]) representation of functor Tor , it is easy to see that the functors $\text{Tor}^{\pi_*}(-, Z_p)$ and $\text{Tor}^{p\pi_*}(-, Z_p)$ are equal. Since Z_p , the integers mod p , has weak dimension equal to ∞ , so there are infinitely many n such that, $\text{Tor}_{n,*}^{p\pi_*}(-, Z_p) \neq 0$. Hence $G.D({}_p\pi_*) = \infty$.

II. Next we will devote ourselves to the proof of Theorem 2. First let us recall a Kunneth type theorem for stable homotopy groups as a generalized theory. The details are in [1] or [12]. Let $W \wedge Y$ be the smash product and let $\pi_*(X, W)$ be the graded abelian group of the stable homotopy classes of maps from X to W . Let $h_\alpha: X^\alpha \rightarrow Y$ be a family of maps, and let $\bigvee_\alpha X^\alpha$ be the wedge of X^α 's. Then there is a map

$$h: X^\alpha \longrightarrow Y$$

such that its restriction on X^α is h_α , i.e., $h|X^\alpha = h_\alpha$. We will call this map h the wedge of h_α and denote it by $h = \bigvee_\alpha h_\alpha$.

THEOREM 4.8. *Let X, Y, W be spectra. Then there exist spectral sequence $\{E_{*,*}^r\}$ and $\{\tilde{E}_{*,*}^r\}$ with*

$$\begin{aligned} E_{*,*}^2 &= \text{Tor}^{\pi_*}(\pi_*(W), \pi_*(Y)) ; \\ \tilde{E}_{*,*}^2 &= \text{Tor}^{\pi_*}(\pi_*(W), \pi^*(X)) \end{aligned}$$

and if X is a finite spectrum and Y, Z are connected spectra, then both spectral sequence converge

$$\begin{aligned} E_{*,*}^r &\Longrightarrow \pi_*(W \wedge Y) ; \\ \tilde{E}_{*,*}^r &\Longrightarrow \pi_*(X, W) . \end{aligned}$$

Moreover the spectral sequences are functor of their arguments and the filtrations

$$\begin{aligned} F^o \subset F^1 \subset \dots \subset F^n \dots; \cup F^n &= \pi_*(W \wedge Y); \\ \tilde{F}^o \subset \tilde{F}^1 \subset \dots \subset \tilde{F}^n \dots; \cup \tilde{F}^n &= \pi_*(X, W), \end{aligned}$$

which satisfy

$$\begin{aligned} F^n / F^{n-1} &= E_{n,*}^\infty \\ \tilde{F}^n / \tilde{F}^{n-1} &= \tilde{E}_{n,*}^\infty, \end{aligned}$$

are given by

$$\begin{aligned} F^n &= \text{Ker}\{\pi_*(W \wedge Y) \longrightarrow \pi_*(W_n \wedge Y)\}; \\ \tilde{F}^n &= \text{Ker}\{\pi_*(X, W) \longrightarrow \pi_*(X, W_n)\}, \end{aligned}$$

where W_n is defined inductively as follows: Let $W_{-1} = W$, and assume that W_n is defined. Let $\{g_\alpha: S^{n_\alpha} \rightarrow W_n\}$ be the set of generators of $\pi_*(W_n)$ as a π_* -module. Let $M_{n+1} = \bigvee_\alpha S^{n_\alpha}$, and let $f_{n+1} = \bigvee_\alpha g_\alpha: M_{n+1} \rightarrow W_n$ be the wedge of maps g_α . Then W_{n+1} is defined to be the mapping cone of f_{n+1} .

REMARK. Let $\pi_*(X) \otimes \pi_*(Y)$ be the tensor product over π_* .¹ Then the edge homomorphism of the spectral sequence $\{E_{*,*}^r\}$ in the above theorem is a map

$$\pi_*(X) \otimes \pi_*(Y) \longrightarrow \pi_*(X \wedge Y)$$

which sends $f \otimes g$ into $f \wedge g$ for $f \in \pi_*(X)$ and $g \in \pi_*(Y)$. We will call this map smash product (This map is stronger than the usual “smash product”, because we require the tensor product be over π_*).

COROLLARY 4.9. *If $\pi_*(W)$ is a flat π_* -module, then the smash product*

$$\pi_*(W) \otimes \pi_*(Y) \longrightarrow \pi_*(W \wedge Y),$$

is an isomorphism.

COROLLARY 4.10. *Let $\pi_*(W)$ be a flat π_* -module. Then all the primary and higher order homology operations of degree ≥ 1 are all zero on $H_*(W)$.*

Proof. The notations are the same as Theorem 4.8. Since $\pi_*(W)$ is flat, the spectral sequence $\{E_{*,*}^r\}$ in the Theorem 4.8 has

¹ Throughout the whole paper \otimes will always mean tensor product over the ring π_* .

$$E_{n,*}^2 = \text{Tor}_{n,*}^{\pi_*}(\pi_*(W), \pi_*(Y)) = 0 \quad n \neq 0.$$

Hence

$$E_{n,*}^\infty = 0, \quad \text{for } n \neq 0.$$

In other words

$$F^0 = F^1 = F^2 = \cdots = \bigcup_i F^i = \pi_*(W \wedge Y)$$

or equivalently,

$$F^0 = \pi_*(W \wedge Y) = \text{Ker}\{\pi_*(W \wedge Y) \longrightarrow \pi_*(W_0 \wedge Y)\}.$$

That is,

$$\pi_*(W \wedge Y) \longrightarrow \pi_*(W_0 \wedge Y) \quad \text{is a zero map.}$$

Then by the Puppe sequence of the mapping cone sequence $M_0 \rightarrow W \rightarrow W_0$ (constructed in the statement of Theorem 4.8), we have an epimorphism

$$\pi_*(M_0 \wedge Y) \longrightarrow \pi_*(W \wedge Y) \longrightarrow 0.$$

Take Y to be the Eilenberg-MacLane spectrum $K(G)$, then the above epimorphism reduces to the epimorphism

$$H_*(M_0; G) \longrightarrow H_*(W; G) \longrightarrow 0$$

where G is any coefficient group. Since M_0 is a wedge of spheres (see Theorem 4.8), all the homology operations of degree ≥ 1 are zero on $H_*(M_0)$. Thus, by naturality of homology operations, all the homology operations of degree ≥ 1 on $H_*(W)$ are zero.

Next we will look at the Spanier-Whithead duality and its relation with cohomology (homology) operations. Let X be a finite spectrum (roughly speaking, that is a spectrum with finite number of stable cells; see [6, pp. 7, C. 7] for precise definition). Then, from [6, pp. 33, K. 12] we can choose its Spanier-Whithead dual DX to be finite too. Then it is obvious to observe that.

LEMMA 4.11. *Let X be a finite spectrum. Then X is a wedge of spheres if and only if DX is.*

Let us recall from [19] the cohomology operations. Let X be an arbitrary spectrum, and let α be one of the lowest order nonzero cohomology operations on $H^*(X)$. Then α can be represented as a differential of the spectral sequence obtained from some Postnikov system. More precisely, let

$$\begin{array}{ccccccc}
E_0 & \longleftarrow & E_1 & \longleftarrow & \cdots & \longleftarrow & E_n & \longleftarrow & \cdots \\
\downarrow & & \downarrow & & & & \downarrow & & \\
K_0 & & K_1 & & & & K_n & &
\end{array}$$

be a Postnikov system, where K_i are Eilenberg-MacLane objects. By applying the functor $\pi_*(X, -)$ to the system we get a spectral sequence with $E_{*,*}^1 = \pi_*(X, K_*) = H^*(X; K_*)$. Then since α is the lowest order nonzero cohomology operation (and hence has zero indeterminacy), the differential α representing this operation is a non-zero group-homomorphism

$$\alpha: H^q(X; K_{n_1}) \longrightarrow H^{q+r}(X; K_{n_2}).$$

Next observe that $H^*(X, K_n) = \pi_*(DX \wedge K_n) = H_*(DX; K_n)$, by Spanier-Whithead duality (see [19]). Then the spectral sequence can be written as

$$E_{*,*}^1 = H^*(X; K_*) = H_*(DX; K_*)$$

and hence α is a nonzero homology operation on $H_*(DX)$. Thus we have proved the following proposition.

PROPOSITION 4.12. *Let X be a finite spectrum. Then there is a nonzero cohomology operation of degree ≥ 1 on $H^*(X)$ if and only if there is a nonzero homology operation of degree ≥ 1 on $H_*(DX)$.*

We need an algebraic Lemma on localization.

LEMMA 4.13. *Let $\delta: G_1 \rightarrow G_2$ be a nonzero homomorphism of abelian groups from G_1 to G_2 . Then there is a prime integer p such that*

$$\delta \otimes_{\mathbb{Z}} \mathbb{Z}(p): G_1 \otimes_{\mathbb{Z}} \mathbb{Z}(p) \rightarrow G_2 \otimes_{\mathbb{Z}} \mathbb{Z}(p)$$

is a nonzero homomorphism, where $\mathbb{Z}(p)$ is the local ring obtained by localizing the ring \mathbb{Z} of integers at the prime ideal $p\mathbb{Z}$, and $\otimes_{\mathbb{Z}}$ is the tensor product of abelian groups.

Proof. It is sufficient to show that there is a prime p such that $(\text{Im } \delta) \otimes_{\mathbb{Z}} \mathbb{Z}(p) \neq 0$. Let $C \subset \text{Im } \delta$ be a cyclic subgroup of $\text{Im } \delta$, then $C \otimes_{\mathbb{Z}} \mathbb{Z}(p) \subset \text{Im } \delta \otimes_{\mathbb{Z}} \mathbb{Z}(p)$. (Note that $\mathbb{Z}(p)$ is a torsion free group; hence the inclusion is preserved). Thus it is sufficient to show that for any cyclic group C , there is some $\mathbb{Z}(p)$ such that $C \otimes_{\mathbb{Z}} \mathbb{Z}(p) \neq 0$; and this is obvious.

PROPOSITION 4.14. *Let all the homology operations of degree ≥ 1 be zero on $H_*(Y)$. Then the integral homology group $H_*(Y; \mathbb{Z})$ is a*

torsion free abelian group.

Proof. Suppose $H_m = H_m(Y; Z)$ be the lowest dimension group which has torsion. Let G be a group such that $\text{Tor}^2(H_m, G) \neq 0$. Let

$$(1) \quad 0 \longrightarrow P_1 \xrightarrow{d_1} P_0 \longrightarrow G \longrightarrow 0$$

be a Z -free resolution of G . Then we have the exact sequence of Tor :

$$\begin{aligned} 0 = \text{Tor}^Z(H_m, P_0) &\longrightarrow \text{Tor}^Z(H_m, G) \longrightarrow H_m \otimes_Z P_1 \xrightarrow{H_m \otimes_Z d_1} H_m \otimes_Z P_0 \\ &\longrightarrow H_m \otimes_Z G \longrightarrow 0. \end{aligned}$$

Since $\text{Tor}^Z(H_m, G) \neq 0$, so $H_m \otimes_Z d_1$ can not be monomorphic. On the other hand, the long exact sequence of homology groups associated to (1) is

$$\cdots \longrightarrow H_{m+1}(Y, G) \xrightarrow{\beta} H_m(Y, P_1) \xrightarrow{d'_1} H_m(Y, P_2) \longrightarrow \cdots$$

Since P_i are free, so $H_m(Y, P_i) = H_m \otimes_Z P_i$ and $d'_1 = H_m \otimes_Z d_1$. Therefore $d'_1 = H_m \otimes_Z d_1$ is not a monomorphism. Thus, by exactness, the Bockstein operator β is not zero. This proves the Proposition.

PROPOSITION 4.15. *Let L_p be the spectrum constructed in Theorem 3.5. Then all the primary and higher order homology operations of degree ≥ 1 are zero on $H_*(Y)$ if all the primary and higher order homology operations of degree ≥ 1 are zero on $H_*(Y \wedge L_p)$ for every prime p .*

Proof. Let us assume, to the contrary, that there are nonzero homology operations of degree ≥ 1 on $H_*(Y)$. Let α be one of the lowest order of nonzero homology operations of degree ≥ 1 . Then α can be expressed as a differential of the spectral sequence of the following exact couple

$$\begin{array}{ccc} \sum_n \pi_*(Y \wedge E_n) & \longrightarrow & \sum_n \pi_*(Y \wedge E_{n-1}) \\ & \swarrow \quad \searrow & \\ & \sum_n \pi_*(Y \wedge K_n) = \sum_n H_*(Y; K_n) & \end{array}$$

where $\{E_n\}$ is a Postnikov system and for each n , K_n is an Eilenberg-MacLane spectrum, Since α is the lowest order nonzero operation (and hence the indeterminacy is zero), so the differential, which represents this operation α , gives rise to a nonzero group homomorphism

α' on E_{q_1, n_1}^1 - component, i.e.,

$$(2) \quad \alpha': H_{q_1}(Y; K_{n_1}) = E_{q_1, n_1}^1 \longrightarrow H_{q_2}(Y; K_{n_2}) = E_{q_2, n_2}^1$$

is a nonzero homomorphism of groups. Observe that $\pi_*(L_p) = A_p$ is a flat π_* -module, therefore

$$(3) \quad \begin{array}{ccc} \sum \pi_*(Y \wedge E_n) \otimes A_p & \longrightarrow & \sum \pi_*(Y \wedge E_n) \otimes A_p \\ & \nwarrow \quad \nearrow & \\ \sum \pi_*(Y \wedge K_n) \otimes A_p & = & \sum H_*(Y; K_n) \otimes A_p \end{array}$$

is still an exact couple.

Moreover, from properties of tensor product, it is readily seen that

$$H_*(Y; K_n) \otimes A_p = H_*(Y; K_n) \otimes_Z Z(p), \quad \text{as groups,}$$

where $Z(p)$ is the ring Z localized at pZ . Thus the differential $\alpha \otimes A_p$ of (3), when it is restricted to E_{q_1, n_1}^1 , is just a homomorphism of groups

$$\alpha' \otimes_Z Z(p): H_{q_1}(Y; K_{n_1}) \otimes_Z Z(p) \longrightarrow H_{q_2}(Y; K_{n_2}) \otimes_Z Z(p).$$

Then, by Lemma 4.13, we can choose a prime p such that

$$\alpha' \otimes_Z Z(p) \neq 0.$$

In other words, there is some prime p such that the differential

$$(4) \quad \alpha \otimes A_p \neq 0$$

is not a zero differential in (3), since $\alpha \otimes A_p|_{E_{q_1, n_1}^1} = \alpha' \otimes_Z Z(p)$.

Next, note that the exact couple (3), by Corollary 4.9, can be written as

$$(5) \quad \begin{array}{ccc} \sum \pi_*(Y \wedge E_n \wedge L_p) & \longrightarrow & \sum \pi_*(Y \wedge E_n \wedge L_p) \\ & \nwarrow \quad \nearrow & \\ \sum \pi_*(Y \wedge K_n \wedge L_p) & = & \sum H_*(Y \wedge L_p; K_n). \end{array}$$

Therefore the differential $\tilde{\alpha}$ of (5), which corresponds to the differential $\alpha \otimes A_p$ of (3), is not a zero differential, i.e., by (4) we have,

$$0 \neq \tilde{\alpha} = \alpha \otimes A_p: H_{q_1}(Y \wedge L_p; K_{n_1}) \longrightarrow H_{q_2}(Y \wedge L_p; K_{n_2}).$$

Moreover $\tilde{\alpha}$ can be regarded as a homology operation (of degree ≥ 1) on $H_*(Y \wedge L_p)$; thus $\tilde{\alpha} \neq 0$ contradicts the assumption of the proposition. This completes the proof.

We also need a well known folk theorem about F. Peterson's work on detecting maps by cohomology operations (e.g. see [13]).

THEOREM 4.16. *A connected spectrum is a wedge of spheres if and only if all the primary and higher order cohomology operations of degree ≥ 1 are zero.*

Now we are ready for the proof of Theorem 2.

4.17. *The proof of Theorem 2.*

From Example 2.8 and Proposition 2.9, we see that it is sufficient to show only for the projective dimension case. Let us assume that $P.d_{\pi_*}(\pi_*(Y)) < \infty$. Then from standard homological algebra (e.g. see [17]) we have

$$P.d_{A_p}(\pi_*(Y) \otimes A_p) < \infty .$$

By Theorem 3.2, $f/P.D(A_p) = 0$. It then follows that

$$P.d_{A_p}(\pi_*(Y) \otimes A_p) = 0 .$$

Thus, by Corollary 2.14, $\pi_*(Y) \otimes A_p$ is a free A_p -module. Moreover, by Corollary 4.9, we have

$$\pi_*(Y \wedge L_p) = \pi_*(Y) \otimes A_p ,$$

where L_p is the spectrum constructed in Theorem 3.5, and $\pi_*(L_p) = A_p$. Thus $\pi_*(Y \wedge L_p)$, a free A_p -module, is a flat π_* -module. Then, by Corollary 4.10, all the homology operations of degree ≥ 1 are all zero on $H_*(Y \wedge L_p)$; and this is true for every prime p . By Proposition 4.15, all the homology operations of degree ≥ 1 are zero on $H_*(Y)$.

Since Y is connected, there is a homology decomposition, or co-Postnikov system $\{Y_n\}$ (e.g. see [8])

$$(6) \quad \begin{array}{ccccc} Y_m & \subset & Y_{m+1} & \subset & \cdots \subset Y_n & \subset & \cdots; & \bigcup_{n \geq m} Y_n = W \simeq Y \\ \uparrow k'_m & & \uparrow k'_{m+1} & & & & \uparrow k'_n & \\ N_m & & N_{m+1} & & & & N_n & \end{array}$$

such that

- (i) $H_q(Y; Z) \cong H_q(Y_n; Z) \quad q \leq n$
- (ii) $H_q(Y_n; Y) = 0 \quad q > n$
- (iii) N_n is the Moore space of type $(H_{n+1}(Y), n)$
- (iv) $Y_{n+1} = Y_n \cup k'_n CN_n$, the mapping cone of k'_n .
- (v) $k'_*: H_*(N_n) \longrightarrow H_*(Y_n)$ is 0.

Note that, in the last paragraph, we have shown that all the homology operations of degree ≥ 1 are zero on $H_*(Y)$. Thus, by Proposition 4.14, $H_*(Y; Z)$ is torsion free. By the universal coefficient theorem,

$$H_*(Y; G) = H_*(Y; Z) \otimes_Z G.$$

Thus (i) and (ii) of (6) can be improved to

$$\begin{aligned} \text{(i')} \quad H_q(Y; G) &\cong H_p(Y_n; G) & q \leq n \\ \text{(ii')} \quad H_q(Y; G) &= 0 & q > n. \end{aligned}$$

Since all the homology operations (of degree ≥ 1) are all zero on $H_*(Y)$, by (i') and (ii'), we see immediately that all the homology operations of degree ≥ 1 are also zero on $H_*(Y_n)$ for every n . Thus, by Proposition 4.12, all the cohomology operations of degree ≥ 1 are all zero on $H^*(DY_n)$ for every n , where DY is Spanier-Whithead dual of Y . By Theorem 4.16, DY_n is a wedge spheres for each n . Recall that, by assumption, Y has finite skeleton at each dimension. Therefore we can choose each Y_n in (6) to be finite. Thus, by Lemma 4.11 and the fact $DDY_n \simeq Y_n$ [6, pp. 33, K. 12], we can conclude that each Y_n is a wedge of sphere (since we have just shown above that DY_n is a wedge of spheres). Thus, by Peterson's theorem on detecting map, we conclude that k'_{n-1} is null-homotopic; and this is true for every n . Hence Y is a wedge of spheres.

REFERENCES

1. J. F. Adams, *Lectures on generalized cohomology*, Lectures notes in Math. No. 99, Category theory, homology theory and their application III, pp. 1-138, Spring-Verlag 1968.
2. Adams, *Stable homotopy theory*, Lecture Notes in Math. No. 3 Spring-Verlag 1964.
3. H. Bass, *Finitistic dimension and homological generalization of semi-primary ring*, Trans. Amer. Math. Soc., **95** (1960), 466-488.
4. ———, *Big projective modules are free*. Illinois J. Math. (1963), 24-31.
5. ———. *Algebraic K-Theory*, Benjamin 1968.
6. J. Boardman, *Stable homotopy theory*, Mimeograph notes, Univ. of Warwick, 1965.
7. P. Freyd, *Stable homotopy*, Proc. of LaJolla Conf. pp. 121-172, Spring-Verlag 1965.
8. P. Hilton, *Homotopy theory and duality*, Gordon and Breach, 1965, New York 1964.
9. J. P. Jans, *Rings and homology*, Holt, Rinehart, and Winston, New York 1964.
10. D. W. Kahn, *On stable homotopy modules I*, Invent. Math. I, (1966), 375-379.
11. M. Kervaire and J. Milnor, *Bernoulli numbers, homotopy groups and a theorem of Rohlin*, Proc. Int. Congress, Edinburgh, 1958.
12. T. Y. Lin, *A spectral sequence in stable homotopy category*, Thesis, Yale Univ. 1968.
13. ———, *Cohomological characterization of stable spheres* (unpublished).
14. ———, *Homological dimensions of π_* -modules and their geometric characterization* (in preparation).
15. P. May, *The Algebraic Eilenberg-Moore spectral Sequence* (to appear).
16. ———, *Homological algebra*, lectures, Yale Univ. 1965.
17. D. G. Northcott. *An introduction to homological algebra*. Cambridge Univ. Press, New York. 1960.
18. O. Ore, *Linear equation in non-commutative fields*, Ann. of Math. **32** (1931), 463-477.
19. F. Peterson, *Functional cohomology operations*, Trans. Amer. Math. Soc., **86** (1957), 197-211.
20. Toda, *Composition method in homotopy group of spheres*, Ann. of Math, Studies, No. 49, Princeton, 1962.

Received June 16, 1970 and in revised form November 9, 1970. The preparation of this paper was sponsored partly by the National Science Foundation Grant GP-21062. The abstract was announced in the Notices of AMS, February 1970, #70-G37.

PURDUE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

C. R. HOBBY
University of Washington
Seattle, Washington 98105

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 38, No. 1

March, 1971

Bruce Alan Barnes, <i>Banach algebras which are ideals in a Banach algebra</i>	1
David W. Boyd, <i>Inequalities for positive integral operators</i>	9
Lawrence Gerald Brown, <i>Note on the open mapping theorem</i>	25
Stephen Daniel Comer, <i>Representations by algebras of sections over Boolean spaces</i>	29
John R. Edwards and Stanley G. Wayment, <i>On the nonequivalence of conservative Hausdorff methods and Hausdorff moment sequences</i>	39
P. D. T. A. Elliott, <i>On the limiting distribution of additive functions (mod 1)</i>	49
Mary Rodriguez Embry, <i>Classifying special operators by means of subsets associated with the numerical range</i>	61
Darald Joe Hartfiel, <i>Counterexamples to a conjecture of G. N. de Oliveira</i>	67
C. Ward Henson, <i>A family of countable homogeneous graphs</i>	69
Satoru Igari and Shigehiko Kuratsubo, <i>A sufficient condition for L^p-multipliers</i>	85
William A. Kirk, <i>Fixed point theorems for nonlinear nonexpansive and generalized contraction mappings</i>	89
Erwin Kleinfeld, <i>A generalization of commutative and associative rings</i>	95
D. B. Lahiri, <i>Some restricted partition functions. Congruences modulo 11</i>	103
T. Y. Lin, <i>Homological algebra of stable homotopy ring π_* of spheres</i>	117
Morris Marden, <i>A representation for the logarithmic derivative of a meromorphic function</i>	145
John Charles Nichols and James C. Smith, <i>Examples concerning sum properties for metric-dependent dimension functions</i>	151
Asit Baran Raha, <i>On completely Hausdorff-completion of a completely Hausdorff space</i>	161
M. Rajagopalan and Bertram Manuel Schreiber, <i>Ergodic automorphisms and affine transformations of locally compact groups</i>	167
N. V. Rao and Ashoke Kumar Roy, <i>Linear isometries of some function spaces</i>	177
William Francis Reynolds, <i>Blocks and F-class algebras of finite groups</i>	193
Richard Rochberg, <i>Which linear maps of the disk algebra are multiplicative</i>	207
Gary Sampson, <i>Sharp estimates of convolution transforms in terms of decreasing functions</i>	213
Stephen Scheinberg, <i>Fatou's lemma in normed linear spaces</i>	233
Ken Shaw, <i>Whittaker constants for entire functions of several complex variables</i>	239
James DeWitt Stein, <i>Two uniform boundedness theorems</i>	251
Li Pi Su, <i>Homomorphisms of near-rings of continuous functions</i>	261
Stephen Willard, <i>Functionally compact spaces, C-compact spaces and mappings of minimal Hausdorff spaces</i>	267
James Patrick Williams, <i>On the range of a derivation</i>	273