

Pacific Journal of Mathematics

TWO UNIFORM BOUNDEDNESS THEOREMS

JAMES DEWITT STEIN

TWO UNIFORM BOUNDEDNESS THEOREMS

JAMES D. STEIN JR.

A geodesically convex space is a metric space in which each two points can be connected by a unique segment (a path of minimal length). An affine transformation between two geodesically convex spaces is a map which takes segments into segments. It is shown that, if the domain is complete, a pointwise-bounded family of continuous affine transformations is uniformly bounded. Under a mild additional hypothesis, the following stronger theorem holds: if

$$\mathcal{F} = \{T_\alpha \mid \alpha \in A\}$$

is a pointwise-bounded family of affine transformations and T_α is continuous on a closed geodesically convex S_α with

$$\bigcap_{\alpha \in A} S_\alpha \neq \emptyset,$$

then $\exists \alpha_1, \dots, \alpha_n$ such that \mathcal{F} is uniformly bounded on

$$\bigcap_{k=1}^n S_{\alpha_k}.$$

Let $(X, d), (Y, d')$ be metric spaces, and $\mathcal{F} = \{T_\alpha \mid \alpha \in A\}$ a collection of maps from X to Y . We say \mathcal{F} is pointwise-bounded if, for fixed $x, y \in X$, $\sup \{d'(T_\alpha x, T_\alpha y) \mid \alpha \in A\}$ is finite. If $x_0 \in S \subseteq X$, we say \mathcal{F} is uniformly bounded on S if $\sup \{d'(T_\alpha x, T_\alpha x_0) \mid x \in S, \alpha \in A\}$ is finite. A uniform boundedness theorem is one in which uniform boundedness (for some family \mathcal{F}) is deduced from pointwise-boundedness.

Let $\gamma: [0, 1] \rightarrow X$ be continuous, $0 = t_0 < \dots < t_n = 1$ a partition P of $[0, 1]$, define $\ell(\gamma, P) = \sum_{k=1}^n d(\gamma(t_k), \gamma(t_{k-1}))$, and define $\ell(\gamma)$ to be the supremum over all partitions P of the $\ell(\gamma, P)$. For $x, y \in X$, define $d_g(x, y) = \inf \{\ell(\gamma) \mid \gamma: [0, 1] \rightarrow X, \gamma(0) = x, \gamma(1) = y\}$; this is the geodesic or intrinsic distance between x and y . d_g is a generalized metric, and γ is said to be a segment from x to y if

$$\gamma(0) = x, \gamma(1) = y,$$

and $\ell(\gamma) = d_g(x, y) < \infty$.

DEFINITION 1. X is said to be geodesically convex if for any x, y in X there is a unique segment from x to y . We denote by $\Phi_g(x, y, t)$ the intrinsic parametrization of this segment (if $0 \leq t \leq s \leq 1$, $d_g(\Phi_g(x, y, t), \Phi_g(x, y, s)) = (s - t)d_g(x, y)$); T is said to be an affine map between geodesically convex spaces if $T(\Phi_g(x, y, t)) = \Phi_g(Tx, Ty, t)$.

A term often used for a geodesically convex space is a space with

unique segments. Throughout this paper we assume $d = d_g$.

Our first theorem is a generalization to geodesically convex spaces of the classical Banach-Steinhaus Theorem.

THEOREM 1. *Let (X, d) be a geodesically convex complete metric space and let (Y, d') be geodesically convex. Let $\mathcal{F} = \{T_\alpha | \alpha \in A\}$ be a pointwise-bounded family of geodesically affine maps from X to Y , each of which is continuous. Then for each $x_0 \in X$,*

$$\sup \{d'(T_\alpha x, T_\alpha x_0) | \alpha \in A, d(x, x_0) \leq 1\}$$

is finite.

We shall need the following lemma.

LEMMA 1. *For each $\alpha \in A, z_0 \in X$ and $p > 0$,*

$$r(\alpha, z_0, p) = \sup \{d'(T_\alpha x, T_\alpha z_0) | d(x, z_0) \leq p\}$$

is finite.

Proof. By continuity of T_α at z_0 , $\exists \delta > 0$ such that

$$d(x, z_0) < \delta \implies d'(T_\alpha x, T_\alpha z_0) < 1;$$

we can clearly assume $\delta < p$. If $x \in X, d(x, z_0) \leq p$, let $z = \Phi_g(z_0, x, \delta/2p)$, then $d(z_0, z) = \delta/2pd(z_0, x) < \delta$, so $d'(T_\alpha z_0, T_\alpha z) < 1$. But

$$T_\alpha(\Phi_g(z_0, x, \delta/2p)) = \Phi_g(T_\alpha z_0, T_\alpha x, \delta/2p),$$

and so $d'(T_\alpha z_0, T_\alpha z) = \delta/2pd'(T_\alpha z_0, T_\alpha x) < 1$, so $d'(T_\alpha z_0, T_\alpha x) < 2p/\delta$.

For purposes of simplicity, we prove the following lemma.

LEMMA 2. *Assume the conclusion of the theorem is false. Let $M > 0, x_1, \dots, x_n \in X$ and $T_1, \dots, T_n \in \mathcal{F}$ be given, with $d(x_0, x_k) < 1 (1 \leq k \leq n)$. Then $\exists x_{n+1} \in X, T_{n+1} \in \mathcal{F}$ with $d(x_0, x_{n+1}) < 1, d(x_n, x_{n+1}) < 1/2^{n+1}, d'(T_{n+1}x_{n+1}, T_{n+1}x_0) > M$, and*

$$d'(T_k x_n, T_k x_{n+1}) < 1/2^{n+1}$$

for $1 \leq k \leq n$.

Proof. For $x \in X$, let $S(x) = \sup \{d'(T_\alpha x, T_\alpha x_0) | \alpha \in A\}$. Let

$$\alpha = 1/3 \min (2^{-n-1}r(1, x_n, 2)^{-1}, \dots, 2^{-n-1}r(n, x_n, 2)^{-1}, 2^{-n-1}, 1 - d(x_n, x_0)),$$

then $\alpha > 0$. If the theorem is false, then for any $K > 0$ there is a $z \in X$ with $d(x_0, z) < 1$ and a $T \in \mathcal{F}$ with $K < d'(Tx_0, Tz)$, consequently

$$K < d'(Tx_0, Tz) \leq d'(Tx_0, Tx_n) + d'(Tx_n, Tz) \leq S(x_n) + d'(Tx_n, Tz).$$

This means that we can always find a $z \in X$ and $T \in \mathcal{F}$ with

$$d(x_0, z) < 1$$

and $d'(Tx_n, Tz)$ arbitrarily large. Having defined α , choose $y \in X$, $T (= T_{n+1}) \in \mathcal{F}$ with $d(y, x_0) < 1$, $\alpha d'(Tx_n, Ty) - S(x_n) > M$, and let $x_{n+1} = \Phi_g(x_n, y, \alpha)$. Then $d(x_n, x_{n+1}) = \alpha d(x_n, y) \leq 2^{-n-1}$. For $1 \leq k \leq n$, we have

$$\begin{aligned} d'(T_k x_n, T_k x_{n+1}) &= d'(T_k \Phi_g(x_n, y, 0), T_k \Phi_g(x_n, y, \alpha)) \\ &= d'(\Phi_g(T_k x_n, T_k y, 0), \Phi_g(T_k x_n, T_k y, \alpha)) \\ &= \alpha d'(T_k x_n, T_k y) \leq \alpha r(k, x_n, 2) < 2^{-n-1}. \end{aligned}$$

We also have

$$\begin{aligned} d(x_0, x_{n+1}) &\leq d(x_0, x_n) + d(x_n, x_{n+1}) \\ &\leq d(x_0, x_n) + d(\Phi_g(x_n, y, 0), \Phi_g(x_n, y, \alpha)) \\ &= d(x_0, x_n) + \alpha d(x_n, y) < d(x_0, x_n) + 2\alpha \\ &< d(x_0, x_n) + 1 - d(x_0, x_n) \\ &= 1. \end{aligned}$$

Finally,

$$\begin{aligned} \alpha d'(Tx_n, Ty) &= d'(Tx_n, Tx_{n+1}) \\ &\leq d'(Tx_n, Tx_0) + d(Tx_0, Tx_{n+1}) \\ &\leq S(x_n) + d'(Tx_0, Tx_{n+1}) \Rightarrow d'(Tx_0, Tx_{n+1}) \\ &\geq \alpha d'(Tx_n, Ty) - S(x_n) > M, \end{aligned}$$

completing the proof.

We return to the proof of the theorem. Assume the theorem is false. Then $\exists x_1 \in X, T_1 \in \mathcal{F}$ with

$$d(x_0, x_1) < 1, d'(T_1 x_0, T_1 x_1) > 2.$$

Having chosen $x_1, \dots, x_n \in X, T_1, \dots, T_n \in \mathcal{F}$ with

$$d(x_0, x_k) < 1 (1 \leq k \leq n),$$

by Lemma 2 choose $x_{n+1} \in X, T_{n+1} \in \mathcal{F}$ with $d(x_0, x_{n+1}) < 1, d(x_n, x_{n+1}) < 2^{-n-1}, d'(T_{n+1} x_0, T_{n+1} x_{n+1}) > n + 2$ and $d'(T_k x_n, T_k x_{n+1}) < 2^{-n-1}$ for $1 \leq k \leq n$. Since $d(x_n, x_{n+1}) < 2^{-n-1}$, the sequence $\{x_n | n = 1, 2, \dots\}$ is Cauchy ($n < m \Rightarrow d(x_n, x_m) < \sum_{k=n}^{m-1} 2^{-k-1}$); by completeness $x_n \rightarrow x \in X$. By continuity of T_n we have $\lim_{m \rightarrow \infty} d'(T_n x, T_n x_{m+1}) = 0$, so

$$\begin{aligned}
 d'(T_n x_0, T_n x_n) &\leq d'(T_n x_0, T_n x) + d'(T_n x, T_n x_n) \leq \dots \\
 &\leq d'(T_n x_0, T_n x) + \sum_{k=n}^m d'(T_n x_k, T_n x_{k+1}) + d'(T_n x, T_n x_{m+1}) ;
 \end{aligned}$$

letting $m \rightarrow \infty$ we obtain

$$\begin{aligned}
 d'(T_n x_0, T_n x_n) &\leq d'(T_n x_0, T_n x) + \sum_{k=n}^{\infty} d'(T_n x_k, T_n x_{k+1}) \\
 &< d'(T_n x_0, T_n x) + \sum_{k=n}^{\infty} 2^{-k-1} < d'(T_n x_0, T_n x) + 1 ,
 \end{aligned}$$

since $k \geq n \Rightarrow d'(T_n x_k, T_n x_{k+1}) < 2^{-k-1}$. So

$$n + 1 < d'(T_n x_0, T_n x_n) \leq d'(T_n x_0, T_n x) + 1 \Rightarrow d'(T_n x_0, T_n x) > n ,$$

contradicting the pointwise-boundedness of \mathcal{F} .

We now make an additional hypothesis, which will enable us to prove a stronger version of this theorem. Let $\Phi = \Phi_g$.

DEFINITION 2. If $0 < \alpha < 1$, define

$$M(\alpha) = \sup \{d(\Phi(x, y, \alpha), \Phi(x, z, \alpha))/d(y, z) \mid x, y, z \in X, y \neq z\} ,$$

and define $M'(\alpha)$ similarly in Y . Note that, if $M(\alpha) < \infty$, then

$$x, y, z \in X \Rightarrow d(\Phi(x, y, \alpha), \Phi(x, z, \alpha)) \leq M(\alpha)d(y, z) .$$

For the remainder of this paper we shall make the following assumption: $\exists \alpha \in (0, 1)$ such that both $M(\alpha)$ and $M'(\alpha)$ are finite. This α will be fixed from now on.

DEFINITION 3. Let $\{x_n \mid n = 1, 2, \dots\} \subseteq X$, and let $x_0 \in X$. Define $z_1^{(n)} = \Phi(x_n, x_0, \alpha)$, and for $2 \leq k \leq n$ define $z_k^{(n)} = \Phi(x_{n+1-k}, z_{k-1}^{(n)}, \alpha)$. Now define $y_n = z_n^{(n)}$ for $n = 1, 2, \dots$.

If X were a Banach space and $x_0 = 0$, then we would have

$$y_n = \sum_{k=1}^n (1 - \alpha)^k x_k .$$

In general, however, we have $y_n = \Phi(x_1, \Phi(x_2, \dots, \Phi(x_n, x_0, \alpha), \dots, \alpha))$, which will henceforth be abbreviated $\Phi(x_1, \dots, \Phi(x_n, x_0, \alpha), \dots, \alpha)$.

LEMMA 3. *Given*

$$\{x_n \mid n = 1, 2, \dots\} \subseteq X ,$$

$x_0 \in X$, define $\{y_n \mid n = 1, 2, \dots\}$ as in Definition 3. Then

$$d(y_n, y_{n-1}) \leq M(\alpha)^{n-1}(1 - \alpha)d(x_n, x_0)$$

if $n \geq 2$.

Proof. Clearly, we have

$$\begin{aligned} & d(y_n, y_{n-1}) \\ &= d(\Phi(x_1, \dots, \Phi(x_n, x_0, \alpha), \dots, \alpha), \Phi(x_1, \dots, \Phi(x_{n-1}, x_0, \alpha), \dots, \alpha)) \\ &\leq M(\alpha)d(\Phi(x_2, \dots, \Phi(x_n, x_0, \alpha), \dots, \alpha), \Phi(x_2, \dots, \Phi(x_{n-1}, x_0, \alpha), \dots, \alpha)) \\ &\leq \dots \leq M(\alpha)^{n-2}d(\Phi(x_{n-1}, \Phi(x_n, x_0, \alpha), \alpha), \Phi(x_{n-1}, x_0, \alpha)) \\ &\leq M(\alpha)^{n-1}d(\Phi(x_n, x_0, \alpha), x_0) \\ &= (1 - \alpha)M(\alpha)^{n-1}d(x_n, x_0) . \end{aligned}$$

LEMMA 4. Let S be a convex subset of X , $p > 0$, and let $x_0 \in S$, $\mathcal{F} = \{T_\lambda | \lambda \in A\}$ a collection of affine functions on X . If \mathcal{F} is not uniformly bounded on $S \cap S(x_0, p)$, then given $M > 0$, $\varepsilon > 0$, we can find a $T \in \mathcal{F}$ and an $x \in S \cap S(x_0, p)$ such that $d(x_0, x) < \varepsilon$ and

$$d'(Tx, Tx_0) > M .$$

Proof. We can assume without loss of generality that $\varepsilon < p$. Choose $T \in \mathcal{F}$, $y \in S \cap S(x_0, p)$ such that $d'(Ty, Tx_0) > Mp/\varepsilon$. Let $x = \Phi(x_0, y, \varepsilon/p)$; $x \in S$ by the convexity of S . Now

$$d(x, x_0) = (\varepsilon/p)d(y, x_0) < \varepsilon ,$$

and

$$\begin{aligned} d'(Tx, Tx_0) &= d'(T\Phi(x_0, y, \varepsilon/p), Tx_0) \\ &= d'(\Phi(Tx_0, Ty, \varepsilon/p), Tx_0) \\ &= \varepsilon/pd'(Tx_0, Ty) > M , \end{aligned}$$

completing the proof.

The next lemma will be critical in proving the desired theorem.

LEMMA 5. Let $\{S_n | n = 1, 2, \dots\}$ be a collection of closed convex subsets of X , and let $\{T_n | n = 1, 2, \dots\}$ be a collection of affine functions on X such that $T_n|S_n$ is continuous for $n = 1, 2, \dots$. Assume that $x_{n+1} \in \bigcap_{k=1}^n S_k$ for $n = 1, 2, \dots$, and that $d(x_n, x_0)$ is sufficiently small to make $\{y_n | n = 1, 2, \dots\}$ (as defined in Definition 2) a Cauchy sequence (we do this by requiring $\sum_{n=1}^\infty M(\alpha)^{n-1}d(x_n, x_0)$ to converge). By completeness of X , let $y = \lim_{n \rightarrow \infty} y_n$. Then for each integer N , $T_N y = \lim_{n \rightarrow \infty} T_N y_n$.

Proof. Observe first that, if $\lim_{n \rightarrow \infty} u_n = u$ (in either X or Y), then $\lim_{n \rightarrow \infty} \Phi(v, u_n, \alpha) = \Phi(v, u, \alpha)$, as

$$d(\Phi(v, u_n, \alpha), \Phi(v, u, \alpha)) \leq M(\alpha)d(u_n, u) \rightarrow 0 .$$

If $n > N$, let $z_n = \Phi(x_{N+1}, \dots, \Phi(x_n, x_0, \alpha), \dots, \alpha)$. As in Lemma 1, we can show that $d(z_n, z_{n-1}) \leq (1 - \alpha)M(\alpha)^{n-N-1}d(x_n, x_0)$, and since $\sum_{n=1}^{\infty} M(\alpha)^{n-1}d(x_n, x_0)$ converges, we can define $z = \lim_{n \rightarrow \infty} z_n$. Note that $n > N \Rightarrow z_n \in S_N$, as $x_{N+1}, \dots, x_n \in S_N$ and S_N is convex. Since S_N is closed, $z \in S_N$, and so $T_N z_n \rightarrow T_N z$ by the continuity of $T_N|S_N$. If $n > N$, we have

$$\begin{aligned} T_N y_n &= T_N \Phi(x_1, \dots, \Phi(x_n, z_n, \alpha), \dots, \alpha) \\ &= \Phi(T_N x_1, \dots, \Phi(T_N x_n, T_N z_n, \alpha), \dots, \alpha) , \end{aligned}$$

and so

$$\begin{aligned} \lim_{n \rightarrow \infty} T_N y_n &= \Phi\left(T_N x_1, \dots, \lim_{n \rightarrow \infty} \Phi(T_N x_n, T_N z_n, \alpha), \dots, \alpha\right) \\ &= \Phi\left(T_N x_1, \dots, \Phi\left(T_N x_n, \lim_{n \rightarrow \infty} T_N z_n, \alpha\right), \dots, \alpha\right) \\ &= \Phi(T_N x_1, \dots, \Phi(T_N x_n, T_N z, \alpha), \dots, \alpha) . \end{aligned}$$

Since $y_n = \Phi(x_1, \dots, \Phi(x_n, z_n, \alpha), \dots, \alpha)$ and

$$\begin{aligned} y &= \lim_{n \rightarrow \infty} y_n = \Phi\left(x_1, \dots, \lim_{n \rightarrow \infty} \Phi(x_n, z_n, \alpha)\right) \\ &= \Phi(x_1, \dots, \Phi(x_n, z, \alpha), \dots, \alpha) , \end{aligned}$$

we see that $T_N y = \Phi(T_N x_1, \dots, \Phi(T_N x_n, T_N z, \alpha), \dots, \alpha) = \lim_{n \rightarrow \infty} T_N y_n$.

It is now necessary to perform some calculations. Assume

$$\{x_n | n = 1, 2, \dots\} \subseteq X ,$$

$x_0 \in X$, and $\{y_n | n = 1, 2, \dots\}$ is defined as in Definition 3. Now define

$$z_k = \Phi(x_k, \dots, \Phi(x_n, x_0, \alpha), \dots, \alpha) = \Phi(x_k, z_{k+1}, \alpha)$$

(for the purpose of these calculations, n will be assumed to be fixed) for $k \leq n - 1$, $z_n = \Phi(x_n, x_0, \alpha)$. We now have $d(x_0, \Phi(x_n, x_0, \alpha)) = d(x_0, z_n) \leq d(x_0, y_n) + \sum_{k=1}^{n-1} (d(z_k, z_{k+1}))$, as clearly $z_1 = y_n$. Observe further that

$$\begin{aligned} d(z_k, z_{k+1}) &= d(\Phi(x_k, z_{k+1}, \alpha), z_{k+1}) \\ &= (1 - \alpha)d(x_k, z_{k+1}) \\ &\leq (1 - \alpha)[d(x_k, x_0) + d(x_0, z_{k+1})] \end{aligned}$$

for $k \leq n - 1$.

We now prove some computational lemmas.

LEMMA 6. If $k \leq n - 2$,

$$d(x_0, z_{k+1}) \leq (1 + \alpha)d(x_{k+1}, x_0) + \alpha d(x_0, z_{k+2}).$$

Proof.

$$\begin{aligned} d(x_0, z_{k+1}) &\leq d(z_{k+1}, x_{k+1}) + d(x_{k+1}, x_0) \\ &= d(\Phi(x_{k+1}, z_{k+2}, \alpha), x_{k+1}) + d(x_{k+1}, x_0) \\ &= \alpha d(x_{k+1}, z_{k+2}) + d(x_{k+1}, x_0) \\ &\leq \alpha[d(x_{k+1}, x_0) + d(x_0, z_{k+2})] + d(x_{k+1}, x_0) \\ &= (1 + \alpha)d(x_{k+1}, x_0) + \alpha d(x_0, z_{k+2}). \end{aligned}$$

LEMMA 7. If $k \leq n - 2$, then

$$d(x_0, z_{k+1}) \leq (1 + \alpha) \sum_{j=0}^{n-k-2} \alpha^j d(x_0, x_{k+1+j}) + (1 - \alpha)^{n-k-1} d(x_0, x_n).$$

Proof. If $j \leq n - k - 2$, we shall verify the inequality

$$d(x_0, z_{k+1}) \leq (1 + \alpha) \sum_{i=0}^j \alpha^i d(x_0, x_{k+1+i}) + \alpha^{j+1} d(x_0, z_{k+j+2}).$$

If $j = 0$, this inequality is the conclusion of Lemma 4. Inductively, assume it is true for j . By Lemma 6, we have

$$\alpha^{j+1} d(x_0, z_{k+j+2}) \leq \alpha^{j+1} [(1 + \alpha)d(x_0, x_{k+j+2}) + \alpha d(x_0, z_{k+j+3})];$$

adding this term to the j^{th} inequality yields the inequality for $j + 1$. When $j = n - k - 2$, we therefore have

$$\begin{aligned} d(x_0, z_{k+1}) &\leq \sum_{j=0}^{n-k-2} (1 + \alpha) \alpha^j d(x_0, x_{k+1+j}) + \alpha^{n-k-1} d(x_0, z_n) \\ &= (1 + \alpha) \sum_{j=0}^{n-k-2} \alpha^j d(x_0, x_{k+1+j}) + (1 - \alpha) \alpha^{n-k-1} d(x_0, x_n). \end{aligned}$$

A consequence of Lemma 7 and a previous observation is that

$$\begin{aligned} d(z_k, z_{k+1}) &\leq (1 - \alpha)[d(x_k, x_0) + d(x_0, z_{k+1})] \\ &\leq (1 - \alpha)[d(x_k, x_0) + (1 + \alpha) \sum_{j=0}^{n-k-2} \alpha^j d(x_0, x_{k+1+j}) \\ &\quad + (1 - \alpha) \alpha^{n-k-1} d(x_0, x_n)]. \end{aligned}$$

Now let $1 \leq k \leq n - 1$. We make the following definition for $k \leq j \leq n$.

$$\begin{aligned} \mu_j^{(k)} &= 1 - \alpha && \text{if } j = k \\ &= (1 - \alpha^2) \alpha^{j-k-1} && \text{if } k < j < n \\ &= (1 - \alpha)^2 \alpha^{n-k-1} && \text{if } j = n. \end{aligned}$$

Then $d(z_k, z_{k+1}) \leq \sum_{j=k}^n \mu_j^{(k)} d(x_j, x_0)$, and so

$$\begin{aligned} (1 - \alpha)d(x_0, x_n) &= d(x_0, \Phi(x_n, x_0, \alpha)) \\ &\leq d(x_0, y_n) + \sum_{k=1}^{n-1} d(z_k, z_{k+1}) \\ &\leq d(x_0, y_n) + \sum_{k=1}^{n-1} \left(\sum_{j=k}^n \mu_j^{(k)} d(x_j, x_0) \right) \\ &= d(x_0, y_n) + \sum_{k=1}^{n-1} \left(\sum_{j=1}^k \mu_k^{(j)} \right) d(x_k, x_0) + \sum_{j=1}^{n-1} \mu_n^{(j)} d(x_n, x_0). \end{aligned}$$

If $1 \leq k \leq n - 1$, let $\beta_k = \sum_{j=1}^k \mu_k^{(j)}$, and let

$$\beta_n = \sum_{j=1}^{n-1} \mu_n^{(j)} - (1 - \alpha).$$

Obviously $\beta_k > 0$ if $1 \leq k \leq n - 1$, and also

$$\begin{aligned} \sum_{j=1}^{n-1} \mu_n^{(j)} &= (1 - \alpha)^2 \sum_{j=1}^{n-1} \alpha^{n-j-1} \\ &= (1 - \alpha)^2 \sum_{j=0}^{n-2} \alpha^j \\ &= (1 - \alpha)^2 [(1 - \alpha^{n-1}) / (1 - \alpha)] \\ &= (1 - \alpha)(1 - \alpha^{n-1}) < 1 - \alpha, \end{aligned}$$

and so $\beta_n < 0$. Since this calculation has been performed for the integer n , we shall relabel the constants just obtained $\beta_1^{(n)}, \dots, \beta_n^{(n)}$.

The last inequality proved shows that

$$0 \leq d(x_0, y_n) + \sum_{k=1}^n \beta_k^{(n)} d(x_0, x_k),$$

which implies that $d(x_0, y_n) \geq (-\beta_n^{(n)})d(x_0, x_n) - \sum_{k=1}^{n-1} \beta_k^{(n)} d(x_0, x_k)$. A reexamination of the work done subsequent to Lemma 3 shows that, if $T: X \rightarrow Y$ is affine, then

$$d'(Tx_0, Ty_n) \geq (-\beta_n^{(n)})d'(Tx_0, Tx_n) - \sum_{k=1}^{n-1} \beta_k^{(n)} d'(Tx_0, Tx_k).$$

We have therefore proved the following:

LEMMA 8. *Let $\mathcal{F} = \{T_\lambda | \lambda \in \Lambda\}$ be a pointwise-bounded family of affine functions from X into Y , and let $\{x_n | n = 1, 2, \dots\}$ be given in X , $\{y_n | n = 1, 2, \dots\}$ as in Definition 2. If*

$$S(x) = \sup \{d'(Tx, Tx_0) | T \in \mathcal{F}\},$$

then $d'(Tx_0, Ty_n) \geq (-\beta_n^{(n)})d'(Tx_0, Tx_n) - \sum_{k=1}^{n-1} \beta_k^{(n)} S(x_k)$ for any $T \in \mathcal{F}$.

Proof. Immediate from previous work and the fact that

$$d'(Tx_0, Tx_k) \leq S(x_k)$$

for all $T \in \mathcal{F}$.

We come now to the desired theorem.

THEOREM 2. *Let $(X, d), (Y, d')$ be spaces with unique segments, let X be complete, and assume there is an $\alpha \in (0, 1)$ such that $M(\alpha), M'(\alpha)$ are finite. Let $\mathcal{F} = \{T_\lambda | \lambda \in A\}$ be a pointwise-bounded family of affine maps from X into Y , and let S_λ be a closed convex subset of X such that $\bigcap_{\lambda \in A} S_\lambda \neq \varnothing$ and $T_\lambda|_{S_\lambda}$ is continuous for each $\lambda \in A$. Then $\exists \lambda_1, \dots, \lambda_n \in A$ such that \mathcal{F} is uniformly bounded on $\bigcap_{k=1}^n S_{\lambda_k}$.*

Proof. Let $x_0 \in \bigcap_{\lambda \in A} S_\lambda$, $p > 0$, and assume that \mathcal{F} is not uniformly bounded on the intersection of $S(x_0, p)$ and any finite intersection of the $\{S_\lambda | \lambda \in A\}$. We assert that we can prove the following: given $x_1, \dots, x_n \in X$, $T_1, \dots, T_n \in \mathcal{F}$ with $T_k|_{S_k}$ continuous, $1 \leq k \leq n$ and $x_k \in \bigcap_{j=1}^{k-1} S_j$ for $2 \leq k \leq n$, and given $M > 0$, let y_1, \dots, y_n be derived from x_1, \dots, x_n as in Definition 3. Then we can find $x_{n+1} \in \bigcap_{k=1}^n S_k$ and $T_{n+1} \in \mathcal{F}$ such that, if we let y_{n+1} be derived from x_1, \dots, x_{n+1} as in Definition 3,

$$d(x_0, y_{n+1}) < p, d(y_n, y_{n+1}) < 1/2^{n+1}, d'(T_{n+1}y_{n+1}, T_{n+1}x_0) > M,$$

and $d'(T_k y_n, T_k y_{n+1}) < 1/2^{n+1}$ for $1 \leq k \leq n$.

Since $x_0 \in \bigcap_{k=1}^n S_k$, choose δ_k ($1 \leq k \leq n$) such that $x \in S_k$,

$$d(x, x_0) < \delta_k \implies d'(T_k x, T_k x_0) < 1/2^{n+1}(1 - \alpha)M'(\alpha)^n;$$

then if we define $y = \Phi(x_1, \dots, \Phi(x_n, \Phi(x, x_0, \alpha), \alpha), \dots, \alpha)$, by Lemma 3 we have $x \in S_k$, $d(x, x_0) < \delta_k \implies d'(T_k y_n, T_k y) < 1/2^{n+1}$. Now let

$$\gamma = 2^{-1} \min(p, \delta_1, \dots, \delta_n, (p - d(x_0, y_n))/(1 - \alpha)M(\alpha)^n, 1/(1 - \alpha)M(\alpha)^{n2^{n+1}}).$$

Finally, by Lemma 4 choose $x_{n+1} \in \bigcap_{k=1}^n S_k$ and $T(=T_{n+1}) \in \mathcal{F}$ with $d(x_0, x_{n+1}) < \gamma$ and $(-\beta_{n+1}^{(n+1)})d'(Tx_0, Tx_{n+1}) > M + \sum_{k=1}^n \beta_k^{(n+1)}S(x_k)$. Define $y_{n+1} = \Phi(x_1, \dots, \Phi(x_{n+1}, x_0, \alpha), \dots, \alpha)$. We have already observed that $1 \leq k \leq n \implies d'(T_k y_n, T_k y_{n+1}) < 1/2^{n+1}$. Now by Lemma 3

$$d(y_n, y_{n+1}) \leq (1 - \alpha)M(\alpha)^n d(x_0, x_{n+1}) < 1/2^{n+1},$$

and also

$$\begin{aligned} d(x_0, y_{n+1}) &\leq d(x_0, y_n) + d(y_n, y_{n+1}) \\ &\leq d(x_0, y_n) + (1 - \alpha)M(\alpha)^n d(x_0, x_{n+1}) \\ &< d(x_0, y_n) + (p - d(x_0, y_n)) \\ &= p. \end{aligned}$$

By Lemma 8 we see that

$$d'(Ty_{n+1}, Tx_0) \geq (-\beta_{n+1}^{(n+1)})d'(Tx_0, Tx_{n+1}) - \sum_{k=1}^n \beta_k^{(n+1)}S(x_k) > M.$$

Construct $\{y_n | n = 1, 2, \dots\}$ by this procedure to insure that

$$d(x_0, y_{n+1}) < p, d(y_n, y_{n+1}) < 1/2^{n+1}$$

and choose $\{T_n | n = 1, 2, \dots\} \subseteq \mathcal{F}$ with $d'(T_{n+1}y_{n+1}, T_{n+1}x_0) > n + 2$ and $d'(T_k y_n, T_k y_{n+1}) < 1/2^{n+1}$ for $1 \leq k \leq n$. Now $\{y_n | n = 1, 2, \dots\}$ is Cauchy, so let $y = \lim_{k \rightarrow \infty} y_n$. By Lemma 5, for each integer n we have

$$T_n y = \lim_{m \rightarrow \infty} T_n y_m,$$

and so far any n we have $\lim_{m \rightarrow \infty} d'(T_n y, T_n y_{m+1}) = 0$. So

$$\begin{aligned} d'(T_n x_0, T_n y_n) &\leq d'(T_n x_0, T_n y) + d'(T_n y, T_n y_n) \leq \dots \\ &\leq d'(T_n x_0, T_n y) + \sum_{k=n}^m d'(T_n y_k, T_n y_{k+1}) + d'(T_n y, T_n y_{m+1}); \end{aligned}$$

as $m \rightarrow \infty$ we obtain

$$\begin{aligned} d'(T_n x_0, T_n y_n) &\leq d'(T_n x_0, T_n y) + \sum_{k=n}^{\infty} d'(T_n y_k, T_n y_{k+1}) \\ &< d'(T_n x_0, T_n y) + \sum_{k=n}^{\infty} 2^{-k-1} \\ &< d'(T_n x_0, T_n y) + 1, \end{aligned}$$

since $k \geq n \Rightarrow d'(T_n y_k, T_n y_{k+1}) < 1/2^{k+1}$. So

$$n + 1 < d'(T_n x_0, T_n y_n) \leq d'(T_n x_0, T_n y) + 1 \Rightarrow d'(T_n x_0, T_n y) > n,$$

contradicting the pointwise-boundedness of \mathcal{F} and completing the proof.

In conclusion, although spaces such that $M(\alpha)$ is infinite for every $\alpha \in (0, 1)$ are highly pathological, it would be nice to know whether or not the restriction that some $M(\alpha)$ and $M'(\alpha)$ be finite can be removed.

REFERENCES

1. G. Choquet, *Topology*, New York, Academic Press, 1966.
2. N. Dunford and J. Schwartz, *Linear Operators*, New York, Interscience, 1958.

Received March 17, 1970, and in revised form April 14, 1971. The preparation of this paper was sponsored in part by the U.S. Army Research Office under Grant DA-31-124-ARO(D)-355. Reproduction in whole or in part is permitted for any purpose of the United States Government.

UNIVERSITY OF CALIFORNIA, LOS ANGELES
 AND
 ÅRHUS UNIVERSITY, ÅRHUS, DENMARK

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 38, No. 1

March, 1971

Bruce Alan Barnes, <i>Banach algebras which are ideals in a Banach algebra</i>	1
David W. Boyd, <i>Inequalities for positive integral operators</i>	9
Lawrence Gerald Brown, <i>Note on the open mapping theorem</i>	25
Stephen Daniel Comer, <i>Representations by algebras of sections over Boolean spaces</i>	29
John R. Edwards and Stanley G. Wayment, <i>On the nonequivalence of conservative Hausdorff methods and Hausdorff moment sequences</i>	39
P. D. T. A. Elliott, <i>On the limiting distribution of additive functions (mod 1)</i>	49
Mary Rodriguez Embry, <i>Classifying special operators by means of subsets associated with the numerical range</i>	61
Darald Joe Hartfiel, <i>Counterexamples to a conjecture of G. N. de Oliveira</i>	67
C. Ward Henson, <i>A family of countable homogeneous graphs</i>	69
Satoru Igari and Shigehiko Kuratsubo, <i>A sufficient condition for L^p-multipliers</i>	85
William A. Kirk, <i>Fixed point theorems for nonlinear nonexpansive and generalized contraction mappings</i>	89
Erwin Kleinfeld, <i>A generalization of commutative and associative rings</i>	95
D. B. Lahiri, <i>Some restricted partition functions. Congruences modulo 11</i>	103
T. Y. Lin, <i>Homological algebra of stable homotopy ring π_* of spheres</i>	117
Morris Marden, <i>A representation for the logarithmic derivative of a meromorphic function</i>	145
John Charles Nichols and James C. Smith, <i>Examples concerning sum properties for metric-dependent dimension functions</i>	151
Asit Baran Raha, <i>On completely Hausdorff-completion of a completely Hausdorff space</i>	161
M. Rajagopalan and Bertram Manuel Schreiber, <i>Ergodic automorphisms and affine transformations of locally compact groups</i>	167
N. V. Rao and Ashoke Kumar Roy, <i>Linear isometries of some function spaces</i>	177
William Francis Reynolds, <i>Blocks and F-class algebras of finite groups</i>	193
Richard Rochberg, <i>Which linear maps of the disk algebra are multiplicative</i>	207
Gary Sampson, <i>Sharp estimates of convolution transforms in terms of decreasing functions</i>	213
Stephen Scheinberg, <i>Fatou's lemma in normed linear spaces</i>	233
Ken Shaw, <i>Whittaker constants for entire functions of several complex variables</i>	239
James DeWitt Stein, <i>Two uniform boundedness theorems</i>	251
Li Pi Su, <i>Homomorphisms of near-rings of continuous functions</i>	261
Stephen Willard, <i>Functionally compact spaces, C-compact spaces and mappings of minimal Hausdorff spaces</i>	267
James Patrick Williams, <i>On the range of a derivation</i>	273