

Pacific Journal of Mathematics

NORMS OF DERIVATIONS OF $\mathcal{L}(X)$

BARRY E. JOHNSON

NORMS OF DERIVATIONS ON $\mathcal{L}(\mathfrak{X})$

B. E. JOHNSON

If \mathfrak{X} is a real or complex Banach space and $\mathcal{L}(\mathfrak{X})$ is the algebra of bounded linear endomorphisms of \mathfrak{X} then each element T of $\mathcal{L}(\mathfrak{X})$ defines an operator D_T on $\mathcal{L}(\mathfrak{X})$ by $D_T(A) = AT - TA$. Clearly $\|D_T\| \leq 2 \inf_{\lambda \in \mathbf{K}} \|T + \lambda I\|$ and Stampfli has shown that when \mathfrak{X} is a complex Hilbert space equality holds. In this paper it is shown, by methods which apply to a large class of uniformly convex spaces, that this formula for $\|D_T\|$ is false in l^p and $L^p(0, 1)$ $1 < p < \infty$, $p \neq 2$. For L^1 spaces the formula is true in the real case but not in the complex case when the space has dimension 3 or more.

Stampfli's results appear in [1] stated for complex Hilbert space but the same proofs yield the corresponding result for real spaces.

Throughout this paper K will denote either \mathbf{R} or \mathbf{C} . We begin by describing the construction of an operator T of rank 1 with $\|D_T\| < d_T = 2 \inf_{\lambda \in K} \|T + \lambda I\|$. The reason that this fails in Hilbert space is precisely because for an ellipse, conjugacy is a symmetric relation on the set of diameters; more precisely if x, y are two points on the unit ball then y is parallel to the tangent plane at x if and only if x is parallel to the tangent plane at y .

DEFINITION 1. Let $x \in \mathfrak{X}$, $\|x\| = 1$. The unit ball \mathfrak{X}_1 is *uniformly convex* at x if whenever $\{y_n\}$ is a sequence with $\|y_n\| \leq 1$, $\|x + y_n\| \rightarrow 2$ then $y_n \rightarrow x$.

PROPOSITION 2. Let \mathfrak{X} be a normed space over K and let $x, y \in \mathfrak{X}$ with the following properties

- (i) $\|x\| = 1$ and there is $f \in \mathfrak{X}^*$ with $\|f\| = 1$ and such that if $\{x_n\}$ is a sequence with $\|x_n\| \leq 1$, $f(x_n) \rightarrow 1$ then $x_n \rightarrow x$.
- (ii) $\|y\| = 1$ and the unit ball \mathfrak{X}_1 is uniformly convex at y .
- (iii) For some $\lambda \in K$, $\|x + \lambda y\| < 1$.
- (iv) For all λ in K , $\|y + \lambda x\| \geq 1$.

Define $T \in \mathcal{L}(\mathfrak{X})$ by $Tz = f(z)y$. Then $2\|T\| = d_T > \|D_T\|$.

Proof. $\|T + \lambda I\| \geq \|(T + \lambda I)x\| = \|y + \lambda x\| \geq \|y\| = 1$ by (iv) and $\|T\| = 1$ so $d_T = 2$. Suppose $\|D_T\| = 2$ and choose sequences $\{A_n\}$ from $\mathcal{L}(\mathfrak{X})$ and $\{x_n\}$ from \mathfrak{X} with $\|A_n\| = 1 = \|x_n\|$ and $\|D_T(A_n)x_n\| \rightarrow 2$. As $\|TA_nx_n\| \leq 1$, $\|A_nTx_n\| \leq 1$ we have $\|TA_nx_n\| \rightarrow 1$, $\|A_nTx_n\| \rightarrow 1$

and hence $\|A_n x_n\| \rightarrow 1$, $\|Tx_n\| \rightarrow 1$. This shows $|f(x_n)| \rightarrow 1$ and so, replacing x_n by $w_n x_n$ if necessary where $\{w_n\}$ is a sequence of elements of K with $|w_n| = 1$, we may assume $f(x_n) \rightarrow 1$. Condition (i) now implies $x_n \rightarrow x$ and hence $Tx_n \rightarrow y$. In the same way $\|TA_n x_n\| \rightarrow 1$ implies $|f(A_n x_n)| \rightarrow 1$ and replacing A_n by $w'_n A_n$ if necessary we can assume $f(A_n x_n) \rightarrow 1$ from which we see $A_n x_n \rightarrow x$, $TA_n x_n \rightarrow y$. As $\|A_n\| \leq 1$ we have $A_n Tx_n - A_n y \rightarrow 0$ and so $\|A_n Tx_n - TA_n x_n\| \rightarrow 2$ implies $\|A_n y - y\| \rightarrow 2$. Condition (ii) now shows $A_n y \rightarrow -y$ so that $A_n(x + \lambda y) \rightarrow x - \lambda y$. However if λ satisfies condition (iii) then $\|x - \lambda y\| > 1$, as otherwise $2 = 2\|x\| \leq \|x + \lambda y\| + \|x - \lambda y\| < 2$, and so $\lim \|A_n(x + \lambda y)\| = \|x - \lambda y\| > 1$ which is impossible because $\|A_n(x + \lambda y)\| \leq \|A_n\| \|x + \lambda y\| < 1$.

PROPOSITION 3. *Let \mathfrak{X} be a uniformly convex Banach space, $x, y \in \mathfrak{X}$, $f, g \in \mathfrak{X}^*$ with $\|x\| = \|y\| = \|f\| = \|g\| = f(x) = g(y) = 1$, $g(x) = 0$, $f(y) \neq 0$ and suppose f is the only element h of \mathfrak{X}^* with $\|h\| = h(x) = 1$. Then x, y, f satisfy the conditions of Proposition 2.*

Proof. (i) If $\|x_n\| \leq 1$, $f(x_n) \rightarrow 1$ then $f(x + x_n) \rightarrow 2$ and as $\|x + x_n\| \leq 2$, $\|f\| = 1$ we have $\|x + x_n\| \rightarrow 2$ so $x_n \rightarrow x$ by uniform convexity.

(ii) is clearly part of the present hypotheses.

(iii) x and y are linearly independent as $g(x) = 0$, $g(y) = 1$, $x \neq 0$. If $\|x + \lambda y\| \geq 1$ for all $\lambda \in K$ then $\alpha x + \beta y \mapsto \alpha$ is a norm one linear functional on the space spanned by x and y and so has an extension h in \mathfrak{X}^* with $\|h\| = 1$, $h(x) = 1$ but $h \neq f$ because $h(y) = 0 \neq f(y)$.

(iv) As $g(y + \lambda x) = g(y) = 1$, for all λ in K and $\|g\| = 1$ we have $\|y + \lambda x\| \geq 1$ for all λ in K .

COROLLARY 4. *If $1 < p < 2$ or $2 < p < \infty$ and $\mathfrak{X} = l^p(0, \infty)$ or $\mathfrak{X} = L^p(-1, +1)$ is the corresponding K Banach space of K valued functions then there is $T \in \mathcal{L}(\mathfrak{X})$ with $\|D_T\| \neq d_T$.*

Proof. The spaces are uniformly convex and at each point z of \mathfrak{X} with $\|z\| = 1$ the element h of \mathfrak{X}^* with $h(z) = 1 = \|h\|$ is unique. Thus the construction in Proposition 2 applies once we find two suitable points x, y and these exist in such abundance that we can take anything but multiples of characteristic functions for x . First of all we give the construction in the two dimensional space $l^p(1, 2)$.

If $x = (x_1, x_2)$, $x_1 > 0$, $x_2 > 0$, $x_1^p + x_2^p = 1$ then $f(z) = x_1^{p-1}z_1 + x_2^{p-1}z_2$ so y can be taken as $\alpha(x_2^{p-1}, -x_1^{p-1})$ where $\alpha^{-p} = x_1^{p(p-1)} + x_2^{p(p-1)}$ and

$g(z) = \alpha^{p-1}(x_2^{(p-1)^2} z_1 - x_1^{(p-1)^2} z_2)$. Then $g(x) = \alpha^{p-1}(x_1 x_2^{(p-1)^2} - x_1^{(p-1)^2} x_2)$ which will be zero if and only if $x_1 = x_2$. Thus taking say $x = 3^{-1/p}(2^{1/p}, 1)$ and y, f, g as above the result is shown in $l^p(1, 2)$.

As $l^p(0, \infty)$ and $L^p(-1, +1)$ each contain subspaces isometric with $l^p(1, 2)$ we can construct x, y, f, g in this subspace and then extend f and g to \mathfrak{X} using the Hahn-Banach theorem.

In order to prove the results for spaces of measures we establish the equation $d_T = \|D_T\|$ for finite dimensional l^1 spaces.

PROPOSITION 5. *Let n be a positive integer and \mathfrak{X} be the real Banach space \mathbf{R}^n with norm $\|x\| = \sum |x_i|$. Let $T \in \mathcal{L}(\mathfrak{X})$. Then $\|D_T\| = 2 \inf_{\lambda \in \mathbf{R}} \|T + \lambda I\|$.*

Proof. Suppose T is given by the matrix a_{ij} in the standard basis e_1, e_2, \dots, e_n . We have $\|T\| = \sup_j \sum_i |a_{ij}|$. Suppose $\sum_i |a_{ij}| = \|T\|$ for $j = 1, \dots, m$ but not for $j > m$. The condition $\|T\| = \frac{1}{2} d_T$ is equivalent to saying that 0 is in the convex hull of a_{11}, \dots, a_{mm} since if 0 does not lie in this convex hull then either $|a_{jj} + \lambda| < |a_{jj}|$ for $j = 1, \dots, m$ and small positive λ or for small negative λ and so there are small values of λ with $\|T + \lambda I\| < \|T\|$ whereas if 0 lies in this hull and $\lambda \neq 0$ there is j with $1 \leq j \leq m$ and $|a_{jj} + \lambda| > |a_{jj}|$ so that $\|T + \lambda I\| > \|T\|$.

It is clearly sufficient to prove the result when $\|T\| = \frac{1}{2} d_T$. First of all consider the case $m \geq 2$ and suppose $a_{11} \geq 0 \geq a_{22}$. Let $A \in \mathcal{L}(\mathfrak{X})$ be an operator of the form $Ae_1 = e_2, Ae_2 = \pm e_1, Ae_i = \pm e_i$ $i = 3, \dots, n$. Clearly $\|A\| = 1$ and

$$\begin{aligned} \|D_T(A)e_1\| &= \|ATE_1 - Te_2\| \\ &= |\pm a_{21} - a_{12}| + |a_{11} - a_{22}| + \sum_{i=3}^n |\pm a_{i1} - a_{i2}| \\ &= \sum_{i=1}^n |a_{i1}| + \sum_{i=1}^n |a_{i2}| \\ &= 2 \|T\| \end{aligned}$$

for a suitable choice of signs of the Ae_i since each sign to be chosen corresponds to exactly one term $|\pm a_{i1} - a_{i2}|$.

If $m = 1$ then $a_{11} = 0$ because 0 lies in the convex hull of a_{11}, \dots, a_{mm} , and we define A by $Ae_1 = e_1, Ae_j = -e_j$ $j = 2, \dots, n$ which gives $\|A\| = 1$ and $ATE_1 = -Te_1$ so that

$$\|D_T(A)e_1\| = \|ATE_1 - TAe_1\| = 2 \|Te_1\| = 2 \|T\|.$$

PROPOSITION 6. *Let Ω be a compact topological space and \mathfrak{X} a closed linear subspace of the (real) Banach space of real valued measures on Ω with the property that if $\mu \in \mathfrak{X}$ then every measure*

absolutely continuous with respect to μ is in \mathfrak{X} . Let $T \in \mathcal{L}(\mathfrak{X})$. Then $\|D_T\| = 2 \inf_{\lambda \in \mathbf{R}} \|T + \lambda I\|$.

Proof. We may assume $d_T = 2 \|T\|$. Let $\varepsilon > 0$. For each $\nu > 0$ in \mathfrak{X} let $E_\nu(\mu)$ be the part of $\mu \in \mathfrak{X}$ which is absolutely continuous with respect to ν . The E_ν form a system of commuting idempotents of norm 1 and $E_\nu E_{\nu'} = E_\nu$ if $\nu' > \nu$, so that $\|E_\nu S E_\nu\|$, where the elements ν are directed by the usual ordering of measures, is a monotonic direct net. It is easy to see that $\|E_\nu S E_\nu\| \rightarrow \|S\|$. Thus applying Dini's theorem to the functions $\lambda \mapsto \|E_\nu(T + \lambda I)E_\nu\|$ we can find $\nu \in \mathfrak{X}, \nu > 0$ with $\|E_\nu(T + \lambda I)E_\nu\| > \|T + \lambda I\| - \varepsilon \geq \|T\| - \varepsilon$ for $|\lambda| \leq 2 \|T\|$.

For each dissection $\Delta = (\Omega_1, \dots, \Omega_n)$ of Ω into disjoint measurable sets of positive ν measure we define

$$P_\Delta(\mu) = (E_\nu \mu(\Omega_1), \dots, E_\nu \mu(\Omega_n))$$

$$Q_\Delta(\xi) = (\sum c_i \xi_i \nu(\Omega_i)^{-1}) \nu$$

where $\mu \in \mathfrak{X}, \xi \in \mathbf{R}^n, P_\Delta: \mathfrak{X} \rightarrow \mathbf{R}^n, Q_\Delta: \mathbf{R}^n \rightarrow \mathfrak{X}$ and c_i is the characteristic function of Ω_i . Directing the dissections in the usual way it is easy to see that for each $S \in \mathcal{L}(\mathfrak{X})$ $\|P_\Delta E_\nu S E_\nu Q_\Delta\|$, where \mathbf{R}^n has the l^1 norm, is a monotonic directed set with limit $\|E_\nu S E_\nu\|$. Applying Dini's theorem again we see that there is a dissection Δ with

$$(*) \quad \|P_\Delta E_\nu(T + \lambda I)E_\nu Q_\Delta\| > \|T\| - \varepsilon$$

for all $|\lambda| \leq 2 \|T\|$. For convenience we now denote $E_\nu, P_\Delta, Q_\Delta$ by E, P, Q . As these operators have norm 1 we see that inequality (*) holds for all values of λ . As $PE = P, EQ = Q, PEQ = PQ =$ identity on \mathbf{R}^n , (*) shows that $d_{PTQ} \geq 2(\|T\| - \varepsilon)$. By proposition 5 there is $A \in \mathcal{L}(\mathbf{R}^n)$ with $\|D_{PTQ}(A)\| = d_{PTQ}, \|A\| = 1$. As Q is an isometry and P maps the unit ball of \mathfrak{X} onto that of \mathbf{R}^n we have

$$d_{PTQ} = \|QD_{PTQ}(A)P\|$$

$$= \|QAPTQP - QPTQAP\|$$

$$= \|QPD_T(QAP)QP\|$$

$$\leq \|D_T(QAP)\|.$$

As $\|QAP\| = 1$ we have $\|D_T\| \geq d_{PTQ} \geq 2(\|T\| - \varepsilon)$ for each $\varepsilon > 0$ and the result follows.

In the complex space $l^1(1, 2)$ Proposition 5 is true and the proof is similar to that for the real case. However the result is false in higher dimensions for complex spaces, e.g., in $l^1(1, 2, 3)$ let T be the linear transformation given by the matrix

$$\begin{matrix} 1 & -\omega & -\omega^2 \\ 1 & \omega & -\omega^2 \\ 1 & \omega & \omega^2 \end{matrix}$$

where $\omega^3 = 1, \omega \neq 1$. The situation is similar to that at the beginning of the proof of Proposition 5 with $m = n = 3$ and the argument given there shows that because 0 is a convex combination of diagonal entries we have $\inf_{\lambda \in \mathbb{C}} \|T + \lambda I\| = \|T\| = 3$. If $\|x\| = 1, \|A\| = 1$ and $\|D_T(A)x\| = 6$ then $\|Tx\| = 3$ and since $|x_1 \pm \omega x_2 \pm \omega^2 x_3| \leq 1$ we see that $|x_1 - \omega x_2 - \omega^2 x_3| = |x_1 + \omega x_2 - \omega^2 x_3| = |x_1 + \omega x_2 + \omega^2 x_3| = |x_1| + |x_2| + |x_3|$ which occurs only if two of x_1, x_2, x_3 are 0. Multiplying by a complex number of absolute value 1, if necessary, we can assume $x = e_1$ or e_2 or e_3 . In the same way $Ax = e_1$ or e_2 or e_3 . If $x = e_1 = Ax$ then

$$\begin{aligned} \|D_T(A)e_1\| &= \|e_1 + Ae_2 + Ae_3 - e_1 - e_2 - e_3\| \\ &= \|Ae_2 + Ae_3 - e_2 - e_3\| \\ &\leq 4 \end{aligned}$$

and if $x = e_1, Ax = e_2$ then

$$\begin{aligned} \|D_T(A)e_1\| &= \|e_2 + Ae_2 + Ae_3 + \omega e_1 - \omega e_2 - \omega e_3\| \\ &= \|(1 - \omega)e_2 + Ae_2 + Ae_3 - \omega e_1 - \omega e_3\| \\ &\leq \sqrt{3} + 4. \end{aligned}$$

The other four possibilities give similar results and so we cannot in fact have $\|D_T\| = 6$.

A similar construction in the complex spaces $l^1(1, n), l^1(0, \infty), L^1(0, 1), M(0, 1)$ shows that Proposition 6 is false in these spaces too.

REFERENCE

1. J. G. Stampfli, *On the norm of a derivation*, Pacific J. Math., **33** (1970), 737-747.

Received November 2, 1970. The author gratefully acknowledges financial support from the National Science Foundation Grant GP-21193.

YALE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics

Vol. 38, No. 2

April, 1971

Richard Davis Anderson and Thomas Ashland Chapman, <i>Extending homeomorphisms to Hilbert cube manifolds</i>	281
Nguyen Huu Anh, <i>Restriction of the principal series of $SL(n, \mathbf{C})$ to some reductive subgroups</i>	295
David W. Boyd, <i>Indices for the Orlicz spaces</i>	315
William Garfield Bridges, <i>The polynomial of a non-regular digraph</i>	325
Billie Chandler Carlson, Robert K. Meany and Stuart Alan Nelson, <i>Mixed arithmetic and geometric means</i>	343
H. A. Çelik, <i>Commutative associative rings and anti-flexible rings</i>	351
Hsin Chu, <i>On the structure of almost periodic transformation groups</i>	359
David Allyn Drake, <i>The translation groups of n-uniform translation Hjelmslev planes</i>	365
Michael Benton Freeman, <i>The polynomial hull of a thin two-manifold</i>	377
Anthony Alfred Gioia and Donald Goldsmith, <i>Convolutions of arithmetic functions over cohesive basic sequences</i>	391
Leslie C. Glaser, <i>A proof of the most general polyhedral Schoenflies conjecture possible</i>	401
Thomas Lee Hayden and Ted Joe Suffridge, <i>Biholomorphic maps in Hilbert space have a fixed point</i>	419
Roger Alan Horn, <i>Schlicht mappings and infinitely divisible kernels</i>	423
Norman Ray Howes, <i>On completeness</i>	431
Hideo Imai, <i>Sario potentials on Riemannian spaces</i>	441
A. A. Iskander, <i>Subalgebra systems of powers of partial universal algebras</i>	457
Barry E. Johnson, <i>Norms of derivations of $\mathcal{L}(X)$</i>	465
David Clifford Kay and Eugene W. Womble, <i>Axiomatic convexity theory and relationships between the Carathéodory, Helly, and Radon numbers</i>	471
Constantine G. Lascarides, <i>A study of certain sequence spaces of Maddox and a generalization of a theorem of Iyer</i>	487
C. N. Linden, <i>On Blaschke products of restricted growth</i>	501
John S. Lowndes, <i>Some triple integral equations</i>	515
Declan McCartan, <i>Bicontinuous preordered topological spaces</i>	523
S. Moedomo and J. Jerry Uhl, Jr., <i>Radon-Nikodým theorems for the Bochner and Pettis integrals</i>	531
Calvin Cooper Moore and Joseph Albert Wolf, <i>Totally real representations and real function spaces</i>	537
Reese Trego Prosser, <i>A form of the moment problem for Lie groups</i>	543
Larry Smith, <i>A note on annihilator ideals of complex bordism classes</i>	551
Joseph Warren, <i>Meromorphic annular functions</i>	559