

Pacific Journal of Mathematics

ON THE DENSITY OF (k, r) INTEGERS

Y. K. FENG AND M. V. SUBBA RAO

ON THE DENSITY OF (k, r) INTEGERS

Y. K. FENG AND M. V. SUBBARAO

Let k and r be integers such that $0 < r < k$. We call a positive integer n , $a(k, r)$ -integer if it is of the form $n = a^k b$, where a and b are natural numbers and b is r -free. Clearly, $a(\infty, r)$ -integer is a r -free integer. Let $Q_{k,r}$ denote the set of (k, r) -integers and let $\delta(Q_{k,r})$, $D(Q_{k,r})$ respectively denote the asymptotic and Schnirelmann densities of the set $Q_{k,r}$. In this paper, we prove that $\delta(Q_{k,r}) > D(Q_{k,r}) \cong \zeta(k)(1 - \sum_p p^{-r}) - 1/k(1 - (1/k))^{k-1}$, and deduce the known results for r -free integers.

1. Introduction and Notation. In some recent papers, ([4, 5]) we introduced a generalized class of r -free integers, which we called the (k, r) -integers. For given integers k, r with $0 < r < k$, $a(k, r)$ -integer is one whose k -free part is also r -free. In the limiting case when $k = \infty$, we get the r -free integers. It is clear that $a(k, r)$ -integer is an integer of the form $a^k b$, where a and b are natural numbers and b is r -free. Let $Q_{k,r}, Q_r$ denote the set of all (k, r) -integers and the set of all r -free integers respectively. Also let $Q_{k,r}(x)$ denote the number of (k, r) -integers not exceeding x , with corresponding meaning for $Q_r(x)$. We write $\delta(Q_{k,r})$ for the asymptotic density of the (k, r) -integers, that is,

$$\delta(Q_{k,r}) = \lim_{x \rightarrow \infty} \frac{Q_{k,r}(x)}{x},$$

(provided this limit exists), and $D(Q_{k,r})$ for their Schnirelmann density given by

$$D(Q_{k,r}) = \inf_n \frac{Q_{k,r}(n)}{n}.$$

We define $\delta(Q_r)$ and $D(Q_r)$ analogously. Let $\psi(n)$ be the characteristic function of $Q_{k,r}$ and $\lambda(n)$ be defined by

$$\sum_{d|n} \lambda(d) = \psi(n).$$

It is easily proved (see [3]) that the function $\psi(n)$ and $\lambda(n)$ are multiplicative and for any prime p

$$\lambda(p^a) = \begin{cases} 1 & a \equiv 0 \pmod{k}, \\ -1 & a \equiv r \pmod{k}, \\ 0 & \text{otherwise.} \end{cases}$$

Further,

$$(1.1) \quad \sum_{n=1}^{\infty} \frac{\lambda(n)}{n^s} = \frac{\zeta(ks)}{\zeta(rs)}, \quad \operatorname{Re}(s) > \frac{1}{r},$$

where $\zeta(s)$ is the Riemann Zeta function. In a previous paper [5], we showed that

$$Q_{k,r}(x) = \frac{x^{\zeta(k)}}{\zeta(r)} + E(x),$$

where the error term $E(x)$ is $O(x^{\frac{1}{r}})$, for $r > 1$, uniformly in k . (We actually gave an improved estimate for the error term, but this is not required here.)

It follows that

$$\delta(Q_{k,r}) = \frac{\zeta(k)}{\zeta(r)}.$$

In this paper we will show that

$$\delta(Q_{k,r}) > D(Q_{k,r}).$$

The corresponding result for Q_2 was first proved by Rogers [2], and for Q_r , for all $r > 1$ by Stark [6]. We also obtain a lower bound for $D(Q_{k,r})$, from which we obtain as a special case a result of Duncan [1] on a lower bound for $D(Q_r)$. The actual value of $D(Q_{k,r})$ is unknown except for the case Q_2 ; Rogers [3] proved that

$$D(Q_2) = \frac{53}{88}.$$

2. Theorem.

$$\delta(Q_{k,r}) > D(Q_{k,r}) \geq \zeta(k) \left(1 - \sum_p p^{-r}\right) - \frac{1}{k} \left(1 - \frac{1}{k}\right)^{k-1}.$$

The proof will be given in two parts, corresponding to the two results:

$$(2.1) \quad D(Q_{k,r}) \geq \zeta(k) \left(1 - \sum_p p^{-r}\right) - \frac{1}{k} \left(1 - \frac{1}{k}\right)^{k-1};$$

$$(2.2) \quad \delta(Q_{k,r}) > D(Q_{k,r}).$$

Proof of (2.1). The case $r > 1$.

It is clear that

$$Q_r(n) \geq n - \sum_p \left[\frac{n}{p^r} \right],$$

p ranging over all the primes.

Since

$$Q_{k,r}(n) = \sum_{a=1}^{\infty} Q_r\left(\left[\frac{n}{a^k}\right]\right),$$

it follows that

$$\begin{aligned} Q_{k,r}(n) &\geq \sum_{a=1}^{\infty} \left(\left[\frac{n}{a^k}\right] - \sum_p \left[\frac{a^{n/k}}{p^r}\right] \right) \\ &> \sum_{a=1}^{\infty} \left(\frac{n}{a^k} - \sum_p \frac{n}{a^k p^r} \right) - (n^{1/k} - 1). \end{aligned}$$

Hence we have

$$\begin{aligned} \frac{Q_{k,r}(n)}{n} &> \sum_{a=1}^{\infty} \left(\frac{1}{a^k} - \sum_p \frac{1}{a^k p^r} \right) - \frac{n_{k/1} - 1}{n} \\ &= \zeta(k)(1 - \sum_p^{P-r}) + \frac{1 - n^{1/k}}{n}. \end{aligned}$$

Let

$$f(x) = \frac{1 - x^{1/k}}{x} = \frac{1}{x} - x^{1/k-1};$$

then

$$f'(x) = \frac{1}{x^2} - \left(\frac{1}{k} - 1\right)x^{1/k-2},$$

so that

$$f'(x) > 0 \text{ if } \left(1 - \frac{1}{k}\right)x^{(1/k)-2} > \frac{1}{x^2}, \text{ i.e., } \left(1 - \frac{1}{k}\right)x^{1/k} > 1.$$

Thus

$$f'(x) \begin{cases} > 0 \text{ when } x > \frac{1}{\left(1 - \frac{1}{k}\right)^k}, \\ < 0 \text{ when } x < \frac{1}{\left(1 - \frac{1}{k}\right)^k}. \end{cases}$$

when $x = (1 - (1/k))^{-k}$ we get the minimum value of f , which is equal to

$$f\left(\left(1 - \frac{1}{k}\right)^{-k}\right) = \frac{1 - \left(1 - \frac{1}{k}\right)^{-1}}{\left(1 - \frac{1}{k}\right)^{-k}} = -\frac{1}{k} \left(1 - \frac{1}{k}\right)^{k-1}.$$

Hence

$$\frac{Q_{k,r}(n)}{n} > \zeta(k)(1 - \sum_p p^{-r}) - \frac{1}{k} \left(1 - \frac{1}{k}\right)^{k-1},$$

and

$$D(Q_{k,r}) \geq \zeta(k)(1 - \sum_p p^{-r}) - \frac{1}{k} \left(1 - \frac{1}{k}\right)^{k-1}.$$

For the case $r = 1$,

$$\begin{aligned} Q_{k,1}(n) &= [n^{1/k}] \\ \delta(Q_{k,1}) &= \lim_{n \rightarrow \infty} \frac{[n^{1/k}]}{n} = 0, \text{ since } k \geq 2. \\ D(Q_{k,1}) &= \inf_n \frac{[n^{1/k}]}{n} = 0. \end{aligned}$$

So the result still holds in this case.

REMARK 2.3. The above proof is easily seen to hold even when $k = \infty$. The corresponding result, namely,

$$D(Q_r) > 1 - \sum_p p^{-r},$$

is due to R. L. Duncan [1].

To prove the result in (2.2), we first obtain the following lemma.

LEMMA 2.4. For any $\varepsilon > 0$, we have

- (i) $E(n) > n^{(1/2r)-\varepsilon}$, for infinitely many integers n ,
- (ii) $E(n) < -n^{(1/2r)-\varepsilon}$, for infinitely many integers n .

Proof. Let

$$\sum \left(\psi(n) - \frac{\zeta(k)}{\zeta(r)} \right) n^{-s} = R_1(s).$$

Since

$$\sum \left(\psi(n) - \frac{\zeta(k)}{\zeta(r)} \right) n^{-s} = \frac{\zeta(ks)\zeta(s)}{\zeta(rs)} - \frac{\zeta(k)\zeta(s)}{\zeta(r)},$$

we have

$$\begin{aligned} R_1(s) &= \sum \left(\psi(n) - \frac{\zeta(k)}{\zeta(r)} \right) n^{-s} \\ &= \sum (E(n) - E(n-1))n^{-s} \\ &= \sum E(n)(n^{-s} - (n+1)^{-s}). \end{aligned}$$

Also, let

$$\begin{aligned} s \sum E(n)n^{-s-1} &= R_2(s) , \\ \sum E(n)n^{-s-1} &= R_3(s) , \\ \sum n^{(1/2r)-\varepsilon} \cdot n^{-s-1} &= R_4(s) , \\ \sum (n^{(1/2r)-\varepsilon} - E(n))n^{-s-1} &= R_5(s) , \\ \sum (n^{(1/2r)-\varepsilon} + E(n))n^{-s-1} &= R_6(s) . \end{aligned}$$

Now suppose that for all $n \geq n_0$, $E(n) \leq n^{(1/2r)-\varepsilon}$. Then the series $R_5(s)$ converges for $a > (1/2r) - \varepsilon$ ($a = \operatorname{Re}(s)$), and all but a finite number of coefficients of $R_5(s)$ are nonnegative. Hence the abscissa of convergence of $R_5(s)$ must be less than or equal to $(1/2r) - \varepsilon$. Let α be its abscissa of convergence, that is $\alpha \leq (1/2r) - \varepsilon$. Note that (see [2], P. 661)

$$|n^{-s} - (n+1)^{-s} - s \cdot n^{-s-1}| \leq |s| |s+1| n^{-a-2} .$$

This implies $R_1(s)$ also converges for $a > \alpha$. But this is false because $R_1(s)$ has singularities on $a = (1/2r)$. Thus we must have

$$E(n) > n^{(1/2r)-\varepsilon}$$

for infinitely many integers n .

Next suppose that for all $n \geq n_0$, $E(n) \geq -n^{(1/2r)-\varepsilon}$, then we consider the series $R_6(s)$, proceed as in (i) and arrive at the same contradiction.

Proof of the result (2.2). By the above lemma, there are infinitely many integers n for which $E(n) < 0$. For such n ,

$$\frac{Q_{k,r}(n)}{n} = \frac{\zeta(k)}{\zeta(r)} + \frac{E(n)}{n} < \frac{\zeta(k)}{\zeta(r)} ,$$

which proves the theorem.

REFERENCES

1. R. L. Duncan, *The Schnirelmann density of the k -free integers*, Proc. Amer. Math. Soc., **16** (1965), 1090-1091.
2. E. Landau, *Handbuch der Lehre von der Verteilung der primzahlen*, Chelsea (reprint), 1953.
3. K. Rogers, *The Schnirelmann density of the square-free integers*, Proc. Amer. Math. Soc., **15** (1964), 515-516.
4. M. V. Subbarao, and V. C. Harris, *A new generalization of Ramanujan's sum*, J. London Math. Soc., **41** (1966), 595-604.
5. M. V. Subbarao, and Y. K. Feng, *On the distribution of generalized r -free integers in residue classes*, (To appear in Duke Math.).

6. H. M. Stark *On the asymptotic density of the k -free integers*, Proc. Amer. Math. Soc., 17 (1966), 1211-1214.

Received July 28, 1970 and in revised form January 26, 1971.

UNIVERSITY OF ALBERTA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics

Vol. 38, No. 3

May, 1971

J. T. Borrego, Haskell Cohen and Esmond Ernest Devun, <i>Uniquely representable semigroups on the two-cell</i>	565
Glen Eugene Bredon, <i>Some examples for the fixed point property</i>	571
William Lee Bynum, <i>Characterizations of uniform convexity</i>	577
Douglas Derry, <i>The convex hulls of the vertices of a polygon of order n</i>	583
Edwin Duda and Jack Warren Smith, <i>Reflexive open mappings</i>	597
Y. K. Feng and M. V. Subba Rao, <i>On the density of (k, r) integers</i>	613
Irving Leonard Glicksberg and Ingemar Wik, <i>Multipliers of quotients of L_1</i>	619
John William Green, <i>Separating certain plane-like spaces by Peano continua</i>	625
Lawrence Albert Harris, <i>A continuous form of Schwarz's lemma in normed linear spaces</i>	635
Richard Earl Hodel, <i>Moore spaces and w Δ-spaces</i>	641
Lawrence Stanislaus Husch, Jr., <i>Homotopy groups of PL-embedding spaces. II</i>	653
Yoshinori Isomichi, <i>New concepts in the theory of topological space—supercondensed set, subcondensed set, and condensed set</i>	657
J. E. Kerlin, <i>On algebra actions on a group algebra</i>	669
Keizō Kikuchi, <i>Canonical domains and their geometry in C^n</i>	681
Ralph David McWilliams, <i>On iterated w^*-sequential closure of cones</i>	697
C. Robert Miers, <i>Lie homomorphisms of operator algebras</i>	717
Louise Elizabeth Moser, <i>Elementary surgery along a torus knot</i>	737
Hiroshi Onose, <i>Oscillatory properties of solutions of even order differential equations</i>	747
Wellington Ham Ow, <i>Wiener's compactification and Φ-bounded harmonic functions in the classification of harmonic spaces</i>	759
Zalman Rubinstein, <i>On the multivalence of a class of meromorphic functions</i>	771
Hans H. Storrer, <i>Rational extensions of modules</i>	785
Albert Robert Stralka, <i>The congruence extension property for compact topological lattices</i>	795
Robert Evert Stong, <i>On the cobordism of pairs</i>	803
Albert Leon Whiteman, <i>An infinite family of skew Hadamard matrices</i>	817
Lynn Roy Williams, <i>Generalized Hausdorff-Young inequalities and mixed norm spaces</i>	823