# Pacific Journal of Mathematics

## GENERALIZED HAUSDORFF-YOUNG INEQUALITIES AND MIXED NORM SPACES

LYNN ROY WILLIAMS

Vol. 38, No. 3

## GENERALIZED HAUSDORFF-YOUNG INEQUALITIES AND MIXED NORM SPACES

## LYNN R. WILLIAMS

We generalize the Hausdorff-Young Theorem for a locally compact connected group G by showing that if  $f \in L^p(G)$ , 1 , then the Fourier transform of f is in a mixed norm $space properly contained in <math>L^{p'}(\Gamma)$ , where  $\Gamma$  is the dual group and 1/p + 1/p' = 1. In the last section we apply the above theorem to obtain new results concerning sets of uniqueness for functions in  $L^p(G)$ , and we give new sufficient conditions which insure that the product of a continuous function and a pseudomeasure is the zero distribution.

1. Introduction. The classical Hausdorff-Young Theorem states that if f belongs to the Lebesgue space  $L^p$  (of the unit circle),  $1 and <math>\{f_n\}$  is its sequence of Fourier coefficients, then

(1.1) 
$$\left\{\sum_{n=-\infty}^{\infty} |f_n|^{p'}\right\}^{1/p'} \leq ||f||_p \text{ where } \frac{1}{p} + \frac{1}{p'} = 1.$$

The companion dual result asserts that a sufficient condition for the sequence  $\{f_n\}$  to be the Fourier coefficients of a function in  $L^{p'}$  is

(1.2) 
$$\left\{\sum_{n=-\infty}^{\infty}|f_n|^p\right\}^{1/p}<\infty$$

In a recent paper [7] Kellogg applied a multiplier theorem of Hedlund to obtain a significant improvement of these inequalities. Precisely, he replaced (1.1) by

(1.3) 
$$\left\{\sum_{k=-\infty}^{\infty} \left(\sum_{n \in I_k} |f_n|^{p'}\right)^{2/p'}\right\}^{1/2} \leq A_p ||f||_p$$

and (1.2) by

(1.4) 
$$\left\{\sum_{k=-\infty}^{\infty}\left(\sum_{n\in I_k}|f_n|^p\right)^{2/p}\right\}^{1/2} < \infty$$

where  $A_p$  is a constant depending only on p and the sets  $I_k$  are the lacunary blocks defined by  $I_k = \{j \in Z: 2^{k-1} \leq j < 2^k\}$  if k > 0,  $I_0 = \{0\}$ and  $I_k = -I_{-k}$  if k < 0. These inequalities rest ultimately upon an extension of the Riesz-Thorin interpolation theorem and the following result of Paley and Hardy and Littlewood [3]: A complex number sequence  $\{\lambda_n\}$  has the property  $\sum_{n=0}^{\infty} \lambda_n a_n z^n \in H^2$  whenever  $\sum_{n=0}^{\infty} a_n z^n \in$  $H^1$  if and only if

(1.5) 
$$\sup_{k\geq 0} \sum_{n \in I_k} |\lambda_n|^2 < \infty$$

This suggests that the proper generalization of inequalities (1.3) and (1.4) is to the setting in which the unit circle T is replaced by a locally compact abelian (LCA) group G with ordered dual  $\Gamma$ -for the notion of lacunary decomposition extends naturally to ordered groups.

In this paper we carry out this extension and give some applications of these generalized inequalities. Specifically, in §3 we strengthen a theorem of Katznelson [6] and of Figà-Talamanca and Gaudry [2] on sets of uniqueness in  $L^{p}(G)$ . We also give stronger sufficient conditions than those found in Edwards [1] for the distribution fSto be zero where S is a pseudomeasure and f is an element of C(T)with absolutely convergent Fourier expansion.

I wish to acknowledge the work of J. H. Wells in extending Kellogg's result to the real line and to express my appreciation for his assistance in the preparation of this paper.

2. Main results. Let G be a compact connected abelian group with dual  $\Gamma$  ordered by a set P of positive elements. For  $1 \leq p \leq \infty$  define  $H^p(G)$  to be the set of all  $f \in L^p(G)$  such that  $\hat{f}(\gamma) = 0$ for  $\gamma \notin P$ . (Here and subsequently  $\hat{f}$  denotes the Fourier transform of f.)

DEFINITION. A lacunary decomposition of P is a countable collection  $\mathscr{D} = \{D_i\}_{i=1}^{\infty}$  of subsets of P satisfying: (1)  $D_i \cap D_j = \phi$  if  $i \neq j$ , and

(2) for each i there exists an  $lpha_i \in P$  such that

$$D_i = \{lpha: \ lpha_i < lpha \leqq 2 lpha_i\}$$
 .

Corresponding to such a decomposition  $\mathscr{D}$  we define the mixed norm space  $L^{r,s}_{\mathscr{P}}(P)$  to be the set of all  $f \in \mathscr{I}^{\infty}(P)$  with support contained in  $\bigcup_{i=1}^{\infty} D_i$  and  $||f||_{r,s,\mathscr{D}} < \infty$  where

Also we define  $L^{r,\infty}(P)$  to be the set of all  $f \in \mathscr{L}^{\infty}(P)$  such that  $||f||_{r,\infty} = \sup_{\alpha \in P} (\sum_{\beta \in I_{\alpha}} |f(\beta)|^{r})^{1/r} < \infty$  where  $1 \leq r < \infty$  and  $I_{\alpha} = \{\beta \in P: \alpha < \beta \leq 2\alpha\}$ . It is easy to verify that these spaces are Banach spaces and the Banach conjugate of  $L^{r,s}(P)$  is  $L^{r',s'}(P)$  if  $r, s < \infty$  where 1/r + 1/r' = 1/s + 1/s' = 1.

If r, s, u and v are real numbers in  $[1, \infty]$  then  $(L^{r,s}_{\mathcal{I}}(P), L^{u,v}_{\mathcal{I}}(P))$ , the multipliers from  $L^{r,s}_{\mathcal{I}}(P)$  into  $L^{u,v}_{\mathcal{I}}(P)$ , is the set of all  $\lambda \in \mathscr{I}^{\infty}(P)$ 

824

with the property that  $\lambda f \in L^{u,v}_{\mathscr{D}}(P)$  whenever  $f \in L^{r,s}_{\mathscr{D}}(P)$ . Each such  $\lambda$  determines a bounded linear operator from  $L^{r,s}_{\mathscr{D}}(P)$  into  $L^{u,v}_{\mathscr{D}}(P)$  whose norm will be called the multiplier norm of  $\lambda$ .

We record the following theorem from [7].

THEOREM 1. Let 1/p = 1/u - 1/r if r > u,  $p = \infty$  if  $r \leq u$  and let 1/q = 1/v - 1/s if s > v and  $q = \infty$  if  $s \leq v$ . Then  $(L^{r,s}_{\mathcal{J}}(P), L^{u,v}_{\mathcal{J}}(P)) = L^{p,q}_{\mathcal{D}}(P)$  and if  $\lambda \in L^{p,q}_{\mathcal{J}}(P)$ , its multiplier norm is  $||\lambda||_{p,q:\mathcal{D}}$ .

If A and B are subsets of  $L^{1}(G)$ , the multiplier space  $(\hat{A}, \hat{B})$  is the space of all measurable complex-valued functions  $\lambda$  on  $\Gamma$  such that for every  $f \in A$  there exists  $g \in B$  with  $\lambda \hat{f} = \hat{g}$ . Hedlund [5, p. 54] shows that  $L^{2p/2-p,\infty}(P) \subset (\hat{H}^{p}(G), \hat{H}^{2}(G))$  for  $1 \leq p \leq 2$ . This follows by showing that  $L^{2p/2-p,\infty}(P) \subset (\hat{H}^{p}(G), \hat{H}^{2}(G))$  for any lacunary decomposition  $\mathscr{D}$  and that for  $f \in H^{p}(G)$  and  $\lambda \in L^{2p/2-p,\infty}(P), ||\lambda \hat{f}||_{2} \leq K_{p}$  $||f||_{p} ||\lambda||_{2p/2-p,\infty:\mathscr{D}}$ . Here  $K_{p}$  is a constant independent of the decomposition  $\mathscr{D}$ .

THEOREM 2. If  $\mathscr{D}$  is a lacunary decomposition of P and  $f \in H^p(G), 1 \leq p \leq 2$ , then  $\hat{f} \in L^{p',2}(P)$  and there exists a constant  $A_p$  independent of  $\mathscr{D}$  and f so that  $||\hat{f}||_{p',2:\mathscr{D}} \leq A_p ||f||_p$ .

*Proof.* By Hedlund's result we have  $L^{2p/2-p,\infty}_{\mathscr{P}}(P) \subset (\hat{H}^p(G), \hat{H}^2(G))$  and

(2.1) 
$$||\lambda \hat{f}||_2 = ||\lambda \hat{f}||_{2,2;\mathscr{D}} \leq K_p ||f||_p ||\lambda||_{2p/2-p,\infty;\mathscr{D}}$$

for  $f \in H^{p}(G)$  and  $\lambda \in L^{2p/2-p,\infty}(P)$ , where  $K_{p}$  is a constant independent of  $\mathscr{D}$ . Then by Theorem 1

$$\hat{H}^p(G) \subset (L^{2p/2-p,\infty}_{\mathscr{D}}(P),\,\hat{H}^2(G)) = (L^{2p/2-p,\infty}_{\mathscr{D}}(P),\,L^{2,2}_{\mathscr{D}}(P)) = L^{p',2}_{\mathscr{D}}(P)\;,$$

and  $||\hat{f}||_{p',2:\mathscr{D}}$  is the smallest number which satisfies (2.1) for all  $\lambda \in L^{2p/2-p,\infty}_{\mathscr{D}}(P)$ . The theorem now follows with  $A_p = K_p$ .

The Riesz Projection Theorem allows us to extend this result to  $L^{p}(G)$ .

THEOREM 3. For  $1 there is a constant <math>A_p$  such that if  $\mathscr{D}$  is a lacunary decomposition of P and  $f \in L^p(G)$ , then

$$\left\{\sum_{k=-\infty}^{\infty}\left(\sum_{\gamma \in D_k}|\widehat{f}(\gamma)|^{p\prime}
ight)^{2/p\prime}
ight\}^{1/2} \leq A_p \parallel f \parallel_p$$
 ,

where  $D_k = -D_{-k}$  for  $k < 0, D_0 = \phi$ .

We now state the main result.

THEOREM 4. Let n be a positive integer and G a compact connected abelian group with ordered dual  $\Gamma$ . If  $f \in L^p(\mathbb{R}^n \oplus G)$ , 1 , then

$$\left\{\sum_{k=-\infty}^{\infty} \left(\int_{J_k\oplus_R} {}^{n-1} \sum_{\Gamma} |\widehat{f}(t,\gamma)|^{p'} dt
ight)^{2/p'}
ight\}^{1/2} \leq A_p \, ||f||_p \; ,$$

where  $A_p$  is a constant,  $J_k = [2^{k-1}, 2^k]$  for  $k > 0, J_0 = [-1, 1]$ , and  $J_k = -J_{-k}$  for k < 0.

Now by [8, p.40] any LCA connected group is of the form  $\mathbb{R}^n \oplus G$ where G is compact and connected and  $n \ge 0$ . Hence the combination of Theorems 3 and 4 yields the desired extension of Kellogg's result.

Proof of Theorem 4. We begin with a special version of Theorem 3. Consider the compact connected group  $T^* \oplus G$  which has dual  $Z^* \oplus \Gamma$ . We introduce an order by defining the following set  $\overline{P}$  of positive elements:

$$((k_1, \dots, k_n), \gamma) \in \overline{P}$$
 if either

(1)  $k_j > 0$  where  $1 \leq j \leq n$  and  $k_i = 0$  for i < j; or (2)  $k_j = 0$  for each j and  $\gamma \in P$ , the set of positive elements in  $\Gamma$ . Let  $\mathscr{D}$  be the lacunary decomposition determined by  $\{\alpha_i\}_{i=1}^{\infty}$  where  $\alpha_i = ((2^{i-1}, 0, \dots, 0), 0) \in Z^n \oplus \Gamma$ . By Theorem 3 there is a constant  $B_p$  such that if  $f \in L^p(T^n \oplus G)$  then

$$igg\{\sum_{i=-\infty}^\infty \Bigl(\sum_{D_i} |\widehat{f}\,|^{\,p\prime}\Bigr)^{\!\!2/p\prime}igg\}^{\!\!1/2} \leqq B_p\,||f\,||_p$$
 .

For  $m \in Z$ , let  $I_m$  be the lacunary block described in the introduction. If m > 1,  $I_m \bigoplus Z^{n-1} \bigoplus \Gamma \subset D_{m-1} \cup D_m$  and if m < -1,

$$I_{{\scriptscriptstyle m}} \oplus Z^{{\scriptscriptstyle n-1}} \oplus arGamma \subset D_{{\scriptscriptstyle m}} \,\cup\, D_{{\scriptscriptstyle m+1}}$$
 .

By the standard Hausdorff-Young Theorem

$$\left\{ \left(\sum_{{I_m} \oplus {Z^{n-1}} \oplus {I'}} |\widehat{f}|^{p'} 
ight)^{2/p'} 
ight\}^{1/2} \leqq ||f||_p, \,\, m=\, -\, 1,\, 0,\, 1.$$

Then by the triangular inequality

(2.2) 
$$\left\{\sum_{m=-\infty}^{\infty} \left(\sum_{I_m \oplus \mathbb{Z}^{n-1} \oplus \mathbb{F}} |\hat{f}|^{p\prime}\right)^{2/p\prime}\right\}^{1/2} \leq C_p ||!f||_p$$

for all  $f \in L^p(T^n \bigoplus G)$ , where  $C_p = 2B_p + 3$ .

Now let f be a continuous function on  $\mathbb{R}^n \oplus G$  which has compact support and choose a positive number L such that f(t, x) = 0 for  $(t, x) \in \mathbb{R}^n \bigoplus G$  and  $||t||_{\infty} > L$ . Choose positive integers m and  $\ell$  so that  $m < \ell$  and  $2^{m-\ell}L < \pi$ . Let  $h(s, x) = f(s_1 2^{\ell-m}, \dots, s_n 2^{\ell-m}, x)$  $(2^{\ell-m})^n (2\pi)^n, (s, x) \in T^n \bigoplus G$ , where we identify T with  $\mathbb{R}/2\pi$ . Then  $h \in L^p(T^n \bigoplus G)$  and for  $(k_1, \dots, k_n, \gamma) \in \mathbb{Z}^n \bigoplus \Gamma$ 

$$\begin{split} \hat{h}(k_{1}, \cdots, k_{n}, \gamma) &= \int_{-2^{m-\ell_{L}}}^{2^{m-\ell_{L}}} \cdots \int_{-2^{m-\ell_{L}}}^{2^{m-\ell_{L}}} \int_{G} e^{-ik_{1}s_{1}} \cdots e^{-ik_{n}s_{n}}(\gamma, -x) f(s2^{\ell-m}, x)(2^{\ell-m})^{n} dxds \\ &= \int_{-L}^{L} \cdots \int_{-L}^{L} \int_{G} e^{-ik_{1}2^{m-\ell_{1}}} \cdots e^{-ik_{n}2^{m-\ell_{1}}}(\gamma, -x) f(t, x) dxdt \\ &= \hat{f}(k_{1}2^{m-\ell}, \cdots, k_{n}2^{m-\ell}, \gamma) . \end{split}$$

From (2.2) we obtain

(2.3) 
$$\sum_{j=-\infty}^{\infty} \left( \sum_{I_j \oplus Z^{n-1} \oplus \Gamma} | \hat{h}(k,\gamma) |^{p'} \right)^{2/p'} \leq C_p^2 ||h||_p^2.$$

Since  $||h||_{p} = (2^{\ell-m})^{n/p'} (2\pi)^{n/p'} ||f||_{p}$  we may rewrite (2.3) in the form

$$\sum_{j=-\infty}^{\infty} \left( \sum_{I_j \oplus \mathbb{Z}^{n-1} \oplus \mathbb{F}} |\hat{f}(k_1 2^{m-\ell}, \cdots, k_n 2^{m-\ell}, \gamma)|^{p'} (2^{\ell-m})^{-n} \right)^{2/p'} \leq C_p^2 (2\pi)^{2n/p'} ||f||_p^2$$

so that, in particular,

$$(2.4) \quad \sum_{j=-\ell}^{\ell} \left( \sum_{I_j \oplus \mathbb{Z}^{n-1} \oplus \Gamma} |\hat{f}(k_1 2^{m-\ell}, \cdots, k_n 2^{m-\ell}, \gamma)|^{p'} (2^{m-\ell})^n \right)^{2/p'} \leq A_p^2 ||f||_p^2$$

where  $A_{p} = C_{p}(2\pi)^{n/p'}$ .

Now for  $r = 0, \dots, m - 1$  it is evident that certain partial sums of

$$\sum_{I_{\ell'-r}\oplus Z^{n-1}\oplus I'} |\hat{f}(k_1 2^{m-\ell'}, \cdots, k_n 2^{m-\ell'}, \gamma)|^{p'} (2^{m-\ell'})^n \\ = \sum_{k_1=2}^{2^{\ell'-r-1}} \sum_{k_2, \cdots, k_n \in Z} \sum_{\gamma \in I'} |\hat{f}(k_1 2^{m-\ell'}, \cdots, k_n 2^{m-\ell'}, \gamma)|^{p'} (2^{m-\ell'})^n$$

will converge to  $\int_{2^{m-r}}^{2^{m-r}} \int_{\mathbb{R}^{n-1}} \sum_{r} |\hat{f}(t,\gamma)|^{p'} dt$  as  $\checkmark$  tends to infinity. Similarly, for  $r = 0, -1, \dots, -m+1$ , the limit superior of the sum

$$\sum_{I_{-(\boldsymbol{c}'+r)}\oplus_{\boldsymbol{Z}}^{n-1}\oplus \boldsymbol{\Gamma}} |\hat{f}(k_1 2^{m-\boldsymbol{c}'}, \cdots, k_n 2^{m-\boldsymbol{c}'}, \boldsymbol{\gamma})|^{p'} (2^{m-\boldsymbol{c}'})^n$$

is not less than  $\int_{-2^{m+r-1}}^{-2^{m+r-1}} \int_{\mathbb{R}^{n-1}} \sum_{\Gamma} |\widehat{f}(t,\gamma)|^{p'} dt$ . Also since  $2/p' \leq 1$ ,

$$\sum_{j=-(\ell-m)}^{\ell-m} \left( \sum_{I_{j} \oplus_{Z}^{n-1} \oplus \Gamma} |\hat{f}(k_{1}2^{m-\ell}, \cdots, k_{n}2^{m-\ell}, \gamma)|^{p'} (2^{m-\ell})^{n} \right)^{2/p'} \\ \geq \left\{ \sum_{k_{1}=-2^{\ell-m+1}}^{2^{\ell-m-1}} \sum_{Z^{n-1} \oplus \Gamma} |\hat{f}(k_{1}2^{m-\ell}, \cdots, k_{n}2^{m-\ell}, \gamma)|^{p'} (2^{m-\ell})^{n} \right\}^{2/p'}$$

,

and the limit superior of the second sum dominates the integral

$$\left(\int_{-1}^{1}\int_{\mathbb{R}^{n-1}}\sum_{\Gamma}|\widehat{f}(t,\gamma)|^{p'} dt\right)^{2/p'}$$

Thus, by letting  $\checkmark$  tend to infinity in (2.4), we obtain

$$\left\{\sum_{k=-m}^{m} \left( \int_{J_k \oplus R^{n-1}} \sum_{\Gamma} |\widehat{f}(t,\gamma)|^{p'} dt 
ight)^{2/p'} 
ight\}^{1/2} \leq A_p ||f||_p,$$

and since m is arbitrary, our proof is complete for continuous functions with compact support. The general result now follows since such functions are dense in  $L^{p}(\mathbb{R}^{n} \oplus G)$ .

There is also a dual to this generalized Hausdorff-Young Theorem in case G is a compact group. Suppose  $\{\Gamma_k\}_{k=-\infty}^{\infty}$  is a collection of subsets of  $\Gamma$  for which a generalized Hausdorff-Young Theorem holds (e.g., Theorem 3). For p and q in  $[1, \infty)$  let  $L^{p,q}(\Gamma)$  be the space of all measurable complexvalued functions  $\lambda$  on  $\Gamma$  satisfying

$$||\lambda||_{p,q} = \Big\{\sum_{k=-\infty}^{\infty} \Big( \int_{\Gamma_k} |\lambda(\gamma)|^p d\gamma \Big)^{q/p} \Big\}^{1/q} < \infty$$
 .

THEOREM 5. If  $1 and <math>\lambda \in L^{p,2}(\Gamma)$  then there exists  $f \in L^{p'}(G)$ with  $\hat{f} = \lambda$  and  $||f||_{p'} \leq B_p ||\lambda||_{p,2}$ , where  $B_p$  is a constant depending only on p.

Proof. Since  $L^1(G) * L^p(G) = L^p(G)$  (\* denotes convolution) we have  $\hat{L}^1(G) \subset (\hat{L}^p(G), \hat{L}^p(G))$ . By assumption a generalized Hausdorff-Young Theorem holds in  $L^p(G)$ . Thus  $\hat{L}^p(G) \subset L^{p',2}(\Gamma)$  and the previous inclusion implies that  $\hat{L}^1(G) \subset (\hat{L}^p(G), L^{p',2}(\Gamma))$ . But  $(\hat{L}^p(G), L^{p',2}(\Gamma)) = (L^{p,2}(\Gamma), \hat{L}^{p'}(G))$  by duality; hence  $\hat{L}^1(G) \subset (L^{p,2}(\Gamma), \hat{L}^{p'}(G))$ or, equivalently,  $L^{p,2}(\Gamma) \subset (\hat{L}^1(G), \hat{L}^{p'}(G))$ . However  $(\hat{L}^1(G), \hat{L}^{p'}(G))$  is known to be  $\hat{L}^{p'}(G)$  [1, p.255] so that  $L^{p,2}(\Gamma) \subset \hat{L}^{p'}(G)$ . An application of the closed graph theorem establishes existence of the constant  $B_p$ .

3. Applications. Let A denote the space of all functions f continuous on T which have absolutely convergent trigonometric expansions and norm given by

$$||f||_{\scriptscriptstyle A} = \sum_{n=-\infty}^{\infty} |\widehat{f}(n)|$$
.

With multiplication defined as the pointwise product of functions, A becomes a Banach algebra. Each element of the dual of A may be identified with a pseudomeasure S, that is, a distribution on  $C^{\infty}(T)$ whose sequence of Fourier coefficients  $\hat{S}(n) = S(e^{-in\theta})$   $(n \in Z)$  belong to  $\swarrow^{\infty}$ . Given a pseudomeasure S and an f in A the distribution fS is defined by fS(u) = S(fu) for  $u \in C^{\infty}(T)$ . In [1] Edwards shows that fS is the zero distribution when f and S satisfy the following conditions:

- (a)  $S \in \mathbb{Z}^q, 1 \leq q \leq \infty$ ;
- (b) f vanishes on supp (S) = E; and

(c) there exists a sequence of positive numbers  $\varepsilon_j$  tending to zero such that f(x) = 0  $(\varepsilon_j^{1/2-1/q})$  for all x such that dist  $(x, E) \leq \varepsilon_j$ .

THEOREM 6. Suppose S is a pseudomeasure and  $f \in A$ . If conditions (b) and (c) above hold and either

$$(a_1) \ \widehat{S} \in \mathscr{C}^{q,2} \text{ if } 1 < q \leq 2$$

or

$$(a_2) \ \widehat{S} \in {\sc \prime}^{q,\infty} \ \text{if} \ 2 < q < \infty$$

then fS = 0.

*Proof.* If  $1 < q \leq 2$  it follows from Theorem 5 that  $\hat{S} = \hat{g}$  for some  $g \in L^{q'}(T)$ . Therefore  $S(u) = \int_{-\pi}^{\pi} u(t)g(t) \quad dt/2\pi, u \in C^{\infty}(T)$ , and  $\operatorname{supp}(S) = \operatorname{supp}(g)$ . But  $fS(u) = S(fu) = \int_{-\pi}^{\pi} f(t)u(t)g(t) \quad dt/2\pi, u \in C^{\infty}(T)$ , and  $\operatorname{supp}(f) \cap \operatorname{supp}(g) = \phi$ ; hence fS = 0.

Now  $\swarrow^q \subsetneq \swarrow^{q,2}$  when q < 2, so we have a stronger result in this case.

For the case q > 2, we need only apply our generalized Hausdorff-Young theorem in 13.5.5 and the appropriate version of Hölders inequality in 13.5.9 of the argument in Edwards [1, pp.101-102] to obtain fS = 0 when  $\hat{S} \in \ell^{q,\infty}$  and f satisfies (b) and (c).

The condition  $\hat{S} \in \mathbb{Z}^{q,\infty}$  when  $2 < q < \infty$  is a significant weakening of condition (a) since it does not require that  $\lim_{|n|\to\infty} |\hat{S}(n)| = 0$ .

Our second application concerns sets of uniqueness for  $L^{p}(G)$ ,  $1 . In a recent paper which generalized earlier work of Katznelson [6, p.101-103], Figa-Talamanca and Gaudry proved the existence of sets <math>E \subset G$  of positive Haar measure such that if  $f \in L^{1}(G)$ , supp  $(f) \subset E$  and  $\hat{f} \in L^{p}(G)$  for some  $p, 1 \leq p < 2$ , then f = 0. Here G is a nondiscrete LCA group with dual  $\Gamma$ .

For our generalization we shall only need to assume that there is a decomposition  $\{\Gamma_k\}_{k=-\infty}^{\infty}$  of  $\Gamma$  for which a generalized Hausdorff-Young Theorem holds and that when f is measurable on  $\Gamma$  and 1 then

$$\sum_{k=-\infty}^{\infty} \Bigl( \int_{{arGamma}_k} f(\gamma) \, |^p d\gamma \Bigr)^{\! 2/p} < \infty \, .$$

always implies

#### LYNN R. WILLIAMS

$$\sum_{k=-\infty}^{\infty} \left( \int_{\Gamma_k} |f(eta-\gamma)|^p d\gamma 
ight)^{2/p} < \infty ext{ for each } eta \in \Gamma ext{ .}$$

For example, these conditions hold relative to the decompositions in the statement and proof of Theorem 4. Subject to this restriction on the regularity of the lacunary decomposition  $\{\Gamma_k\}_{k=-\infty}^{\infty}$  we have the following result.

THEOREM 7. There is a subset E of G of positive Haar measure such that if  $f \in L^1(G)$ , supp  $(f) \subset E$  and

$$\|\widehat{f}\|_{p,2} = \left\{\sum_{k=-\infty}^{\infty} \left( \int_{arGamma_k} |\widehat{f}(\gamma)|^p d\gamma 
ight)^{2/p} 
ight\}^{1/2} < \infty$$

for some p, 1 , then <math>f = 0.

Essentially the proof consists of establishing the following lemma.

LEMMA. If  $1 \varepsilon > 0$ , and M is a subset of G of measure 1, then there exists a subset  $E_{\varepsilon,p}$  of M of measure greater than or equal  $1 - \varepsilon$  and a function  $\Phi \in L^1(G) \cap L^{\infty}(G)$  such that  $\Phi = 1$ on  $E_{\varepsilon,p}$  and  $||\hat{\Phi}||_{p',2} < \varepsilon$ .

Proof of Theorem 7. Let  $\varepsilon_n = 1/4^n$ ,  $p_n = 2 - \varepsilon_n$  and choose M a subset of G of Haar measure 1. Let  $E = \bigcap_{n=1}^{\infty} E_{\varepsilon_n, p_n}$ , where  $E_{\varepsilon_n, p_n}$  and  $\Phi_n$  are as in the lemma.

Suppose  $f \in L^1(G)$ , supp  $f \subset E$ , and  $||\hat{f}||_{p,2} < \infty$  for some p,  $1 . Since <math>||\hat{f}||_{\infty} < \infty$  and p/2 < 1 we have  $||\hat{f}||_{2,2} < \infty$ . Let  $\gamma \in \Gamma$ . It follows from our assumption on the regularity of  $\{\Gamma_k\}$  that  $||\hat{f}_{\tau}||_{p,2} < \infty$  and  $||\hat{f}_{\tau}||_{2,2} < \infty$  where  $\hat{f}_{\tau}(\beta) = \hat{f}(\gamma - \beta)$ . Choose N so that n > N implies  $p_n > p$ . By the interpolation theorem [4, p. 1069] there is a constant  $K_{\tau}$  such that  $||\hat{f}_{\tau}||_{p_n,2} \leq K_{\tau}$  for n > N. By Parseval's identity and Hölder's inequality we have

$$\begin{split} |\hat{f}(\gamma)| &= \left| \int_{\mathcal{G}} (\gamma, -x) f(x) \Phi_n(x) dx \right| \\ &= \left| \int_{\Gamma} \hat{f}(\gamma - \beta) \Phi_n(\beta) d\beta \right| \leq ||\hat{f}_{\gamma}||_{p_n, 2} ||\hat{\Phi}_n||_{p'_n, 2} \leq \varepsilon_n K_{\gamma} \end{split}$$

Thus  $\hat{f}(\gamma) = 0$  for  $\gamma \in \Gamma$  and hence f = 0.

Proof of the Lemma. Since our proof closely parallels that in [2] we omit similar details and computations.

Define a sequence  $\{\pi_n\}_{n=0}^{\infty}$  of partitions of M so that  $\pi_0 = \{M\}$ 

830

and  $\pi_n$  is obtained from  $\pi_{n-1}$  by dividing each set of  $\pi_{n-1}$  into two sets of equal measure. Define a sequence  $\{r_n\}_{n=0}^{\infty}$  of Rademacher functions on M with respect to these partitions. (i.e.,  $r_0 = \chi_M$ ,  $r_n$  is constant on each set of  $\pi_n$  with the value  $\pm 1$  and  $\int_A r_n(x) dx = 0$  for each  $A \in \pi_{n-1}$ , n > 0.) Forming all possible finite products of the  $r_n$ yields an orthogonal system  $W = \{w_i\}$  of Walsh functions whose linear span contains all functions supported on M and constant on the sets of  $\pi_n$  for some  $n \geq 0$ .

Since 1 we can choose t, <math>0 < t < 1, so that 1/p = 1 - t/2. Now let N denote a positive integer such that

where  $A_p$  is the constant in the generalized Hausdorff-Young Theorem for G. We will show the existence of disjoint compact sets  $K_1, \dots, K_n$ in  $\Gamma$  and orthogonal linear combinations of Walsh functions  $\phi_1, \dots, \phi_n$ with supp  $(\phi_j) \subset M, 1 \leq j \leq N$ , satisfying the following conditions:

$$(3.2) \qquad \int_{\mathcal{G}} |\phi_j(x)| \ dx \leq 2 \ \text{and} \ \int_{\mathcal{G}} |\phi_j(x)|^2 dx \leq 2^k + 1, 1 \leq j \leq N ,$$

where  $1/2^k < arepsilon/N \leq 1/2^{k-1}$  ;

(3.3) the measure of  $\{x \in M: \phi_j(x) = 1\}$  is greater than or equal  $1 - \varepsilon/N, \ 1 \leq j \leq N$ ; and

(3.4) 
$$\left\{\sum_{k=-\infty}^{\infty} \left( \int_{\Gamma_k \cap K_n} \left| \sum_{j=1}^N \hat{\phi}_j(\beta) \right|^{p'} d\beta \right)^{2/p'} \right\}^{1/2} \leq 4 \| \hat{\phi}_n \|_{p',2}$$

and

$$\left\{\sum_{k=-\infty}^{\infty} \left(\int_{\Gamma_k\setminus K_n} |\hat{\phi}_n(eta)|^{p'} \, deta
ight)^{2/p'}
ight\}^{1/2} \leq arepsilon/2N, \ 1\leq n\leq N$$
 .

Denote the sums in (3.4) by  $||\sum_{j=1}^{N} \hat{\phi}_j||_{p',2;K_n}$  and  $||\hat{\phi}_n||_{p',2;\Gamma\setminus K_n}$ , respectively.

Let  $\phi_1 = r_0$  and choose a compact set  $K_1$  so that

$$\|\widehat{\phi}_{_1}\|_{{}_{p'},2;arGamma \setminus {K_1}} < arepsilon/2N$$
 .

Assume we have n functions and compact sets satisfying the above with (3.4) replaced by

(3.5) 
$$\left\| \sum_{j=1}^{n} \hat{\phi}_{j} \right\|_{p',2;K_{m}} \leq \left(2 + \frac{2n}{N}\right) \|\hat{\phi}_{m}\|_{p',2}$$

and

$$\| \hat{\phi}_{\scriptscriptstyle m} \|_{\scriptscriptstyle p',2; \varGamma \setminus K_m} \leq arepsilon / 2N, \; 1 \leq m \leq n$$
 .

By use of Bessel's inequality and Dini's Theorem we obtain the

existence of a finite set  $F \subset W$  such that

$$\sum_{w_i 
otin F} |\left( \gamma, \, w_i 
ight)|^2 < \Bigl( arepsilon / 2N \sum_{j=1}^n |\, K_j \,| \Bigr)^2 (1/(2^k \,+\, 1)) \; .$$

uniformly for  $\gamma$  in the compact set  $K_1 \cup K_2 \cup \cdots \cup K_n$ , where  $|K_j|$  denotes the measure of  $K_j$ . We have assumed  $|K_j| \neq 0, 1 \leq j \leq n$ , for otherwise  $\phi_j$  is the desired function with  $E_{\varepsilon,p} = \{x: \phi_j(x) = 1\}$ . Suppose further that F is chosen so as to contain all Walsh functions appearing in the expansions of  $\phi_1, \cdots, \phi_n$ . Let m be a positive integer so that the elements of F (and hence  $\phi_1, \cdots, \phi_n$ ) are constant on the sets  $E_1, \cdots, E_{2^m}$  of  $\pi_m$ . For each  $j, 1 \leq j \leq 2^m$ , consider the partition of  $E_j$  into  $2^k$  subsets  $E'_{1j}, \cdots, E'_{2^{k_j}}$  determined by  $\pi_{m+k}$ . Define  $\phi_{n+1}$  to be zero off M and on M, to be as follows

Then for  $1 \leq j \leq 2^m$ ,  $\int_{E_j} \phi_{n+1}(x) dx = 0$  and therefore $\int_M \phi_{n+1}(x) w(x) dx = 0$ 

for each  $w \in F$ . Thus  $\phi_{n+1} = \sum_{w_j \notin F} a_j w_j$  and  $\phi_{n+1}$  is easily seen to satisfy (3.2) and (3.3). For  $\gamma \in \bigcup_{j=1}^n K_j$ ,

$$egin{aligned} &|\, \hat{\phi}_{n+1}(\gamma)\,| &\leq \sum\limits_{w_j 
otin F} \,|\, a_j \int \!\!\! w_j(x) \gamma(\,-\,x) dx\,| \,\leq \sum\limits_{w_j 
otin F} \,|\, a_j\,| \,\, |\, (w_j,\,\gamma)\,|^2 \ &\leq arepsilon_j 2N \sum\limits_{j=1}^n \,|\, K_j\,| \,\,. \end{aligned}$$

If  $1 \leq m_{\scriptscriptstyle 0} \leq n$  and  $||\hat{\phi}_{m_{\scriptscriptstyle 0}}||_{p',z} < \varepsilon/2$  we are done; otherwise

$$arepsilon/N \sum_{j=1}^n |K_j| \leq 2 \, || \hat{\phi}_{m_0} ||_{p',2} / N \, \sum_{j=1}^n |K_j|$$

and therefore

$$|\hat{\phi}_{n+1}||_{p',2:K_{m_0}} \leq |K_{m_{\mathfrak{I}}}| \ ||\hat{\phi}_{n+1}||_{\infty:K_{m_0}} \leq 2 \ ||\hat{\phi}_{m_0}||_{p',2} / N$$
 .

Hence

$$\left\|\sum_{j=1}^{n+1} \hat{\phi}_j\right\|_{p',2:K_{m_0}} \leq \left(2 + \frac{2(n+1)}{N}\right) || \hat{\phi}_{m_0} ||_{p',2}$$

by the triangular inequality. Also since  $||\hat{\phi}_{n+1}||_{p',2:\bigcup_{j=1}^{n}K_j} < \varepsilon/2N$ , there exists a compact set  $K_{n+1}$  disjoint from  $\bigcup_{j=1}^{n}K_j$  such that

$$|| \widehat{\phi}_{\scriptscriptstyle n+1} ||_{\scriptscriptstyle p',2; \varGamma \setminus K_{n+1}} < arepsilon/2N$$
 .

Then

$$\begin{split} \left\|\sum_{j=1}^{n+1} \hat{\phi}_{j}\right\|_{p',2;K_{n+1}} &\leq ||\hat{\phi}_{n+1}||_{p',2} + \sum_{j=1}^{n} ||\hat{\phi}_{j}||_{p',2;I \setminus K_{j}} \\ &\leq ||\hat{\phi}_{n+1}||_{p',2} + \varepsilon/2 \leq \left(2 + \frac{2(n+1)}{N}\right) ||\hat{\phi}_{n+1}||_{p',2} , \end{split}$$

where again we have assumed that  $||\hat{\phi}_{n+1}||_{p',2;K_{n+1}} \ge \varepsilon/2$ . The existence of the functions now follows by induction.

Let  $\Phi = (1/N) \sum_{j=1}^{N} \phi_j$  and  $E_{\varepsilon,p} = \{x \in M: \Phi(x) = 1\}$ . Clearly  $|E_{\varepsilon,p}| \ge 1 - \varepsilon$  and

$$||\hat{\varphi}||_{p',2} \leq ||\hat{\varphi}||_{p',2; \bigcup_{j=1}^{N} K_{j}} + ||\hat{\varphi}||_{p',2; \Gamma \setminus \bigcup_{j=1}^{N} K_{j}} \leq ||\hat{\varphi}||_{p',2; \bigcup_{j=1}^{N} K_{j}} + \varepsilon/2.$$

In order to complete the proof it suffices to show that

$$\|\widehat{\varPhi}\|_{p',2:\mathop{\cup}\limits_{\substack{j=1K_j}}^N} .$$

But

$$\begin{split} ||\hat{\varPhi}||_{p',2:\sum_{m=1}^{N}K_{m}}^{2} &= \sum_{\ell=-\infty}^{\infty} \left( \int_{\Gamma_{\ell} \cap (\sum_{m=1}^{N}K_{m})} \left| \frac{1}{N} \sum_{j=1}^{N} \hat{\phi}_{j}(\gamma) \right|^{p'} d\gamma \right)^{2/p'} \\ &= \frac{1}{N^{2}} \sum_{\ell=-\infty}^{\infty} \left( \sum_{m=1}^{N} \int_{\Gamma_{\ell} \cap K_{m}} \left| \sum_{j=1}^{N} \hat{\phi}_{j}(\gamma) \right|^{p'} d\gamma \right)^{2/p'} \\ &\leq \frac{1}{N^{2}} \sum_{m=1}^{N} \left\| \left| \sum_{j=1}^{N} \hat{\phi}_{j} \right\|_{p',2;K_{m}}^{2} \leq \frac{16}{N^{2}} \sum_{m=1}^{N} ||\hat{\phi}_{m}||_{p',2}^{2} \\ &\leq \frac{16A_{p}^{2}}{N^{2}} \sum_{m=1}^{N} ||\phi_{m}||_{p}^{2} \leq \frac{16A_{p}^{2}}{N^{2}} \sum_{m=1}^{N} (||\phi_{m}||^{1-t} ||\phi_{m}||_{2}^{t})^{2} \\ &\leq \frac{16A_{p}^{2}}{N} 2^{2(1-t)} (2^{k}+1)^{t} \leq [32A_{p}\varepsilon^{-t/2} N^{t/2-1/2}]^{2} < \left(\frac{\varepsilon}{2}\right)^{2}. \end{split}$$

#### References

1. R. E. Edwards, Fourier Series: A Modern Introduction, Vol. II., Holt, Rinehart and Winston, New York, 1967.

2. A. Figà-Talamanca and G. I. Gaudry, Multipliers and sets of uniqueness of  $L^p$ , Mich. J. Math., 17 (1970), 179-191.

3. G. H. Hardy and J. E. Littlewood, Notes on the theory of series (XX): Generalizations of a theorem of Paley, Quart. J. Math. Oxford Ser., 8 (1937), 161-171.

4. J. H. Hedlund, Multipliers of H<sup>p</sup> spaces, J. Math. Mech., 18 (1969), 1067-1074.

5. ——, Multipliers of H<sup>p</sup> spaces, Ph. D. Thesis, Univ. of Mich., 1968.

6. Y. Katznelson, An Introduction to Harmonic Analysis, Wiley, New York, 1968.

7. C. N. Kellogg, An extension of the Hausdorff-Young Theorem, (to appear).

8. W. Rudin, Fourier Analysis on Groups, Wiley (Interscience), New York, 1962.

Received October 6, 1970. Submitted as partial fulfillment of the requirements for the degree of Doctor of Philosophy from the University of Kentucky.

UNIVERSITY OF KENTUCKY

## PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON Stanford University Stanford, California 94805

C. R. HOBBY

University of Washington Seattle, Washington 98105 J. DUGUNDJI Department of Mathematics University of Southern California Los Angeles, California 90007

RICHARD ARENS

University of California Los Angeles, California 90024

### ASSOCIATE EDITORS

E. F. BECKENBACH B. H. NEUMANN F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA CALIFORNIA INSTITUTE OF TECHNOLOGY UNIVERSITY OF CALIFORNIA MONTANA STATE UNIVERSITY UNIVERSITY OF NEVADA NEW MEXICO STATE UNIVERSITY OREGON STATE UNIVERSITY UNIVERSITY OF OREGON OSAKA UNIVERSITY UNIVERSITY OF SOUTHERN CALIFORNIA STANFORD UNIVERSITY UNIVERSITY OF TOKYO UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF WASHINGTON

AMERICAN MATHEMATICAL SOCIETY CHEVRON RESEARCH CORPORATION NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. The editorial "we" must not be used in the synopsis, and items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The Pacific Journal of Mathematics is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

# Pacific Journal of Mathematics Vol. 38, No. 3 May, 1971

J. T. Borrego, Haskell Cohen and Esmond Ernest Devun, <i>Uniquely</i>	5 ( F
Clan Eugana Bradan. Some examples for the fixed point property.	571
William Lee Durpum Changetenizations of uniform convenity	571
Douglos Dorry. The convex hulls of the vertices of a polycon of order n	502
Educing Dude and Look Women Smith Reflexing onen mannings	507
Edwin Duda and Jack Warren Smith, <i>Reflexive open mappings</i>	597
Y. K. Feng and M. V. Subba Rao, On the density of $(k, r)$ integers	613
Irving Leonard Glicksberg and Ingemar Wik, Multipliers of quotients of	610
L1	019
continua	625
Lawrence Albert Harris, A continuous form of Schwarz's lemma in normed	
linear spaces	635
Richard Earl Hodel, <i>Moore spaces and</i> $w \Delta$ <i>-spaces</i>	641
Lawrence Stanislaus Husch, Jr., Homotopy groups of PL-embedding spaces.	
<i>II</i>	653
Yoshinori Isomichi, New concepts in the theory of topological	
space—supercondensed set, subcondensed set, and condensed set	657
J. E. Kerlin, On algebra actions on a group algebra	669
Keizō Kikuchi, <i>Canonical domains and their geometry in</i> C <sup>n</sup>	681
Ralph David McWilliams, On iterated w*-sequential closure of cones	697
C. Robert Miers, <i>Lie homomorphisms of operator algebras</i>	717
Louise Elizabeth Moser, <i>Elementary surgery along a torus knot</i>	737
Hiroshi Onose, Oscillatory properties of solutions of even order differential	
equations	747
Wellington Ham Ow, <i>Wiener's compactification and</i> $\Phi$ -bounded harmonic	
functions in the classification of harmonic spaces	759
Zalman Rubinstein, On the multivalence of a class of meromorphic	
functions	771
Hans H. Storrer, Rational extensions of modules	785
Albert Robert Stralka, The congruence extension property for compact	
topological lattices	795
Robert Evert Stong, On the cobordism of pairs	803
Albert Leon Whiteman, An infinite family of skew Hadamard matrices	817
Lynn Roy Williams, Generalized Hausdorff-Young inequalities and mixed	
norm spaces	823