

# Pacific Journal of Mathematics

**FINITE DIMENSIONAL TORSION FREE RINGS**

THOMAS J. CHEATHAM

## FINITE DIMENSIONAL TORSION FREE RINGS

THOMAS J. CHEATHAM

**In the class of rings with zero singular left ideal, several characterizations of rings with finite left Goldie dimension are given. They include: the direct limit of torsion free modules is torsion free; the direct limit of torsion free injective modules is injective; each absolutely pure torsion free module is injective; each module has a unique (up to isomorphism) torsion free covering module. The latter result gives a converse, in a special case, to a theorem of Mark Teply.**

Throughout,  $R$  will denote an associative ring with identity and module, without further qualification, will mean unitary left  $R$ -module. For a module  $E$ , we use  $S \subseteq' E$  to denote that  $S$  is a large submodule of  $E$ [4, p. 60];  $Z(E)$  will denote the singular submodule of  $E$ , which consists of those elements in  $E$  whose annihilators are large left ideals in  $R$ .

DEFINITION 1. A module  $E$  is torsion free if  $Z(E) = (0)$  and if  $Z(R) = (0)$  we say  $R$  is a torsion free ring.

A submodule  $S$  of a module  $E$  is closed in  $E$  if  $S \subseteq' T \subseteq E$  implies  $T = S$ . The following facts are easily verified.

LEMMA 1. *Let  $S$  be a submodule of a module  $E$ .*

(a) *If  $Z(E/S) = (0)$ ,  $S$  is closed in  $E$ .*

(b) *If  $Z(E) = (0)$ ,  $S$  is closed in  $E$  if and only if  $Z(E/S) = (0)$ .*

*Proof.* See Lemma 2.3 in [8].

DEFINITION 2. A module  $E$  has finite (Goldie) dimension if it contains no infinite direct sum of nonzero submodules. If the module  $R$  has finite dimension we call  $R$  a finite dimensional ring and write  $\dim R$  is finite.

1. **Torsion Free Rings.** Over an integral domain the direct limit of torsion free modules is torsion free. In this section we show that, in the class of torsion free rings, this property characterizes the finite dimensional rings. We also give two noetherian-like characterizations of such rings.

We record a theorem of F. Sandomierski [7] for easy reference.

THEOREM S. *Let  $Z(R) = (0)$ , and  $Q$  the maximal left quotient*

ring of  $R$  [4, p. 106], then the following are equivalent.

- (a)  $QI = Q$  for every  $I \subseteq R$ .
- (b) For  $I \subseteq R$  there are  $a_1, a_2, \dots, a_n \in I$  such that  $\sum Ra_i \subseteq R$ .
- (c) For  $I \subseteq R$  there are  $a_1, a_2, \dots, a_n \in I$  such that  $\sum Ra_i \subseteq I$ .
- (d)  $\dim R$  is finite.
- (e)  $Q$  is a semisimple (artinian) ring.
- (f)  $\text{Ker}(R \otimes_R E \rightarrow Q \otimes_R E) = Z(E)$  for every module  $E$ .

Sandomierski [7, Th. 2.5, p. 118] noted that if  $R$  has finite dimension the direct sum of torsion free injective modules is injective. If  $Z(R) = (0)$  the converse is also known. In fact, under this assumption, it follows from Teply [10, Th. 2.1, p. 451] that  $\dim R$  is finite if and only if any countable direct sum of torsion free injective modules is injective.

A set  $\mathcal{S}$  of submodules of a module is directed if given  $X, Y \in \mathcal{S}$  there is a  $Z \in \mathcal{S}$  such that  $X \cup Y \subseteq Z$ . Clearly the union of a directed set of submodules is a submodule.

We will make use of the following lemma which is an unpublished remark of M. Teply.

LEMMA 2. *Let  $Z(R) = (0)$ . The union of a directed set of closed submodules of a torsion free module is a closed submodule if and only if  $\dim R$  is finite.*

*Proof.* Assume that  $\dim R$  is finite then, to show that the condition is necessary, we proceed as in [9, Prop. 2.1 (3)] using Theorem  $S(b)$ .

Conversely, let  $E_1, E_2, \dots$  be a countable family of torsion free injective modules. Then  $E = \bigoplus_{n=1}^{\infty} E_n$  is a torsion free module so has a torsion free injective hull  $I(E)$ . But  $E$  can be written as the union of the chain  $S_1 \subseteq S_2 \subseteq \dots$  of injective (hence closed) submodules of  $I(E)$ , where  $S_n = \bigoplus_{i=1}^n E_i$ . Hence  $E$  is closed in  $I(E)$ . But  $E \subseteq I(E)$  so  $E = I(E)$ , i.e.,  $E$  is injective. By a remark above  $\dim R$  is finite.

The following useful result is well-known and trivial to prove.

LEMMA 3. *Let  $f, g \in \text{Hom}_R(E, F)$  where  $Z(F) = (0)$ .*

- (1) *If  $f$  and  $g$  agree on a large submodule of  $E$  then  $f = g$ .*
- (2) *If  $E$  is injective,  $f(E)$  is a direct summand of  $F$ .*

THEOREM 1. *Let  $Z(R) = (0)$ . Then the following statements are equivalent.*

- (1)  $\dim R$  is finite.
- (2) *The direct limit of torsion free modules is torsion free.*

(3) *The direct limit of torsion free injective modules is injective.*

*Proof.* (1) implies (2). Let  $\{E, f_a\}$  be the direct limit of the directed system of torsion free modules  $\{E_a; f_a^b, A\}$ . Then  $E = \bigcup_{a \in A} \text{Im } f_a$ , so to show  $E$  is torsion free it suffices to show  $\text{Im } f_a$  is torsion free for each  $a \in A$ . But  $\text{Im } f_a \cong E_a / \text{Ker } f_a$ , and  $\text{Ker } f_a = \bigcup_{b \geq a} \text{Ker } f_a^b$ . Furthermore, for each  $a \in A$ ,  $\{\text{Ker } f_a^b \mid b \geq a, b \in A\}$  is a directed set of submodules of the torsion free module  $E_a$ , and since  $\text{Im } f_a^b \subseteq E_b$ ,  $\text{Ker } f_a^b$  is closed in  $E_a$ . By Lemma 2  $\text{Ker } f_a$  is closed in  $E_a$  and hence by Lemma 1(b)  $\text{Im } f_a$  is torsion free. Hence (1) implies (2).

(2) implies (1). Let  $\{C_a; a \in A\}$  be a directed set of closed submodules of a torsion free module  $E$ . For  $a, b \in A$  such that  $C_a \subseteq C_b$  define a function  $f_a^b: E/C_a \rightarrow E/C_b$  by  $f_a^b(x + C_a) = x + C_b$ . Clearly  $f_a^b \in \text{Hom}_R(E/C_a, E/C_b)$  and one easily checks that  $\{E/C_a; f_a^b, A\}$  is a directed system of torsion free modules with direct limit  $E/\bigcup_{a \in A} C_a$ . So from (2)  $E/\bigcup_{a \in A} C_a$  is torsion free and hence (Lemma 1(a))  $\bigcup_{a \in A} C_a$  is closed in  $E$ . So (1) follows from Lemma 2.

(1) implies (3). Let  $\{E, f_a\}$  be the direct limit of a directed system of torsion free injective modules  $\{E_a; f_a^b, A\}$ . Let  $I$  be any left ideal of  $R$  and  $h \in \text{Hom}_R(I, E)$ . By Theorem S(c) there is a finitely generated left ideal  $J$  of  $R$  such that  $J \subseteq I$ . From condition (2) we know that  $E$  is torsion free so by Lemma 3(1) we see that an extension of  $f' = f|_J: J \rightarrow E$  to all of  $R$  will give the desired extension of  $f$ . To see that  $f'$  can be extended to  $R$  we proceed as follows.

Let  $p: F \rightarrow J$  be an  $R$ -homomorphism of a finitely generated free module  $F$  onto  $J$  and identify  $J$  with  $F/\text{Ker } p$ . It follows easily from Theorem S that  $F$  has finite dimension and that any submodule of  $F$  has a finitely generated large submodule. Thus let  $K$  be a finitely generated large submodule of  $\text{Ker } p$ .

Now  $f' \circ p(F)$  is a finitely generated submodule of  $E = \bigcup_{a \in A} \text{Im } f_a$  so there's an  $a \in A$  such that  $f' \circ p(F) \subseteq \text{Im } f_a$ . Hence, by the projectivity of  $F$ , there's an  $h \in \text{Hom}_R(F, E_a)$  such that  $f_a \circ h = f' \circ p$ . Then  $h(\text{Ker } f' \circ p) \subseteq \text{Ker } f_a = \bigcup_{b \geq a} \text{Ker } f_a^b$  and since  $K$  is finitely generated  $h(K) \subseteq \text{Ker } f_a^b$  for some  $b \geq a$ , i.e.,  $f_a^b \circ h(K) = (0)$ . Therefore  $f_a^b \circ h$  induces an  $R$ -homomorphism  $g$  from  $F/\text{Ker } p = J$  into  $E_b$ . But  $E_b$  is injective so  $g$  can be extended to  $g^* \in \text{Hom}_R(R, E_b)$ . Define  $f^*$  from  $R$  to  $E$  by  $f^* = f_b \circ g^*$ . Then  $f^*$  is an extension of  $f'$  so the desired extension of  $f$ .

(3) implies (1). Clearly (3) implies that any direct sum of torsion free injective modules is injective. So (1) follows by a previous remark. This completes the proof.

As a corollary we give an easy proof of (d) implies (e) in Theorem S.

**COROLLARY.** *A finite dimensional torsion free ring  $R$  has a semisimple (artinian) maximal left quotient ring  $Q$ .*

*Proof.* It suffices to show that every left ideal of  $Q$  is a direct summand of  $Q$  (as a  $Q$ -module). Since  $Q$  is von Neumann regular any finitely generated left ideal of  $Q$  is a direct summand. Hence any such ideal is left  $R$ -injective as  $Q$  is. But  $Z({}_R Q) = (0)$  and it follows, via a direct limit argument and Theorem 1(3), that any left ideal of  $Q$  is  $R$ -injective, hence  $Q$ -injective. Thus any left ideal of  $Q$  is a direct summand.

B. Maddox [5] calls a module  $M$  absolutely pure if for every module  $E$  containing  $M$  as a submodule the sequence  $0 \rightarrow G \otimes M \rightarrow G \otimes E$  is exact for every right  $R$ -module  $G$ . He showed that any direct sum of absolutely pure modules is absolutely pure. C. Megibben [6, Th. 3, p. 564] characterized left noetherian rings by the property "each absolutely pure module is injective." This result was also obtained indepently by Edgar Enochs.

We have a corresponding characterization of finite dimensionality in the class of torsion free rings.

**THEOREM 2.** *Let  $Z(R) = (0)$ . Then  $\dim R$  is finite if and only if each absolutely pure torsion free module is injective.*

*Proof.* Assume that  $\dim R$  is finite, and let  $E$  be an absolutely pure torsion free module and  $f$  an  $R$ -homomorphism of a left ideal  $I$  of  $R$  into  $E$ . Let  $J$  be a finitely generated left ideal of  $R$  such that  $J \subseteq I$ . As remarked above it suffices to extend  $f'$ , the restriction of  $f$  to  $J$ , to all of  $R$ . But this can be done by [6, Cor. 2, p. 562].

Conversely, it suffices to show that the direct sum of torsion free injective modules is injective. But any such sum is torsion free and absolutely pure, hence injective.

**2. Torsion free covers.** The main result of this section (Theorem 3) gives a converse to [9, Th. 2.4, p. 459], in a special case. We begin with a definition.

**DEFINITION 3.** A torsion free cover of a module  $E$  is a homomorphism  $g$  from a torsion free module  $T(E)$  onto  $E$  such that:

- (1)  $\text{Ker } g$  contains no nonzero closed submodule of  $T(E)$ ,

(2) given  $f \in \text{Hom}_R(F, E)$  where  $F$  is torsion free there is an  $h \in \text{Hom}_R(E, T(E))$  such that  $g \circ h = f$ .

This definition was given initially for modules over an integral domain by E. Enochs [3] who proved that, in this case, every module has a unique (up to isomorphism) torsion free covering module. B. Banaschewski [1, p. 59] gave the following construction for the cover of a module  $E$  over an integral domain  $R$  with quotient field  $K$ :  $T(E) = \{f \in \text{Hom}_R(K, I(E)) \mid f(1) \in E\}$ ;  $g(f) = f(1)$ .

M. Teply [8, p. 449] generalized the notion of a torsion free cover to a hereditary torsion theory  $(\mathcal{T}, \mathcal{F})$  [2] of  $R$ -modules. He proved that each module has a  $\mathcal{F}$ -torsion free cover if  $R \in \mathcal{F}$  and the direct sum of  $\mathcal{F}$ -torsion free injective modules is injective.

The Goldie torsion theory  $(\mathcal{G}, \mathcal{F})$  is the torsion theory whose torsion class  $\mathcal{G}$  is generated by all factor modules  $B/A$  where  $A$  is a large submodule of  $B$ . The Goldie torsion free class  $\mathcal{F}$  is precisely the class of torsion free modules given by Definition 1. Teply's result shows that  $\mathcal{G}$ -torsion free covers exist if  $Z(R) = (0)$  and  $\dim R$  is finite. We prove the converse and show that Banaschewski's construction has an obvious analogy in this case.

**THEOREM 3.** *Let  $R$  be a ring with identity and maximal left quotient ring  $Q$ . If every left  $R$ -module has a torsion free cover then  $Z(R) = (0)$  and  $\dim R$  is finite. Moreover, the evaluation map from  $T(E) = \{f \in \text{Hom}_R(Q, I(E)) \mid f(1) \in E\}$  onto  $E$  is a torsion free cover of  $E$ .*

*Proof.* Let  $(0) \rightarrow \text{Ker } g \rightarrow T(R) \xrightarrow{g} R \rightarrow (0)$  be a torsion free cover of the module  $R$ . Since  $R$  is projective this sequence splits and hence  $\text{Ker } g$  is closed in  $T(R)$ . Then  $\text{Ker } g = (0)$  as  $g$  is a cover of  $R$  so  $R \cong T(R) \in \mathcal{F}$ , i.e.,  $Z(R) = (0)$ .

To show that  $\dim R$  is finite it suffices, by Theorem S, to show that  $Q$  is a semisimple ring. But, since  $Z(R) = (0)$ ,  $Q$  is von Neumann regular so it suffices to show that  $Q$  is a finite dimensional ring. If not, there is an infinite set of nonzero elements  $\{x_i: i \in I\}$  of  $Q$  such that the sum  $B = \sum Qx_i$  is direct and a proper large left ideal of  $Q$ . Then  $Q/B \neq (0)$  has a torsion free cover (as an  $R$ -module), say  $g: F \rightarrow Q/B$ .

Since  $Z({}_R Q) = (0)$  the natural  $R$ -homomorphism  $p: Q \rightarrow Q/B$  factors thru  $g$ , i.e., there exist  $h \in \text{Hom}_R(Q, F)$  such that  $g \circ h = p$ . Then  $h(B) \subseteq \text{Ker } g$  and  $p \neq 0$  implies  $h \neq 0$  so by Lemma 3(1)  $h(B) \neq (0)$  — we must note that  $B$  is a large  $R$ -submodule of  $Q$ . Then  $h(Qx_j) \neq (0)$  for some index  $j \in I$ . By Lemma 3(2)  $h(Qx_j)$  is a direct summand of  $F$ . Thus we have a nonzero closed submodule of  $F$  contained in

$\text{Ker } g$ . This gives a contradiction so  $Q$  is a finite dimensional ring, hence semisimple.

Since  $Q$  is a semisimple ring it follows that any  $Q$ -module is torsion free when considered as an  $R$ -module. Hence any  $R$ -submodule of such a module is a torsion free  $R$ -module. Conversely, it follows from Theorem  $S(f)$  that any torsion free  $R$ -module is an  $R$ -submodule of a  $Q$ -module (i.e., is  $Q$ -extendible in Banaschewski's terminology). Banaschewski [1, p. 63] established the existence of  $Q$ -extendible coverings.

Note that if  $q_1, q_2, \dots, q_n \in Q$ , and  $f_1, f_2, \dots, f_n \in T(E) = \{f \in \text{Hom}_R(Q, I(E)) \mid f(1) \in E\}$  are such that  $\sum q_i f_i = 0 \in \text{Hom}_R(Q, I(E))$  then the large left ideal  $I = \{r \in R \mid r q_i \in R \text{ for all } i\}$  annihilates  $\sum q_i \otimes f_i$  in  $Q \otimes_R T(E)$ . Thus  $\sum q_i \otimes f_i \in Z(Q \otimes_R T(E)) = (0)$ . This shows that the  $R$ -homomorphism  $Q \otimes_R T(E) \rightarrow \text{Hom}_R(Q, I(E))$  given by the  $Q$ -module structure of  $\text{Hom}_R(Q, I(E))$  is one-to-one. Therefore by [1, Prop. 3, p. 64] the evaluation map from  $T(E)$  onto  $E$  is a torsion free cover of  $E$ . This completes the proof.

REMARK. This theorem shows that "torsion free" covers do not exist for the ring mentioned by Banaschewski [1, p. 66], that is, the ring of all functions on a set  $X$  with values in a field which are constant except on some finite set. Since  $Z(R) = (0)$  the "torsion free" he is using agrees with our torsion free and covers do not exist.

This paper constitutes a portion of the author's doctoral dissertation at the University of Kentucky. The author is deeply indebted to Professor Edgar Enochs, his advisor, for his generous advice and encouragement.

#### REFERENCES

1. B. Banaschewski, *On covering of modules*, Math. Nachr., **31** (1966), 57-71.
2. S. E. Dickson, *A torsion theory for abelian categories*, Trans. Amer. Math. Soc., **121** (1966), 223-225.
3. E. Enochs, *Torsion free covering modules*, Proc. Amer. Math. Soc., **14** (1963), 884-889.
4. J. Lambek, *Lectures on rings and modules*, Blaisdell Publishing Co., Waltham, Mass., 1966.
5. B. Maddox, *Absolutely pure modules*, Proc. Amer. Math. Soc., **18** (1967), 155-158.
6. C. Megibben, *Absolutely pure modules*, Proc. Amer. Math. Soc., **26** (1970), 561-566.
7. F. Sandomierski, *Semisimple maximal quotient rings*, Trans. Amer. Math. Soc., **128** (1967), 112-120.
8. F. Sandomierski, *Nonsingular rings*, Proc. Amer. Math. Soc., **19** (1968), 225-230.
9. M. Teply, *Torsion free injective modules*, Pacific J. Math., **28** (1969), 441-453.
10. ———, *Some aspects of Goldie's torsion theory*, Pacific J. Math., **29** (1969), 447-459.

Received April 2, 1971 and in revised form July 15, 1971.

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON  
Stanford University  
Stanford, California 94305

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

C. R. HOBBY  
University of Washington  
Seattle, Washington 98105

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NAVAL WEAPONS CENTER



Charles A. Akemann, <i>A Gelfand representation theory for <math>C^*</math>-algebras</i> . . . .	1
Sorrell Berman, <i>Spectral theory for a first-order symmetric system of ordinary differential operators</i> . . . . .	13
Robert L. Bernhardt, III, <i>On splitting in hereditary torsion theories</i> . . . . .	31
J. L. Brenner, <i>Geršgorin theorems, regularity theorems, and bounds for determinants of partitioned matrices. II. Some determinantal identities</i> . . . . .	39
Robert Morgan Brooks, <i>On representing <math>F^*</math>-algebras</i> . . . . .	51
Lawrence Gerald Brown, <i>Extensions of topological groups</i> . . . . .	71
Arnold Barry Calica, <i>Reversible homeomorphisms of the real line</i> . . . . .	79
J. T. Chambers and Shinnosuke Oharu, <i>Semi-groups of local Lipschitzians in a Banach space</i> . . . . .	89
Thomas J. Cheatham, <i>Finite dimensional torsion free rings</i> . . . . .	113
Byron C. Drachman and David Paul Kraines, <i>A duality between transpotence elements and Massey products</i> . . . . .	119
Richard D. Duncan, <i>Integral representation of excessive functions of a Markov process</i> . . . . .	125
George A. Elliott, <i>An extension of some results of Takesaki in the reduction theory of von Neumann algebras</i> . . . . .	145
Peter C. Fishburn and Joel Spencer, <i>Directed graphs as unions of partial orders</i> . . . . .	149
Howard Edwin Gorman, <i>Zero divisors in differential rings</i> . . . . .	163
Maurice Heins, <i>A note on the Löwner differential equations</i> . . . . .	173
Louis Melvin Herman, <i>Semi-orthogonality in Rickart rings</i> . . . . .	179
David Jacobson and Kenneth S. Williams, <i>On the solution of linear G.C.D. equations</i> . . . . .	187
Michael Joseph Kallaher, <i>On rank 3 projective planes</i> . . . . .	207
Donald Paul Minassian, <i>On solvable <math>O^*</math>-groups</i> . . . . .	215
Nils Øvrelid, <i>Generators of the maximal ideals of <math>A(\bar{D})</math></i> . . . . .	219
Mohan S. Putcha and Julian Weissglass, <i>A semilattice decomposition into semigroups having at most one idempotent</i> . . . . .	225
Robert Raphael, <i>Rings of quotients and <math>\pi</math>-regularity</i> . . . . .	229
J. A. Siddiqi, <i>Infinite matrices summing every almost periodic sequence</i> . . . .	235
Raymond Earl Smithson, <i>Uniform convergence for multifunctions</i> . . . . .	253
Thomas Paul Whaley, <i>Multiplicity type and congruence relations in universal algebras</i> . . . . .	261
Roger Allen Wiegand, <i>Globalization theorems for locally finitely generated modules</i> . . . . .	269