

Pacific Journal of Mathematics

A NOTE ON THE LÖWNER DIFFERENTIAL EQUATIONS

MAURICE HEINS

A NOTE ON THE LÖWNER DIFFERENTIAL EQUATIONS

MAURICE HEINS

Charles Löwner in Memoriam

The object of the present note is to indicate a derivation of the Löwner differential equations [1] based on the derivation of an associated differential equation for Green's function of the variable region relative to the defining parameter. Decisive in our treatment is the use of a certain normalized minimal positive harmonic function on the variable region. In fact, our starting point was the feeling that the Poisson kernel asserted its presence so strongly in the Löwner differential equations that the concomitant presence of a normalized minimal positive harmonic function on the variable region should appear naturally in the study of the question. We shall see that this is the case. A technical advantage of the present approach is that the "tip" lemmas of the classical proof are dispensed with.

It would be of interest to see whether the indicated method, which is available for other families of harmonic functions monotone justifying in a parameter, has useful applications to the theory of harmonic functions.

2. Let γ be a Jordan arc with parametric domain $[0, T]$ such that $0 < |\gamma(t)| < 1$ for $0 \leq t < T$ and $|\gamma(T)| = 1$. Let A_t denote the complement of the set $\gamma(\{t \leq s < T\})$ with respect to the open unit disk, $0 \leq t \leq T$. Let g_t denote Green's function for A_t with pole at 0. The continuous dependence of g_t on the parameter t is an elementary matter (minimal property of Green's function, the Phragmén-Lindelöf boundary maximum principle). We let $\alpha(t)$ denote $\lim_{z \rightarrow 0} [g_t(z) + \log |z|]$. We note that $\alpha: t \rightarrow \alpha(t)$ is an increasing continuous function which satisfies $\alpha(T) = 0$. We reparametrize γ , as in the original Löwner argument, by composing γ with

$$t \longrightarrow \text{inv } \alpha[t + \alpha(0)], \quad 0 \leq t \leq -\alpha(0),$$

so that for the new γ we have $T = -\alpha(0)$ and $\alpha(t) = \alpha(0) + t$. [The notation "inv" is used to denote the inverse of a univalent function.]

We let G be defined by

$$G(z, t) = g_t(z), \quad z \in A_t, \quad 0 \leq t \leq T.$$

Given a function F having as domain a subset of $C \times R$, we denote by $D_2F(a, b)$ the derivative of $t \rightarrow F(a, t)$ at b and by $D_1F(a, b)$ and $\bar{D}_1F(a, b)$ the complex differential coefficients of $z \rightarrow F(z, b)$ at a , the obvious conventions holding. Our first step is to establish the existence of D_2G and to obtain information about it. We remark that the logarithmic singularity of g_t at 0 is harmless. The difference quotient

$$(1) \quad \frac{G(z, t) - G(z, s)}{t - s}, \quad 0 \leq s < t \leq T,$$

defines a positive harmonic function on A_s , which takes the value 1 at 0 and vanishes continuously at each point of the frontier of A_s .

To control the limiting behavior of (1) as $(s, t) \rightarrow (\sigma, \sigma)$, $0 \leq \sigma \leq T$, we make use of the boundary behavior of the Riemann mapping function for a simply-connected Jordan region and the following standard lemma of Harnack type.

LEMMA. Let $m(z) = (1 - |z|)(1 + |z|)^{-3}$ and $M(z) = (1 + |z|) \times (1 - |z|)^{-3}$. Let a and b be points of the semi-circular disk $\{\text{Im } z > 0, |z| < 1\}$. Let u be nonnegative and harmonic on this set and vanish continuously on the diameter. Then

$$[M(b)]^{-1} \frac{u(b)}{\text{Im } b} \leq [m(a)]^{-1} \frac{u(a)}{\text{Im } a}.$$

[A proof of this lemma is readily given with the aid of Schwarzian reflexion and the Poisson integral for a circular disk.]

Suppose that $(s_n, t_n) \rightarrow (\sigma, \sigma)$, where $0 \leq s_n < t_n \leq 1$. Then some subsequence of the sequence of difference quotients (1), given by $s = s_n$ and $t = t_n$, converges, uniformly on compact subsets of A_σ , to a positive harmonic function on A_σ which takes the value 1 at 0. Using the boundary behavior of the Riemann mapping function when a Jordan boundary lies at hand and the stated lemma, we see that the limit function in question vanishes continuously at each point of the frontier of A_σ , the "tip" $\gamma(\sigma)$ excepted.

We introduce the normalized Riemann mapping function f_t , mapping the open unit disk onto A_t and satisfying $f_t(0) = 0$, $f_t'(0) > 0$. From the continuity of $t \rightarrow g_t$, we infer the continuity of $t \rightarrow \text{inv } f_t$ and thence the continuity of $t \rightarrow f_t$. Of course, the term "continuity" is to be construed in the sense of uniform limits on compact subsets. We let $\kappa(t)$ denote the unique preimage of $\gamma(t)$ with respect to the continuous extension of f_t to the closed unit disk. If h is a positive harmonic function on A_t taking the value 1 at 0 and vanish-

ing continuously at each point of the frontier of A_t , the “tip” $\gamma(t)$ excepted, then

$$(2) \quad h[f_t(z)] = \operatorname{Re} \left[\frac{\kappa(t) + z}{\kappa(t) - z} \right], \quad |z| < 1.$$

It follows that there is at most one h having the stated property. Using (2) as a defining condition for h , we see that such h exist. We denote the unique h in question (which is a normalized minimal positive harmonic function on A_t) by h_t .

Combining the results of the preceding two paragraphs we conclude that the difference quotient (1) tends to h_σ as $(s, t) \rightarrow (\sigma, \sigma)$ and that, in fact, the uniformity of the limit process holds on compact subsets of A_σ . We let H be defined by

$$H(z, t) = h_t(z), \quad z \in A_t, \quad 0 \leq t \leq T.$$

We see that the following differential equation, which will serve as a basis for the derivation of the Löwner differential equations, holds:

$$(3) \quad D_2 G = H.$$

Continuity of $t \rightarrow h_t$ and $\kappa: t \rightarrow \kappa(t)$. A second application of the boundary behavior of the Riemann mapping function for Jordan regions and the lemma yields the continuity of $t \rightarrow h_t, 0 \leq t \leq T$. It suffices to establish the fact that if $t_n \rightarrow \sigma$, then some subsequence of (h_{t_n}) tends to h_σ . Using the continuity of $t \rightarrow h_t$, the continuity of $t \rightarrow f_t$, and (2) we shall now conclude the continuity of κ . Indeed, if $t_n \rightarrow \sigma$ and $\kappa(t_n) \rightarrow \alpha$, we obtain, using (2), the equality

$$h_\sigma[f_\sigma(z)] = \operatorname{Re} \left(\frac{\alpha + z}{\alpha - z} \right), \quad |z| < 1,$$

and hence $\alpha = \kappa(\sigma)$. The continuity of κ follows.

3. The Löwner differential equations. The equations bear on the functions F, Ψ , and θ , which will now be introduced.

F. We define F by $F(z, t) = f_t(z), |z| < 1, 0 \leq t \leq T$. It is convenient to have available Φ defined by $\Phi(z, t) = \operatorname{inv} f_t(z), z \in A_t, 0 \leq t \leq T$. Its role is auxiliary. The function Φ is useful as a link between F and $G = -\log |\Phi|$.

\Psi. The function Ψ is defined by $\Psi(z, t) =$

$\operatorname{inv} f_t[f_0(z)], |z| < 1, 0 \leq t \leq T$. This is the first function studied by Löwner in his classical paper. There is an identity involving Φ and Ψ :

$$(4) \quad \Psi(z, t) = \Phi[f_0(z), t].$$

θ . The function θ is specified by the requirement that, $z \rightarrow \theta(z, t)$ is the inverse of $z \rightarrow \Psi(z, t)$, $|z| < 1$, $0 \leq t \leq T$. From

$$z = \Psi[\theta(z, t), t] = \text{inv } f_t \{f_0[\theta(z, t)]\}$$

for (z, t) in the domain of θ , we obtain for such (z, t) the identity

$$(5) \quad F(z, t) = f_t(z) = f_0[\theta(z, t)].$$

The equation (3) yields a corresponding equation for Φ . Indeed, let \tilde{h}_t denote the analytic function with domain A_t satisfying $\tilde{h}_t(0) = 1$, $\text{Re } \tilde{h}_t = h_t$, and let \tilde{H} be defined by $\tilde{H}(z, t) = \tilde{h}_t(z)$, $z \in A_t$, $0 \leq t \leq T$. Clearly, the function \tilde{H} is continuous on its domain. To derive an equation bearing on Φ , we introduce G_1 having the same domain as G which satisfies $G_1(z, t) = G(z, t) + \log |z|$, $z \neq 0$, $G_1(0, t) = \alpha(0) + t$, and thereupon \tilde{G}_1 with the same domain and satisfying the condition that $z \rightarrow \tilde{G}_1(z, t)$ is the analytic function with real part $z \rightarrow G_1(z, t)$ satisfying $\tilde{G}_1(0, t) = G_1(0, t)$. It is readily verified that

$$D_2G_1 = H,$$

the limit process being uniform in the sense indicated above. It follows, in view of the normalization made on \tilde{G}_1 , that

$$D_2\tilde{G}_1 = \tilde{H}.$$

Using the relation

$$\Phi(z, t) = z \exp [-\tilde{G}_1(z, t)],$$

we are led to the equation

$$(6) \quad D_2\Phi = -\tilde{H}\Phi.$$

From (4) and (6) we obtain

$$\begin{aligned} D_2\Psi(z, t) &= -\tilde{H}[f_0(z), t]\Psi(z, t) \\ &= -\tilde{h}_t[f_0(z)]\Psi(z, t) \\ &= -[(\tilde{h}_t \circ f_t) \circ (\text{inv } f_t \circ f_0)](z)\Psi(z, t), \end{aligned}$$

and, consequently, the Löwner equation

$$(7) \quad D_2\Psi(z, t) = -\frac{\kappa(t) + \Psi(z, t)}{\kappa(t) - \Psi(z, t)} \Psi(z, t),$$

(z, t) in the domain of Ψ .

The equation for F . From (6) and the continuity of \tilde{H} , Φ , $D_1\Phi$, $\tilde{D}_1\Phi$ (trivially, since it vanishes), we conclude that Φ has the C' property and so is differentiable. Since F is continuous on its domain, $D_1\Phi$ is nowhere zero, and the identity,

$$\Phi[F(z, t), t] = z,$$

$|z| < 1$, $0 \leq t \leq T$, prevails, it is a standard matter of the differential calculus to conclude the existence of D_2F and the identity,

$$(8) \quad D_1\Phi[F(z, t), t]D_2F(z, t) + D_2\Phi[F(z, t), t] = 0, \\ |z| < 1, 0 \leq t \leq T.$$

It is elementary that

$$(9) \quad D_1\Phi[F(z, t), t]D_1F(z, t) = 1,$$

for the same (z, t) . From (8) and (6) we obtain

$$D_1\Phi[F(z, t), t]D_2F(z, t) = z \frac{\kappa(t) + z}{\kappa(t) - z},$$

and thereupon using (9) the equation

$$(10) \quad D_2F(z, t) = D_1F(z, t)z \frac{\kappa(t) + z}{\kappa(t) - z},$$

$$|z| < 1, 0 \leq t \leq T.$$

That θ satisfies the equation (10), θ replacing F , on its domain, is immediate from (10), and the identities obtained from (5) by differentiation. The C' property of F follows from the continuity of $D_1\Phi$, $D_2\Phi$ and F on their respective domains and the identities (8) and (9) as well as the non-vanishing of $D_1\Phi$. The C' property θ is now concluded with the aid of (5).

REFERENCE

1. K. Löwner, *Untersuchungen über schlichte konforme Abbildungen des Einheitskreises I*, Math. Ann., **89** (1923), 103-121.

Received December 7, 1970.

UNIVERSITY OF ILLINOIS

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Charles A. Akemann, <i>A Gelfand representation theory for C^*-algebras</i>	1
Sorrell Berman, <i>Spectral theory for a first-order symmetric system of ordinary differential operators</i>	13
Robert L. Bernhardt, III, <i>On splitting in hereditary torsion theories</i>	31
J. L. Brenner, <i>Geršgorin theorems, regularity theorems, and bounds for determinants of partitioned matrices. II. Some determinantal identities</i>	39
Robert Morgan Brooks, <i>On representing F^*-algebras</i>	51
Lawrence Gerald Brown, <i>Extensions of topological groups</i>	71
Arnold Barry Calica, <i>Reversible homeomorphisms of the real line</i>	79
J. T. Chambers and Shinnosuke Oharu, <i>Semi-groups of local Lipschitzians in a Banach space</i>	89
Thomas J. Cheatham, <i>Finite dimensional torsion free rings</i>	113
Byron C. Drachman and David Paul Kraines, <i>A duality between transpotence elements and Massey products</i>	119
Richard D. Duncan, <i>Integral representation of excessive functions of a Markov process</i>	125
George A. Elliott, <i>An extension of some results of Takesaki in the reduction theory of von Neumann algebras</i>	145
Peter C. Fishburn and Joel Spencer, <i>Directed graphs as unions of partial orders</i>	149
Howard Edwin Gorman, <i>Zero divisors in differential rings</i>	163
Maurice Heins, <i>A note on the Löwner differential equations</i>	173
Louis Melvin Herman, <i>Semi-orthogonality in Rickart rings</i>	179
David Jacobson and Kenneth S. Williams, <i>On the solution of linear G.C.D. equations</i>	187
Michael Joseph Kallaher, <i>On rank 3 projective planes</i>	207
Donald Paul Minassian, <i>On solvable O^*-groups</i>	215
Nils Øvrelid, <i>Generators of the maximal ideals of $A(\bar{D})$</i>	219
Mohan S. Putcha and Julian Weissglass, <i>A semilattice decomposition into semigroups having at most one idempotent</i>	225
Robert Raphael, <i>Rings of quotients and π-regularity</i>	229
J. A. Siddiqi, <i>Infinite matrices summing every almost periodic sequence</i>	235
Raymond Earl Smithson, <i>Uniform convergence for multifunctions</i>	253
Thomas Paul Whaley, <i>Multiplicity type and congruence relations in universal algebras</i>	261
Roger Allen Wiegand, <i>Globalization theorems for locally finitely generated modules</i>	269