

Pacific Journal of Mathematics

ON RANK 3 PROJECTIVE PLANES

MICHAEL JOSEPH KALLAHER

ON RANK 3 PROJECTIVE PLANES

MICHAEL J. KALLAHER

One of the unsolved problems in the theory of projective planes is the following: Is every finite projective plane with a transitive collineation group desarguesian? This problem is investigated under the additional hypothesis that the group has rank 3. It is proven that if a projective plane \mathcal{P} of order $n > 2$ has a rank 3 collineation group then \mathcal{P} is non-desarguesian and either (i) n is odd and $n = m^4$, or (ii) n is even and $n = m^2$ with $m \equiv 0 \pmod{4}$.

One of the older problems in the theory of projective planes is the following: If a finite projective plane \mathcal{P} has a collineation group G transitive on its points, is the plane desarguesian? So far only under one or more additional assumptions has the answer been shown to be yes. The basis for these results is the theorem, due to Wagner [10], that if the transitive group G contains a central collineation, then \mathcal{P} is desarguesian.

Ostrom and Wagner [8] showed that if the group G is doubly transitive then \mathcal{P} is desarguesian, and G contains all elations of \mathcal{P} . Higman and McLaughlin [4] investigated the problem in the case when the group G is transitive on the flags of \mathcal{P} . They proved that under certain restrictions on the order of \mathcal{P} the plane is desarguesian. Keiser [6] and Wagner [10] have shown that under restrictions on the order of \mathcal{P} and the order of G the plane is desarguesian.

The rank of a permutation group G transitive on a set Ω is the number of orbits of G_P , P a point of Ω , in Ω . Hence a transitive group G has rank 2 on a set Ω if and only if G is doubly transitive on Ω . G has rank 3 if and only if for every point $P \in \Omega$ G_P has two orbits besides G_P . Ostrom and Wagner have thus answered the question when the group G has rank 2. It is then natural to ask: If a finite projective plane has a transitive collineation group of rank 3, is the plane desarguesian?

Investigating this question we have found that a more appropriate question is: Which finite projective planes have rank 3 collineation groups? For we will prove in this article the following

MAIN THEOREM. *Let \mathcal{P} be a finite projective plane of order n with rank 3 collineation group G . Then n satisfies one of the statements:*

- (i) $n = 2$
- (ii) n is odd and $n = m^4$

(iii) n is even and $n = m^2$ with $m \equiv 0 \pmod{4}$

Furthermore only in case (i) is \mathcal{P} desarguesian.

The proof consists in showing that G must be a nonsolvable flag-transitive group of even order. It is not known whether \mathcal{P} can actually exist in cases (ii) or (iii). Hence the theorem leads to the following conjecture:

CONJECTURE. The only finite projective plane having a rank 3 collineation group is the desarguesian plane of order two.

The desarguesian plane of order two (Fig. 1) has the rank 3 collineation group G generated by the collineations $\sigma = (P_1P_7P_6P_4P_3P_2P_5)$ and $\tau = (P_7P_6P_3)(P_4P_5P_2)$. Note that G is solvable, sharply flag-transitive, and has order 21.




FIGURE 1

We wish to thank Professor Ostrom for many helpful suggestions and for reading a preliminary draft.

2. Definitions and results required later. We assume the reader is familiar with the theory and terminology of projective planes as appears, for example, in Chapters 3–5 of Dembowski [3]. Also a familiarity with the simpler aspects of permutation group theory (as in Chapter 1 of Wielandt [12]) will be assumed.

A *transitive collineation group* on a projective plane \mathcal{P} is one which is transitive on the points (and hence on the lines by Result II below). A *rank 3 collineation group* of \mathcal{P} is a transitive collineation group which has rank 3 as a permutation group on the points of \mathcal{P} . A *flag* of \mathcal{P} is an incident point-line pair; i.e., a pair P, l with P a point, l a line, of \mathcal{P} and $P \in l$. A *flag-transitive collineation group* is one which is transitive on the flags of \mathcal{P} .

Use will be made of a number of results concerning permutations

and collineations. These are listed below for convenience:

Result I (Ostrom [27]; Dembowski [3], p. 214): If \mathcal{P} is a finite projective plane having a collineation group G which is transitive on non-incident point-line pairs, then G is doubly transitive on the points of \mathcal{P} .

Result II (Dembowski [2], Hughes [5], Parker [9]): A collineation group of a finite projective plane has equally many point and line orbits.

Result III (Higman and McLaughlin [4]): Let \mathcal{P} be a finite projective plane of order n with a flag-transitive collineation group G . If n is odd and not a fourth power, then \mathcal{P} is desarguesian and G contains all elations of \mathcal{P} (See also Dembowski [3], p. 212).

Result IV (Higman and McLaughlin [4]): Let G be a flag-transitive collineation group of a desarguesian projective plane \mathcal{P} of order n . G contains all elations of \mathcal{P} with precisely two exceptions:

- (i) $n = 2$ and G has order 21.
- (ii) $n = 8$ and G has order 657.

Result V (Wagner [10]): If \mathcal{P} is a finite projective plane having a transitive collineation group G which contains a nontrivial central collineation, then \mathcal{P} is desarguesian and G contains all elations of \mathcal{P} .

Result VI (Keiser [6]): Let \mathcal{P} be a finite projective plane of order n with a transitive collineation group G . If G is nonsolvable and if $n = m^2$ with $m \equiv 2 \pmod{4}$ or $m \equiv 3 \pmod{4}$, then \mathcal{P} is desarguesian and G contains all the elations of \mathcal{P} .

Results VII (Dembowski [3], p. 212): Let \mathcal{P} be a finite projective plane of order n . If G is a collineation group which is solvable and primitive on the points of \mathcal{P} , then $n^2 + n + 1$ is a prime.

For the next result we note that a permutation group on a set Ω is *regular* (*sharply transitive*) if it is transitive and G_P consists only of the identity for each point $P \in \Omega$. G is a *Frobenius group* on Ω if (i) it is transitive, (ii) G_P is nontrivial for each point $P \in \Omega$, and (iii) for every two distinct points $P, Q \in \Omega$ $G_{P,Q}$ consists only of the identity.

Result VIII (Wielandt [11], 11.6): A transitive permutation group of prime degree is solvable if and only if it is either regular or a Frobenius group (Due to E. Galois).

3. The investigation. In this section we prove that if a finite projective plane \mathcal{P} has a rank 3 group G of collineations then G is

flag-transitive on \mathcal{P} , G is nonsolvable (if $n > 2$), and \mathcal{P} is not desarguesian (if $n > 2$). We start with

LEMMA 1. *Let \mathcal{P} be a finite projective plane and G a rank 3 group of collineation of \mathcal{P} . Then for every point P of \mathcal{P} , G_P permutes the lines through P in one or two orbits.*

Proof. G_P permutes the points different than P in two orbits \mathcal{O}_1 and \mathcal{O}_2 . Define the sets $\mathcal{M}_i, i = 1, 2$, by: \mathcal{M}_i consists of the lines through P such that l contains at least one point in \mathcal{O}_i . G_P is transitive on \mathcal{M}_1 and \mathcal{M}_2 , and every line through P is either in \mathcal{M}_1 or \mathcal{M}_2 . If $\mathcal{M}_1 \cap \mathcal{M}_2$ is empty, then G_P permutes the lines through P in two orbits. If $\mathcal{M}_1 \cap \mathcal{M}_2$ is nonempty, then G_P is transitive on the lines through P .

THEOREM 1. *Let \mathcal{P} be a finite projective plane and G a rank 3 group of collineations of \mathcal{P} . Then G_P is flag-transitive.*

Proof. Assume G is not flag-transitive. For every point P of \mathcal{P} G_P has 3 point orbits in \mathcal{P} . Hence by Result II G_P has three line orbits in \mathcal{P} . If all the lines through P are in a single orbit under G_P , then G is flag-transitive contrary to our assumption. Thus by Lemma 1, two of these line orbits consist of the lines through P . Thus the third line orbit of G_P must consist of all the lines of \mathcal{P} which do not go through P . Hence for every point P , G_P is transitive on all lines not through P .

Let (P, l) be a non-incident point-line pair of \mathcal{P} and (Q, m) another non-incident point-line pair. There exists a collineation $\sigma \in G$ such that $P\sigma = Q$. Let \bar{l} be the image of l under σ . $Q \notin \bar{l}$ since $P \notin l$. Then there exists a collineation $\tau \in G_Q$ such that $\bar{l}\tau = m$. Then $P\sigma\tau = Q\tau = Q$ and $l\sigma\tau = \bar{l}\tau = m$. This proves G is transitive on non-incident point-line pairs.

Result I implies that G is doubly transitive on the points of \mathcal{P} . This is a contradiction since G has rank 3 on the points of \mathcal{P} . Thus G is flag-transitive.

LEMMA 2. *Let \mathcal{P} be a finite projective plane of order n and G a rank 3 group of collineations of \mathcal{P} , and let P a point of \mathcal{P} .*

(i) G_P permutes the points not equal to P in two orbits \mathcal{O}_1 and \mathcal{O}_2 of lengths $k_1(n+1)$ and $k_2(n+1)$ respectively, where $k_1 + k_2 = n$.

(ii) G_P permutes the lines not through P in two orbits \mathcal{L}_1 and \mathcal{L}_2 of lengths s_1 and s_2 respectively, with $s_1 + s_2 = n^2$.

Proof. G_P is transitive on the lines through P . Thus every line

through P intersects the non-trivial point orbits \mathcal{O}_1 in the same number of points. Thus $|\mathcal{O}_1| = k_1(n + 1)$ where k_1 is the number of points of \mathcal{O}_1 on a line l through P . Similarly $|\mathcal{O}_2| = k_2(n + 1)$, k_2 the number of points of \mathcal{O}_2 on a line l through P . This proves (i) since for every line l through P , a point $\neq P$ on l is either in \mathcal{O}_1 or \mathcal{O}_2 .

G_P has three point orbits. Hence it has three line orbits (Result II). One of these consists of the lines through P . The other two line orbits are made up of the lines not through P and there are n^2 such lines. This gives (ii).

LEMMA 3. *Let \mathcal{P} be a finite projective plane of order n and G a rank 3 group of collineations of \mathcal{P} . If G is solvable, then*

- (i) $n^2 + n + 1$ is a prime;
- (ii) G acts as a Frobenius group on \mathcal{P} ;
- (iii) n is even, $|G| = 1/2(n^2 + n + 1)n(n + 1)$, and $|G_P| = 1/2n(n + 1)$.

If $|G|$ is odd, then we also have

- (iv) $n = 2m$, m odd.

Proof. Since G is flag-transitive, G is primitive on the points of \mathcal{P} (Dembowski [3], p. 212). By Result VII $n^2 + n + 1$ is a prime. Result VIII implies that G is either regular on \mathcal{P} or it is a Frobenius group on \mathcal{P} . Since G is clearly not regular it must be a Frobenius group on \mathcal{P} . This proves (i) and (ii).

By Lemma 2 we have for every point $P \in \mathcal{P}$

$$|G_P| = k_1(n + 1) |G_{P,Q}| = k_2(n + 1) |G_{P,R}|,$$

where $Q \in \mathcal{O}_1$ and $R \in \mathcal{O}_2$, and $k_1 + k_2 = n$. But G a Frobenius group on \mathcal{P} implies $|G_{P,Q}| = 1 = |G_{P,R}|$. Hence $|G_P| = k_1(n + 1) = k_2(n + 1)$ and thus $k_1 = k_2 = n/2$. This implies n is even since k_1 is an integer, and we have $|G| = 1/2(n^2 + n + 1)(n(n + 1))$. This proves (iii).

If $|G|$ is odd, then $n/2$ is odd and this proves (iv).

LEMMA 4. *Let \mathcal{P} be a finite projective plane of order n with a rank 3 group of collineations. If $n > 2$, then $|G|$ is even.*

Proof. Assume $|G|$ is odd. Then G is solvable (by the Feit-Thompson theorem) and the previous lemma implies $n^2 + n + 1$ is a prime, $n = 2m$ with m odd, and $|G| = (n^2 + n + 1)m(n + 1)$. Also for each point P , G_P has two line orbits \mathcal{L}_1 and \mathcal{L}_2 of lengths s_1 and s_2 respectively with

$$(1) \quad s_1 + s_2 = n^2 = 4m^2$$

(Lemma 2). Since $|G_P| = m(n + 1) = m(2m + 1)$, we have

$$(2) \quad s_i \mid m(2m + 1), \quad i = 1, 2.$$

Let $s = \text{g.c.d.}(s_1, s_2)$ and define integers t_1 and t_2 by

$$(3) \quad s_i = st_i, \quad i = 1, 2.$$

Then $\text{g.c.d.}(t_1, t_2) = 1$ and $s(t_1 + t_2) = 4m^2$ (by (1)). Since s is odd (for otherwise $|G_{\mathcal{P}}|$ is even), $s \mid m^2$. Since $s \mid s_1$ (2) implies $s \mid m$ since $\text{g.c.d.}(m, 2m + 1)$. Hence

$$(4) \quad m = su$$

for some integer u , and

$$(5) \quad t_1 + t_2 = 4su^2.$$

If $v = \text{g.c.d.}(t_1, s) > 1$, then $v \mid t_2 - a$ contradiction to the fact that $\text{g.c.d.}(t_1, t_2) = 1$. If $w = \text{g.c.d.}(t_1, u) > 1$, then $w \mid t_2$ again a contradiction. Hence $1 = \text{g.c.d.}(t_1, s) = \text{g.c.d.}(t_1, u)$, which implies $\text{g.c.d.}(t_1, m) = 1$. Similarly $\text{g.c.d.}(t_2, m) = 1$ and thus

$$(6) \quad t_i \mid 2m + 1, \quad i = 1, 2.$$

Then we have, using (5)

$$2(2m + 1) \geq t_1 + t_2 = 4su^2.$$

Applying (4) we get

$$1 \geq 2su(u - 1).$$

Therefore $u = 1$ and

$$(7) \quad m = s.$$

But then

$$(8) \quad t_1 + t_2 = 4m.$$

If $t_1 = t_2$, then $t_1 = 2m$ and this contradicts (6). Thus without loss of generality we may assume $t_1 < t_2$. Then from (8) we get $t_1 < 2m$, $t_2 > 2m$ and this implies $t_2 = 2m + 1$ (by (6)) and $t_1 = 2m - 1$ (by (8)). Hence $2m - 1 \mid 2m + 1$, which implies $m = 1, n = 2$. This proves the lemma.

REMARK. The example at the end of §1 shows that if $n = 2 \mid |G|$ can be odd.

By combining Lemma 3 and Lemma 4 we can show that the rank 3 group G is nonsolvable if $n > 2$:

THEOREM 2. *Let \mathcal{P} be a finite projective plane of order n with a rank 3 group G of collineations. If $n > 2$, then G is nonsolvable.*

Proof. Assume G is solvable and $n > 2$. By Lemma 3(ii) G acts as a Frobenius group on \mathcal{P} . Lemma 4 implies $|G|$ is even. Hence G has an element σ of order 2. Either σ is a central collineation or it fixes a subplane of \mathcal{P} pointwise (Baer [1]). In both cases σ fixes more than two points. But $G_{P,Q}$ consists only of the identity for every two distinct points P, Q of \mathcal{P} . This gives a contradiction. Therefore G is nonsolvable if $n > 2$.

Our last result in this section shows that \mathcal{P} is desarguesian only when $n = 2$.

THEOREM 3. *Let \mathcal{P} be a finite projective plane with a rank 3 group G of collineations. \mathcal{P} is desarguesian if and only if $n = 2$.*

Proof. Assume \mathcal{P} is desarguesian. By Theorem 1 G is flag-transitive. G cannot contain all the elations of \mathcal{P} . For the group H generated by the elations of \mathcal{P} is doubly transitive on the points of \mathcal{P} . H a subgroup of G implies G is doubly transitive on the points of \mathcal{P} -again contradicting the fact that G has rank 3 on the points of \mathcal{P} . By Result IV either $n = 2$ and G has order 21, or $n = 8$ and G has order 657. But the second case cannot occur since $n > 2$ implies $|G|$ is even (Lemma 4). Thus $n = 2$ and G has order 21.

Conversely if $n = 2$, then \mathcal{P} is desarguesian, and the example at the end of §1 shows that in this case a rank 3 group does occur.

4. Proof of the main theorem. We now prove the main theorem stated in §1. Assume n is odd. If n is not a fourth power, then Theorem 1 and Result III implies that \mathcal{P} is desarguesian and G contains all elations of \mathcal{P} . But by Theorem 3 this is impossible.

Assume n is even and $n > 2$. Lemma 4 $|G|$ is even. If n is not a square, then an element in G of order 2 must be an elation (Baer [1]) and \mathcal{P} is desarguesian by Result V. This contradicts Theorem 3. Hence n is a square. If $n = m^2$ with $m \equiv 2 \pmod{4}$, then \mathcal{P} is desarguesian by Result VI since G is nonsolvable (Corollary 2.1). This contradicts Theorem 3 again. The proof of the main theorem is complete.

REFERENCES

1. R. Baer, *Projectivities with fixed points on every line of the plane*, Bull. Amer. Math. Soc., **52** (1946), 273-286.
2. P. Dembowski, *Verallgemeinerungen von Transitivitätsklassen endlicher projektiver Ebenen*, Math. Z., **69** (1958), 59-89.
3. ———, *Finite Geometries*, Springer-Verlag, Berlin-Heidelberg, 1968.
4. D. G. Higman and J. E. McLaughlin, *Geometric ABA-groups*, Ill. J. Math., **5** (1961), 382-397.

5. D. R. Hughes, *Collineations and generalized incidence matrices*, Trans. Amer. Math. Soc., **86** (1957), 284-296.
6. V. H. Keiser, *Finite projective planes with non-solvable transitive collineation groups*, Amer. Math. Monthly, **74** (1967), 556-559.
7. T. G. Ostrom, *Dual transitivity in finite projective planes*, Proc. Amer. Math. Soc., **9** (1958), 55-56.
8. T. G. Ostrom and A. Wagner, *On projective and affine planes with transitive collineation groups*, Math. Z., **71** (1959), 186-199.
9. E. T. Parker, *On collineations of symmetric designs*, Proc. Amer. Math. Soc., **8** (1957), 350-351.
10. A. Wagner, *On Perspectivities of finite projective planes*, Math. Z., **71** (1959), 113-123.
11. H. Wielandt, *Finite permutation groups*, Academic Press, New York and London, 1964.

Received January 12, 1971. This work was partially supported by NSF Contract No. GP. 17461.

WASHINGTON STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California 94305

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

C. R. HOBBY
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 39, No. 1

May, 1971

Charles A. Akemann, <i>A Gelfand representation theory for C^*-algebras</i>	1
Sorrell Berman, <i>Spectral theory for a first-order symmetric system of ordinary differential operators</i>	13
Robert L. Bernhardt, III, <i>On splitting in hereditary torsion theories</i>	31
J. L. Brenner, <i>Geršgorin theorems, regularity theorems, and bounds for determinants of partitioned matrices. II. Some determinantal identities</i>	39
Robert Morgan Brooks, <i>On representing F^*-algebras</i>	51
Lawrence Gerald Brown, <i>Extensions of topological groups</i>	71
Arnold Barry Calica, <i>Reversible homeomorphisms of the real line</i>	79
J. T. Chambers and Shinnosuke Oharu, <i>Semi-groups of local Lipschitzians in a Banach space</i>	89
Thomas J. Cheatham, <i>Finite dimensional torsion free rings</i>	113
Byron C. Drachman and David Paul Kraines, <i>A duality between transpotence elements and Massey products</i>	119
Richard D. Duncan, <i>Integral representation of excessive functions of a Markov process</i>	125
George A. Elliott, <i>An extension of some results of Takesaki in the reduction theory of von Neumann algebras</i>	145
Peter C. Fishburn and Joel Spencer, <i>Directed graphs as unions of partial orders</i>	149
Howard Edwin Gorman, <i>Zero divisors in differential rings</i>	163
Maurice Heins, <i>A note on the Löwner differential equations</i>	173
Louis Melvin Herman, <i>Semi-orthogonality in Rickart rings</i>	179
David Jacobson and Kenneth S. Williams, <i>On the solution of linear G.C.D. equations</i>	187
Michael Joseph Kallaher, <i>On rank 3 projective planes</i>	207
Donald Paul Minassian, <i>On solvable O^*-groups</i>	215
Nils Øvrelid, <i>Generators of the maximal ideals of $A(\bar{D})$</i>	219
Mohan S. Putcha and Julian Weissglass, <i>A semilattice decomposition into semigroups having at most one idempotent</i>	225
Robert Raphael, <i>Rings of quotients and π-regularity</i>	229
J. A. Siddiqi, <i>Infinite matrices summing every almost periodic sequence</i>	235
Raymond Earl Smithson, <i>Uniform convergence for multifunctions</i>	253
Thomas Paul Whaley, <i>Multiplicity type and congruence relations in universal algebras</i>	261
Roger Allen Wiegand, <i>Globalization theorems for locally finitely generated modules</i>	269